

GCSE AQA Biology

Includes Video Solutions & Digital Quiz

Revision Guide


Higher Level





laboratory tested

CGP's checklist for GCSE Biology success!



cgpbooks.co.uk

(CGP accepts no responsibility for books nibbled by rabbits.)



If you're looking for heart-pumping, neurone-jangling fun, then GCSE Biology is where it's at.

And with CGP, it really is fun — we're revision experts, so we leave no stone unturned

(and no joke untold). Before long, you'll be reproducing Biology facts in your sleep.

Looking for more? Well, wrap your retinas round the digital extras. We've included step-by-step video solutions from CGP's Biology boffins, quizzes to test your knowledge.

Yes we know, we're the best.

Revision Guide Higher Level

Contents

Working Scientifically	Topic 3 — Infection and Response		
The Scientific Method	1	Communicable Disease	46
Communication & Issues Created by Science	2	Viral, Fungal and Protist Diseases	
Risk	3	Bacterial Diseases and Preventing Disease	
Designing Investigations	4	Fighting Disease	
Collecting Data	5	Fighting Disease — Vaccination	
Processing and Presenting Data	6	Fighting Disease — Drugs	
Units and Equations	8	Developing Drugs	
Drawing Conclusions	9	Monoclonal Antibodies	
Uncertainties and Evaluations		More on Monoclonal Antibodies	
		Plant Diseases and Defences	
Topic 1 — Cell Biology		Revision Questions for Topic 3	
Cells	11	'	
Microscopy		Topic 4 — Bioenergetics	
More on Microscopy			E7
Cell Differentiation and Specialisation		Photosynthesis and Limiting Factors	
Chromosomes and Mitosis	15	The Rate of Photosynthesis	
Binary Fission	16	Respiration and Metabolism	
Culturing Microorganisms		Aerobic and Anaerobic Respiration	
Stem Cells		Exercise	
Diffusion		Revision Questions for Topic 4	64
Osmosis			
Active Transport		Topic 5 — Homeostasis and	
Exchange Surfaces		Response	
Exchanging Substances		Homeostasis	65
More on Exchanging Substances		The Nervous System	
Revision Questions for Topic 1		Synapses and Reflexes	
		Investigating Reaction Time	68
Topic 2 — Organisation		The Brain	69
	27	The Eye	
Cell Organisation		Correcting Vision Defects	
Enzymes	20	Controlling Body Temperature	
Investigating Enzymatic Reactions		The Endocrine System	
Enzymes and Digestion		Controlling Blood Glucose	
More on Enzymes and Digestion		The Kidneys	
Food Tests		Kidney Failure	
The Lungs		Puberty and the Menstrual Cycle	
Circulatory System — The Heart		Controlling Fertility	
Circulatory System — Blood Vessels		More on Controlling Fertility	
Circulatory System — Blood		Adrenaline and Thyroxine	
Cardiovascular Disease		Plant Hormones	
		Commercial Uses of Plant Hormones	
Health and Disease		Revision Questions for Topic 5	
Risk Factors for Non-Communicable Diseases		. to . topic 5	
Cancer			
Plant Cell Organisation			
Transpiration and Translocation			
Transpiration and Stomata			
Revision Questions for Topic 2	45		

Topic 6 — Inheritance, Variation and Evolution

DNA	84
The Structure of DNA and Protein Synthesis	85
Mutations	86
Reproduction	87
Meiosis	88
More on Reproduction	89
X and Y Chromosomes	90
Genetic Diagrams	91
More Genetic Diagrams	92
Inherited Disorders	93
The Work of Mendel	94
Variation	95
Evolution	96
More About Evolution	97
Selective Breeding	98
Genetic Engineering	99
Cloning	100
Fossils	101
Speciation	102
Antibiotic-Resistant Bacteria	103
Classification	104
Revision Questions for Topic 6	105

Topic 7 — Ecology

Competition	106
Abiotic and Biotic Factors	107
Adaptations	108
Food Chains	109
Using Quadrats	11C
Using Transects	111
Environmental Change & The Water Cycle	112
The Carbon Cycle	
Decay	
Investigating Decay	115
Biodiversity and Waste Management	116
Global Warming	117
Deforestation and Land Use	
Maintaining Ecosystems and Biodiversity	119
Trophic Levels	12C
Pyramids of Biomass	121
Biomass Transfer	122
Food Security and Farming	123
Biotechnology	124
Revision Questions for Topic 7	125
- '	
Practical Skills	
Measuring Substances	126
Heating, Safety and Ethics	
D 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	

Measuring Substances	126
Heating, Šafety and Ethics	127
otometers and Microscopes	128
5ampling	129
Comparing Results	
\nswers	131
n.	

You'll see QR codes throughout the book that you can scan with your smartphone.

A QR code next to a tip box question takes you to a video that talks you through solving the question. You can access all the videos by scanning this code here.



A QR code on a 'Revision Questions' page takes you to a Retrieval Quiz for that topic. You can access all the quizzes by scanning this code here.



You can also find the full set of videos at cgpbooks.co.uk/GCSEBiologyHigher/Videos and the full set of quizzes at cgpbooks.co.uk/GCSEBiologyHigher/Quiz

For useful information about What to Expect in the Exams and other exam tips head to cgpbooks.co.uk/GCSEBiologyHigher/Exams Published by CGP

From original material by Richard Parsons

Editors: Eleanor Crabtree, Mary Falkner, Sarah Pattison and Rachael Rogers.

Contributor: Paddy Gannon

BAR46 -1223 - 28376

With thanks to Lottie Edwards for the copyright research.

Clipart from Corel®

Illustrations by: Sandy Gardner Artist, email sandy@sandygardner.co.uk

Text, design, layout and original illustrations $\,$ © Coordination Group Publications Ltd (CGP) 2021 All rights reserved.

The Scientific Method

This section isn't about how to 'do' science — but it does show you the way most scientists work.

Scientists Come Up With Hypotheses — Then Test Them

- 1) Scientists try to explain things. They start by observing something they don't understand.
- 2) They then come up with a <u>hypothesis</u> a possible <u>explanation</u> for what they've observed.
- 3) The next step is to test whether the hypothesis might be right or not. This involves making a prediction based on the hypothesis and testing it by gathering evidence (i.e. data) from investigations. If evidence from experiments backs up a prediction, you're a step closer to figuring out if the hypothesis is true.



Hundreds of years ago, we thought demons caused illness.

Several Scientists Will Test a Hypothesis

- Normally, scientists share their findings in peer-reviewed journals, or at conferences.
- 2) Peer-review is where other scientists check results and scientific explanations to make sure they're 'scientific' (e.g. that experiments have been done in a sensible way) before they're published. It helps to detect false claims, but it doesn't mean that findings are correct just that they're not wrong in any obvious way.



3) Once other scientists have found out about a hypothesis, they'll start basing their own predictions on it and carry out their own experiments. They'll also try to reproduce the original experiments to check the results—and if all the experiments in the world back up the hypothesis, then scientists start to think the hypothesis is true.

4) However, if a scientist does an experiment that doesn't fit with the hypothesis (and other scientists can reproduce the results) then the hypothesis may need to be modified or scrapped altogether.

If All the Evidence Supports a Hypothesis, It's Accepted — For Now

- Accepted hypotheses are often referred to as theories. Our currently
 accepted theories are the ones that have survived this 'trial by evidence'
 they've been tested many times over the years and survived.
- However, theories never become totally indisputable <u>fact</u>. If <u>new evidence</u> comes along that <u>can't be explained</u> using the existing theory, then the hypothesising and testing is likely to <u>start all over again</u>.



Now we've collected more evidence, we know that illnesses that can be spread between people are due to microorganisms.

Theories Can Involve Different Types of Models

A representational model is a simplified description or picture of what's
going on in real life. Like all models, it can be used to explain observations
and make predictions. E.g. the lock and key model of enzyme action
is a simplified way of showing how enzymes work (see p.28). It can
be used to explain why enzymes only catalyse particular reactions.

Scientists test models by carrying out experiments to check that the predictions made by the model happen as expected.

- 2) Computational models use computers to make simulations of complex real-life processes, such as climate change. They're used when there are a lot of different variables (factors that change) to consider, and because you can easily change their design to take into account new data.
- 3) All models have limitations on what they can explain or predict. Climate change models have several limitations for example, it's hard to take into account all the biological and chemical processes that influence climate. It can also be difficult to include regional variations in climate.

I'm off to the zoo to test my hippo-thesis...

The scientific method has been developed over time, and many people have helped to develop it. From Aristotle to modern day scientists, lots of people have contributed. And many more are likely to contribute in the future.

Communication & Issues Created by Science

Scientific developments can be great, but they can sometimes raise more questions than they answer...

It's Important to Communicate Scientific Discoveries to the General Public

Some scientific discoveries show that people should change their habits, or they might provide ideas that could be developed into new technology. So scientists need to tell the world about their discoveries.

Gene technologies are used in genetic engineering to produce genetically modified crops. Information about these crops needs to be communicated to farmers who might benefit from growing them and to the general public, so they can make informed decisions about the food they buy and eat.

Scientific Evidence can be Presented in a Biased Way

- 1) Reports about scientific discoveries in the media (e.g. newspapers or television) aren't peer-reviewed.
- 2) This means that, even though news stories are often <u>based</u> on data that has been peer-reviewed, the data might be <u>presented</u> in a way that is <u>over-simplified</u> or <u>inaccurate</u>, making it open to <u>misinterpretation</u>.
- 3) People who want to make a point can sometimes <u>present data</u> in a <u>biased way</u>. (Sometimes <u>without knowing</u> they're doing it.) For example, a scientist might overemphasise a relationship in the data, or a newspaper article might describe details of data supporting an idea without giving any evidence against it.

Scientific Developments are Great, but they can Raise Issues

Scientific knowledge is increased by doing experiments. And this knowledge leads to scientific developments, e.g. new technologies or new advice. These developments can create issues though. For example:

Economic issues: Society can't always afford to do things scientists recommend (e.g. investing in alternative energy sources) without cutting back elsewhere.

Social issues: Decisions based on scientific evidence affect people — e.g. should alcohol be banned (to prevent health problems)? Would the effect on people's lifestyles be acceptable...?

Personal issues: Some decisions will affect individuals. For example, someone might support alternative energy, but object if a wind farm is built next to their house.

Environmental issues: Human activity often affects the natural environment — e.g. genetically modified crops may help us to produce more food — but some people think they could cause environmental problems (see p.99).

THE GAZETTE THE

Science Can't Answer Every Question — Especially Ethical Ones

- 1) We don't understand everything. We're always finding out more, but we'll never know all the answers.
- 2) In order to answer scientific questions, scientists need data to provide evidence for their hypotheses.
- Some questions can't be answered <u>yet</u> because the data <u>can't</u> currently be <u>collected</u>, or because there's not enough data to support a theory.
- 4) Eventually, as we get more evidence, we'll answer some of the questions that <u>currently</u> can't be answered, e.g. what the impact of global warming on sea levels will be. But there will always be the "Should we be doing this at all?"-type questions that experiments <u>can't</u> help us to answer...

Think about new drugs which can be taken to boost your 'brain power'.

- Some people think they're good as they could improve concentration or memory.
 New drugs could let people think in ways beyond the powers of normal brains.
- Other people say they're <u>bad</u> they could give you an <u>unfair advantage</u> in exams. And people
 might be <u>pressured</u> into taking them so that they could work more <u>effectively</u>, and for <u>longer hours</u>.

Tea to milk or milk to tea? — Totally unanswerable by science...

Science can't tell you whether or not you should do something. That's for you and society to decide. But there are tons of questions science might be able to answer, like where life came from and where my superhero socks are.

Risk

By reading this page you are agreeing to the risk of a paper cut or severe drowsiness...

Nothing is Completely Risk-Free

- 1) A hazard is something that could potentially cause harm.
- 2) All hazards have a risk attached to them this is the chance that the hazard will cause harm.
- 3) The risks of some things seem pretty <u>obvious</u>, or we've known about them for a while, like the risk of causing <u>acid rain</u> by polluting the atmosphere, or of having a <u>car accident</u> when you're travelling in a car.
- 4) New technology arising from scientific advances can bring new risks, e.g. scientists are unsure whether nanoparticles that are being used in cosmetics and suncream might be harming the cells in our bodies. These risks need to be considered alongside the benefits of the technology, e.g. improved sun protection.
- 5) You can estimate the <u>size</u> of a risk based on <u>how many times</u> something happens in a big sample (e.g. 100 000 people) over a given <u>period</u> (e.g. a year). For example, you could assess the risk of a driver crashing by recording how many people in a group of 100 000 drivers crashed their cars over a year.
- 6) To make <u>decisions</u> about activities that involve <u>hazards</u>, we need to take into account the <u>chance</u> of the hazard causing harm, and how <u>serious</u> the <u>consequences</u> would be if it did. If an activity involves a hazard that's <u>very likely</u> to cause harm, with <u>serious consequences</u> if it does, it's considered <u>high risk</u>.

People Make Their Own Decisions About Risk

- 1) Not all risks have the same <u>consequences</u>, e.g. if you chop veg with a sharp knife you risk cutting your finger, but if you go scuba-diving you risk death. You're much <u>more likely</u> to cut your finger during half an hour of <u>chopping</u> than to die during half an hour of <u>scuba-diving</u>. But most people are happier to accept a higher <u>probability</u> of an accident if the <u>consequences</u> are <u>short-lived</u> and fairly <u>minor</u>.
- 2) People tend to be more willing to accept a risk if they <u>choose</u> to do something (e.g. go scuba diving), compared to having the risk <u>imposed</u> on them (e.g. having a nuclear power station built next door).
- 3) People's <u>perception</u> of risk (how risky they <u>think</u> something is) isn't always <u>accurate</u>. They tend to view <u>familiar</u> activities as <u>low-risk</u> and <u>unfamiliar</u> activities as <u>high-risk</u>. even if that's not the case. For example, cycling on roads is often <u>high-risk</u>, but many people are happy to do it because it's a <u>familiar</u> activity. Air travel is actually pretty <u>safe</u>, but a lot of people perceive it as <u>high-risk</u>.
- 4) People may over-estimate the risk of things with long-term or invisible effects, e.g. ionising radiation.

Investigations Can be Hazardous

- 1) Hazards from science experiments might include:
 - · Microorganisms, e.g. some bacteria can make you ill.
 - · Chemicals, e.g. sulfuric acid can burn your skin and alcohols catch fire easily.
 - · Fire, e.g. an unattended Bunsen burner is a fire hazard.
 - Electricity, e.g. faulty electrical equipment could give you a shock.
- 2) Part of planning an investigation is making sure that it's safe.
- 3) You should always make sure that you identify all the hazards that you might encounter. Then you should think of ways of reducing the risks from the hazards you've identified. For example:
 - If you're working with <u>sulfuric acid</u>, always wear gloves and safety goggles. This will reduce
 the risk of the acid coming into contact with your skin and eyes.
 - If you're using a <u>Bunsen burner</u>, stand it on a heat proof mat.
 This will reduce the risk of starting a fire.

You can find out about potential hazards by looking in textbooks, doing some internet research, or asking your teacher.

Not revising — an unacceptable exam hazard...

The world's a dangerous place, but if you can recognise hazards, decide how to reduce their risks, and be happy to accept some risks, you can still have fun. Just maybe don't go skydiving with a great white shark on Friday 13th.

Hmm... Where did my bacteria sample go?

Designing Investigations

Dig out your lab coat and dust down your badly-scratched safety goggles... it's investigation time.

Investigations Produce Evidence to Support or Disprove a Hypothesis

Scientists observe things and come up with hypotheses to explain them (see p.1).
 You need to be able to do the same. For example:

Observation: People have big feet and spots. Hupothesis: Having big feet causes spots.

- 2) To <u>determine</u> whether or not a hypothesis is <u>right</u>, you need to do an <u>investigation</u> to gather evidence. To do this, you need to use your hypothesis to make a <u>prediction</u> something you think <u>will happen</u> that you can test. E.g. people who have bigger feet will have more spots.
- 3) Investigations are used to see if there are <u>patterns</u> or <u>relationships</u> between <u>two variables</u>, e.g. to see if there's a pattern or relationship between the variables 'number of spots' and 'size of feet'.

Evidence Needs to be Repeatable, Reproducible and Valid

- 1) Repeatable means that if the <u>same person</u> does an experiment again using the <u>same methods</u> and equipment, they'll get <u>similar results</u>.
- 2) Reproducible means that if someone else does the experiment, or a different method or piece of equipment is used, the results will still be similar.



Investigations

include experiments =

4) <u>Valid results</u> are both repeatable and reproducible AND they <u>answer the original question</u>. They come from experiments that were designed to be a FAIR TEST...

To Make an Investigation a Fair Test You Have to Control the Variables

- 1) In a lab experiment you usually change one variable and measure how it affects another variable.
- To make it a fair test, everything else that could affect the results should stay the same
 — otherwise you can't tell if the thing you're changing is causing the results or not.
- 3) The variable you CHANGE is called the INDEPENDENT variable.
- 4) The variable you MEASURE when you change the independent variable is the DEPENDENT variable.
- 5) The variables that you KEEP THE SAME are called CONTROL variables.

You could find how temperature affects the rate of an enzyme-controlled reaction. The independent variable is the temperature. The dependent variable is the rate of reaction. Control variables include the concentration and amounts of reactants, pH, the time period you measure, etc.

6) Because you can't always control all the variables, you often need to use a <u>control experiment</u>. This is an experiment that's kept under the <u>same conditions</u> as the rest of the investigation, but <u>doesn't</u> have anything <u>done</u> to it. This is so that you can see what happens when you don't change anything at all.

The Bigger the Sample Size the Better

- 1) Data based on <u>small samples</u> isn't as good as data based on large samples. A sample should <u>represent</u> the <u>whole population</u> (i.e. it should share as many of the characteristics in the population as possible) a small sample can't do that as well. It's also harder to spot <u>anomalies</u> if your sample size is too small.
- 2) The <u>bigger</u> the sample size the <u>better</u>, but scientists have to be <u>realistic</u> when choosing how big. For example, if you were studying how lifestyle affects people's weight it'd be great to study everyone in the UK (a huge sample), but it'd take ages and cost a bomb. It's more realistic to study a thousand people, with a mixture of ages, gender and race.

This is no high street survey — it's a designer investigation...

Not only do you need to be able to plan your own investigations, you should also be able to look at someone else's plan and decide whether or not it needs improving. Those examiners aren't half demanding.

Collecting Data

You've designed the perfect investigation - now it's time to get your hands mucky and collect some data.

Your Data Should be Repeatable, Reproducible, Accurate and Precise

- To check repeatability you need to repeat the readings and check that the results are similar.
 You need to repeat each reading at least three times.
- To make sure your results are <u>reproducible</u> you can cross check them by taking a second set of readings with another instrument (or a different observer).
- 3) Your data also needs to be ACCURATE. Really accurate results are those that are really close to the true answer. The accuracy of your results usually depends on your method you need to make sure you're measuring the right thing and that you don't miss anything that should be included in the measurements. E.g. estimating the amount of gas released from a reaction by counting the bubbles isn't very accurate because you might miss some of the bubbles and they might have different yolumes. It's more accurate to measure the volume of gas released using a gas suringe.



Beth's result

was a curate.

Data set 1 is more precise
... than data set 2.

If there's no systematic error, \subseteq

then doing repeats and

 Your data also needs to be PRECISE. Precise results are ones where the data is all really close to the mean (average) of your repeated results (i.e. not spread out).

Your Equipment has to be Right for the Job

- 1) The measuring equipment you use has to be <u>sensitive enough</u> to measure the changes you're looking for. For example, if you need to measure changes of 1 cm³ you need to use a measuring cylinder that can measure in 1 cm³ steps it'd be no good truing with one that only measures in 10 cm³ steps.
- 2) The <u>smallest change</u> a measuring instrument can <u>detect</u> is called its <u>RESOLUTION</u>. E.g. some mass balances have a resolution of 1 g, some have a resolution of 0.1 g, and some are even more sensitive.
- 3) Also, equipment needs to be <u>calibrated</u> by measuring a known value. If there's a <u>difference</u> between the <u>measured</u> and <u>known value</u>, you can use this to <u>correct</u> the inaccuracy of the equipment.

You Need to Look out for Errors and Anomalous Results

- 1) The results of your experiment will always vary a bit because of RANDOM ERRORS unpredictable differences caused by things like human errors in measuring. E.g. the errors you make when reading from a measuring cylinder are random. You have to estimate or round the level when it's between two marks so sometimes your figure will be a bit above the real one, and sometimes it will be a bit below.
- You can reduce the effect of random errors by taking repeat readings and finding the mean. This will make your results more precise.
- 3) If a measurement is wrong by the same amount every time, it's called a SYSTEMATIC ERROR. For example, if you measured from the very end of your ruler instead of from the O cm mark every time, all your measurements would be a bit small. Repeating the experiment in the exact same way and calculating a mean won't correct a systematic error.
- 4) Just to make things more complicated, if a systematic error is caused by using equipment that isn't zeroed properly, it's called a ZERO ERROR. For example, if a mass balance always reads 1 gram before you put anything on it, all your measurements will be 1 gram too heavy.
- 5) You can compensate for some systematic errors if you know about them though, e.g. if your mass balance always reads I gram before you put anything on it you can subtract I gram from all your results.
- 6) Sometimes you get a result that doesn't fit in with the rest at all. This is called an ANOMALOUS RESULT. You should investigate it and try to work out what happened. If you can work out what happened (e.g. you measured something totally wrong) you can ignore it when processing your results.

Watch what you say to that mass balance — it's very sensitive...

Weirdly, data can be really precise but not very accurate. For example, a fancy piece of lab equipment might give results that are really precise, but if it's not been calibrated properly those results won't be accurate.

Processing and Presenting Data

Processing your data means doing some <u>calculations</u> with it to make it <u>more useful</u>. Once you've done that, you can present your results in a nice <u>chart</u> or <u>graph</u> to help you <u>spot any patterns</u> in your data.

Data Needs to be Organised

Tables are dead useful for <u>organising data</u>. When you draw a table <u>use a ruler</u> and make sure each column has a heading (including the units).

You Might Have to Process Your Data

- When you've done repeats of an experiment you should always calculate the <u>mean</u> (a type of average).
 To do this <u>add together</u> all the data values and <u>divide</u> by the total number of values in the sample.
- You might also need to calculate the <u>range</u> (how spread out the data is).
 To do this find the largest number and subtract the smallest number from it.

if. results when calculating these.

EXAMPLE

The results of an experiment to find the volume of gas produced in an enzyme-controlled reaction are shown below. Calculate the mean volume and the range.

	Repeat 1 (cm³)	Repeat 2 (cm³)	Repeat 3 (cm³)	Mean (cm³)	Range (cm³)
1	28	37	32	$(28 + 37 + 32) \div 3 = 32$	37 – 28 = 9

3) You might also need to calculate the <u>median</u> or <u>mode</u> (two more types of average). To calculate the <u>median</u>, put all your data in <u>numerical order</u> — the median is the middle value. The number that appears most often in a data set is the mode.

E.g. If you have the data set: 1 2 1 1 3 4 2

The median is: 1 1 1 2 2 3 4. The mode is 1 because 1 appears most often.

If you have an even of number of values, the median is halfway between the middle two values.

Round to the Lowest Number of Significant Figures

The <u>first significant figure</u> of a number is the first digit that's <u>not zero</u>. The second and third significant figures come <u>straight after</u> (even if they're zeros). You should be aware of significant figures in calculations.

- 1) In any calculation, you should round the answer to the lowest number of significant figures (s.f.) given.
- 2) Remember to write down how many significant figures you've rounded to after your answer.
- 3) If your calculation has multiple steps, only round the final answer, or it won't be as accurate.

EXAMPLE


A plant produces 10.2 cm³ of oxygen in 6.5 minutes whilst photosynthesising. Calculate the rate of photosynthesis.

rate = 10.2 cm³ ÷ 6.5 min = 1.5692... = 1.6 cm³/min (2 s.f.)

Final answer should be rounded to 2 s.f.


If Your Data Comes in Categories, Present It in a Bar Chart

- If the independent variable is <u>categoric</u> (comes in distinct categories, e.g. flower colour, blood group) you should use a <u>bar chart</u> to display the data.
- 2) You also use them if the independent variable is discrete (the data can be counted in chunks, where there's no in-between value, e.g. number of bacteria is discrete because you can't have half a bacterium).
- 3) There are some golden rules you need to follow for drawing bar charts:



If Your Data is Continuous, Plot a Graph

If both variables are continuous (numerical data that can have any value within a range, e.g. length, volume, temperature) you should use a graph to display the data.



Graphs Can Give You a Lot of Information About Your Data

The gradient (slope) of a graph tells you how quickly the dependent variable changes if you change the independent variable.

change in y gradient = change in X

munnimuni kananani You can use this method to calculate any rates from a graph, not just the rate of a reaction. Just remember that a rate is how much something changes over time, so x needs to be the time.

This graph shows the volume of gas produced in a reaction against time. The graph is linear (it's a straight line graph), so you can simply calculate the gradient of the line to find out the rate of reaction.

1) To calculate the gradient, pick two points on the line that are easy to read and a good distance apart.

2) Draw a line down from one of the points and a line across from the other to make a triangle. The line drawn down the side of the triangle is the change in y and the line across the bottom is the change in x.

Change in $y = 6.8 - 2.0 = 4.8 \text{ cm}^3$ Change in x = 5.2 - 1.6 = 3.6 schange in $y = \frac{4.8 \text{ cm}^3}{1.3 \text{ cm}^3/\text{s}} = 1.3 \text{ cm}^3/\text{s}$ or $1.3 \text{ cm}^3\text{s}^{-1}$

If you've got a curved graph, you can find the rate at any point by drawing a tangent — a straight line that touches a single point on a curve. You can then find the gradient of the tangent in the usual way, to give you the rate at that point. The units of the gradient are (units of y)/(units of x). cm3/s can also be written as cm3s-1.

Time (s)

8 (cm³)

6-

4 t

2 /olume

2) The intercept of a graph is where the line of best fit crosses one of the axes. The x-intercept is where the line of best fit crosses the x-axis and the <u>y-intercept</u> is where it crosses the <u>y-axis</u>.

Graphs Show the Relationship Between Two Variables

1) You can get three types of correlation (relationship) between variables:

Rate = gradient =

2) Just because there's correlation. it doesn't mean the change in one variable is causing the change in the other - there might be other factors involved (see page 9).





relationship between the two variables.

I love eating apples — I call it core elation...

Science is all about finding relationships between things. And I don't mean that chemists gather together in corners to discuss whether or not Devini and Sebastian might be a couple... though they probably do that too.

Units and Equations

Graphs and maths skills are all very well, but the numbers don't mean much if you can't get the units right.

SI Units Are Used All Round the World

- It wouldn't be all that useful if I defined volume in terms of <u>bath tubs</u>, you defined it in terms of <u>egg-cups</u> and my pal Sarwat defined it in terms of <u>balloons</u> — we'd never be able to compare our data.
- To stop this happening, scientists have come up with a set of standard units, called SI units, that all scientists use to measure their data. Here are some SI units you'll see in Biology:

Quantity	SI Base Unit
mass	kilogram, kg
length	metre, m
time	second, s

Scaling Prefixes Can Be Used for Large and Small Quantities

- Quantities come in a huge range of sizes. For example, the volume of a swimming pool might be around 2 000 000 000 cm³, while the volume of a cup is around 250 cm³.
- 2) To make the size of numbers more <u>manageable</u>, larger or smaller units are used. These are the <u>SI base unit</u> (e.g. metres) with a <u>prefix</u> in front:

prefix	tera (T)	giga (G)	mega (M)	kilo (k)	deci (d)	centi (c)	milli (m)	micro (μ)	nano (n)
multiple of unit	1012	10 ⁹	1 000 000 (106)	1000	0.1	0.01	0.001	0.000001 (10-6)	10-9

- These <u>prefixes</u> tell you <u>how much bigger</u> or <u>smaller</u> a unit is than the base unit. So one <u>kilometre</u> is <u>one thousand</u> metres.
- The conversion factor is the number of times the smaller unit goes into the larger unit.
- To swap from one unit to another, all you need to know is what number you have to divide or multiply by to get from the original unit to the new unit — this is called the conversion factor.
 - To go from a bigger unit (like m) to a smaller unit (like cm), you multiply by the conversion factor.
 - To go from a smaller unit (like g) to a bigger unit (like kg), you divide by the conversion factor.
- 5) Here are some conversions that'll be useful for GCSE biology:

Mass can have units of kg and g.

× 1000 kg g Length can have lots of units, including mm, μ m and nm.



Time can have units of min and s.



Volume can have units of m³, dm³ and cm³.

Always Check The Values Used in Equations Have the Right Units

- Equations (sometimes called formulas) show relationships between variables.
- To rearrange an equation, make sure that whatever you do to one side of the equation you also do to the other side.
- You can find the <u>magnification</u> of something using the equation: magnification = image size ÷ real size (see p.12). You can rearrange this equation to find the <u>image size</u> by <u>multiplying</u> each side by the real size: image size = magnification × real size.
- 3) To use an equation, you need to know the values of <u>all but one</u> of the variables. <u>Substitute</u> the values you do know into the equation, and do the calculation to work out the final variable.
- 4) Always make sure the values you put into an equation have the <u>right units</u>. For example, if you're calculating the magnification of something, but your image size is in mm and the real size is in μm, you'll have to <u>convert</u> both measurements into the same unit (either mm or μm) before you start.
- 5) To make sure your units are correct, it can help to write down the units on each line of your calculation.

I wasn't sure I liked units, but now I'm converted...

It's easy to get in a muddle when converting between units, but there's a handy way to check you've done it right. If you're moving from a smaller unit to a larger unit (e.g. g to kg) the number should get smaller, and vice versa.

Drawing Conclusions

Congratulations — uou're nearly at the end of a gruelling investigation, time to draw conclusions.

You Can Only Conclude What the Data Shows and NO MORE

1) Drawing conclusions might seem pretty straightforward — you just look at your data and say what pattern or relationship you see between the dependent and independent variables.

The table on the right shows the heights of pea plant seedlings grown for three weeks with different fertilisers.

Fertiliser	Mean growth / mm
Α	13.5
В	19.5
No fertiliser	5.5

CONCLUSION: Fertiliser B makes pea plant seedlings grow taller over a three week period than fertiliser A.

2) But you've got to be really careful that your conclusion matches the data you've got and doesn't go any further. You can't conclude that fertiliser B makes any other type of plant grow taller than fertiliser A the results could be totally different.

3) You also need to be able to use your results to justify your conclusion (i.e. back up

Over the three week period, fertiliser B made the pea your conclusion with some specific data). - plants grow 6 mm more on average than fertiliser A.

4) When writing a conclusion you need to refer back to the original hypothesis and say whether the data supports it or not: _____ The hupothesis for this experiment might have been that adding fertiliser would increase the growth of plants and that different types of fertiliser would affect growth by different amounts. If so, the data supports the hypothesis.

Correlation DOES NOT Mean Cause

If two things are correlated (i.e. there's a relationship between them) it doesn't necessarily mean a change in one variable is causing the change in the other — this is REALLY IMPORTANT — DON'T FORGET IT. There are three possible reasons for a correlation:

1) CHANCE: It might seem strange, but two things can show a correlation purely due to chance.

For example, one study might find a correlation between people's hair colour and how good they are at frisbee. But other scientists don't get a correlation when they investigate it — the results of the first study are just a fluke.

2) LINKED BY A 3RD VARIABLE: A lot of the time it may look as if a change in one variable is causing a change in the other, but it isn't — a third variable links the two things.

For example, there's a correlation between water temperature and shark attacks. This isn't because warmer water makes sharks crazy. Instead, they're linked by a third variable — the number of people swimming (more people swim when the water's hotter, and with more people in the water you get more shark attacks).



3) CAUSE: Sometimes a change in one variable does cause a change in the other. You can only conclude that a correlation is due to cause when you've controlled all the variables that could, just could, be affecting the result.

For example, there's a correlation between smoking and lung cancer. This is because chemicals in tobacco smoke cause lung cancer. This conclusion was only made once other variables (such as age and exposure to other things that cause cancer) had been controlled and shown not to affect people's risk of getting lung cancer.



I conclude that this page is a bit dull...

...although, just because I find it dull doesn't mean that I can conclude it's dull (you might think it's the most interesting thing since that kid got his head stuck in the railings near school). In the exams you could be given a conclusion and asked whether some data supports it - so make sure you understand how far conclusions can go.

Uncertainties and Evaluations

Hurrah! The end of another investigation. Well, now you have to work out all the things you did wrong.

Uncertainty is the Amount of Error Your Measurements Might Have

- When you repeat a measurement, you often get a <u>slightly different</u> figure each time you do it due to random error. This means that each result has some uncertainty to it.
- The measurements you make will also have some uncertainty in them due to limits in the resolution of the equipment you use (see page 5).

 The description of the equipment you use (see page 5).
- 3) This all means that the mean of a set of results will also have some uncertainty to it. You can calculate the uncertainty of a mean result using the equation:
- 4) The larger the range, the less precise your results are and the more uncertainty there will be in your results. Uncertainties are shown using the '±' symbol.

The range is the largest value minus the smallest value (p.6).



EXAMPLE

The table below shows the results of a respiration experiment to determine the volume of carbon dioxide produced. Calculate the uncertainty of the mean.

Repeat	1	2	3	mean
Volume of CO ₂ produced (cm ³)	20.1	19.8	20.0	20.0

1) First work out the range: Range = 20.1 - 19.8 = 0.300 cm³

Use the range to find the uncertainty:


Uncertainty = range \div 2 = 0.300 \div 2 = 0.150 cm³. So the uncertainty of the mean = 20.0 \pm 0.150 cm³

5) Measuring a greater amount of something helps to reduce uncertainty. For example, in a rate of reaction experiment, measuring the amount of product formed over a longer period compared to a shorter period will reduce the percentage uncertainty in your results.

Evaluations — Describe How it Could be Improved

An evaluation is a critical analysis of the whole investigation.

- You should comment on the <u>method</u> was it <u>valid</u>?
 Did you control all the other variables to make it a <u>fair test</u>?
- Comment on the quality of the results was there enough evidence to reach a valid conclusion? Were the results repeatable, reproducible, accurate and precise?



- 4) All this analysis will allow you to say how confident you are that your conclusion is right.
- 5) Then you can suggest any changes to the method that would improve the quality of the results, so that you could have more confidence in your conclusion. For example, you might suggest changing the way you controlled a variable, or increasing the number of measurements you took. Taking more measurements at parrower intervals could give you a more accurate result. For example:

Enzymes have an optimum temperature (a temperature at which they work best). Say you do an experiment to find an enzyme's optimum temperature and take measurements at 10 °C, 20 °C, 30 °C, 40 °C and 50 °C. The results of this experiment tell you the optimum is 40 °C. You could then repeat the experiment, taking more measurements around 40 °C to a get a more accurate value for the optimum.

 You could also make more <u>predictions</u> based on your conclusion, then <u>further experiments</u> could be carried out to test them.

When suggesting improvements to the investigation, always make sure that you say why you think this would make the results better.

Evaluation — next time, I'll make sure I don't burn the lab down...

So there you have it — Working Scientifically. Make sure you know this stuff like the back of your hand. It's not just in the lab that you'll need to know how to work scientifically. You can be asked about it in the exams as well.



Cells

When someone first peered down a microscope at a slice of cork and drew the boxes theu saw. little did they know that they'd seen the building blocks of every organism on the planet...

Organisms can be Prokarvotes or Eukarvotes

- 1) All living things are made of cells.
- 2) Cells can be either prokaryotic or eukaryotic. Eukaryotic cells are complex and include all animal and plant cells. Prokaryotic cells are smaller and simpler, e.g. bacteria (see below).
- 3) Eukaryotes are organisms that are made up of eukaryotic cells.
- 4) A prokaruote is a prokaruotic cell (it's a single-celled organism).

Plant and Animal Cells have Similarities and Differences

The different parts of a cell are called subcellular structures.

Most animal cells have the following subcellular structures — make sure you know them all:




- Nucleus contains genetic material that controls the activities of the cell.
 - Cutoplasm gel-like substance where most of the chemical reactions happen. It contains enzumes (see page 28) that control these chemical reactions.
 - 3) Cell membrane holds the cell together and controls what goes in and out.
 - Mitochondria these are where most of the reactions for aerobic respiration take place (see page 62). Respiration transfers energy that the cell needs to work.
 - Ribosomes these are where proteins are made in the cell.

Plant cells usually have all the bits that animal cells have, plus a few extra things that animal cells don't have:

- 1) Rigid <u>cell wall</u> made of <u>cellulose</u>. It <u>supports</u> the cell and strengthens it.
- 2) Permanent vacuole contains cell sap, a weak solution of sugar and salts.
- 3) Chloroplasts these are where photosynthesis occurs, which makes food for the plant (see page 57). They contain a green substance called chlorophull, which absorbs the light needed for photosynthesis.






Bacterial Cells Are Much Smaller

Bacteria are prokaryotes. Here's what a bacterial cell might look like:

MINITED THE PROPERTY OF THE PR You might see the sizes of cells written in standard form (see the next page).



Cell structures — become an estate agent...

You could get asked to estimate the area of a subcellular structure in your exam. If you do, treat it as a regular shape. For example, if it's close to a rectangle, use the area formula 'area = length × width'.

Give two differences in structure between prokaryotic and eukaryotic cells. 01

[2 marks]

Microscopy

Microscopes are pretty important for biology. So here's a couple of pages all about them...

Cells are Studied Using Microscopes

- Microscopes let us see things that we can't see with the naked eye. The microscopy techniques
 we can use have developed over the years as technology and knowledge have improved.
- Light microscopes use light and lenses to form an image of a specimen and magnify it (make it look bigger). They let us see individual cells and large subcellular structures, like nuclei.
- 3) <u>Electron microscopes</u> use <u>electrons</u> instead of light to form an image. They have a much <u>higher magnification</u> than light microscopes.
- 4) They also have a <u>higher resolution</u>. (Resolution is the ability to <u>distinguish</u> between two points, so a higher resolution gives a sharper image.)
- 5) Electron microscopes let us see much <u>smaller things</u> in <u>more detail</u>, like the <u>internal structure</u> of <u>mitochondria</u> and <u>chloroplasts</u>. They even let us see <u>tinier</u> things like <u>ribosomes</u> and <u>plasmids</u>.

See the next page = for how to use a = light microscope.

You Need to be Able to Use the Formula for Magnification

You can calculate the magnification of an image using this formula:

$$magnification = \frac{image\ size}{real\ size}$$

Image size and real size should have the same units. If they don't, you'll need to convert them first (see page 8). If you want to work out the <u>image size</u> or the <u>real size</u> of the object, you can rearrange the equation using this <u>formula triangle</u>:



Cover up the thing you're trying to find. The parts you can still see are the formula you need to use.

EXAMPLE

A specimen is 50 µm wide. Calculate the width of the image of the specimen under a magnification of × 100. Give your answer in mm.

- 1) Rearrange the formula.
- 2) Fill in the values you know.
- 3) Remember the units in your answer.
- 4) Convert the units.

image size = magnification \times real size image size = 100×50

= 5000 μm = 5 mm Remember, to convert from micrometres (µm) to millimetres (mm), you need to divide by 1000 (see p.8).
E.g. 5000 µm ÷ 1000 = 5 mm

You Need to Know How to Work With Numbers in Standard Form

- 1) Because microscopes can see such tinu objects, sometimes it's useful to write numbers in standard form.
- This is where you change very big or small numbers with lots of zeros into something more manageable, e.g. 0.017 can be written 1.7 × 10⁻².
- To do this you just need to <u>move</u> the <u>decimal point</u> left or right.
- 4) The number of places the decimal point moves is then represented by a <u>power of 10</u> this is <u>positive</u> if the decimal point's moved to the <u>left</u>, and <u>negative</u> if it's moved to the <u>right</u>.

EXAMPLE

A mitochondrion is approximately 0.0025 mm long. Write this figure in standard form.

 The first number needs to be <u>between 1 and 10</u> so the decimal point needs to move after the '2'.

0.0025

Count how many places the decimal point has moved

 this is the power of 1O. Don't forget the minus sign because the decimal point has moved right.

2.5 × 10⁻³

Your resolution to revise should be increasing right now...

Keep an eye on the units for that equation — if they're not the same, it just won't work.

Q1 A cheek cell is viewed under a microscope with × 40 magnification. The image of the cell is 2.4 mm wide. Calculate the real width of the cheek cell. Give your answer in µm. [2 marks]



More on Microscopy



Water drop mixed

with indine

Specimen stained with

iodine

Cover slip

It's all very well knowing what microscopes do — you also have to know how to actually use one.

You Need to Prepare Your Slide


If you want to look at a specimen (e.g. plant or animal cells) under a light microscope, you need to put it on a <u>microscope slide</u> first. A slide is a strip of clear <u>glass</u> or <u>plastic</u> onto which the specimen is <u>mounted</u>. Here's how to prepare a slide to view onion cells:

- 1) Add a drop of water to the middle of a clean slide.
- Cut up an onion and separate it out into <u>layers</u>. Use <u>tweezers</u> to peel off some <u>epidermal tissue</u> from the bottom of one of the layers.
- 3) Using the tweezers, place the epidermal tissue into the water on the slide.
- Add a drop of <u>iodine solution</u>. Iodine solution is a <u>stain</u>.
 Stains are used to highlight objects in a cell by adding <u>colour</u> to them.
- 5) Place a cover slip (a square of thin, transparent plastic or glass) on top. To do this, stand the cover slip upright on the slide, next to the water droplet. Then carefully tilt and lower it so it covers the specimen. Try not to get any air bubbles under there they'll obstruct your view of the specimen.

Use a Light Microscope to Look at Your Slide


To look at your prepared slides, you need to know how to use a light microscope:

- 1) Clip the slide you've prepared onto the stage.
- 2) Select the <u>lowest-powered objective lens</u>
 (i.e. the one that produces the lowest magnification).
- Use the coarse adjustment knob to move the stage up to just below the objective lens.
- Look down the <u>eyepiece</u>. Use the coarse adjustment knob to move the stage downwards until the image is <u>roughly in focus</u>.
- 5) Adjust the <u>focus</u> with the <u>fine adjustment knob</u>, until you get a <u>clear image</u> of what's on the slide.
- 6) If you need to see the slide with greater magnification, swap to a higher-powered objective lens and refocus.



Draw Your Observations Neatly with a Pencil

- 1) Draw what you see under the microscope using a pencil with a sharp point.
- Make sure your drawing takes up at least half of the space available and that it is drawn with clear, unbroken lines.
- 3) Your drawing should not include any colouring or shading.
- If you are drawing <u>cells</u>, the <u>subcellular structures</u> should be drawn in <u>proportion</u>.
- Remember to include a <u>title</u> of what you were observing and write down the <u>magnification</u> that it was observed under.
- 6) Label the important features of your drawing (e.g. nucleus, chloroplasts), using straight, uncrossed lines.



You can work out the real size of a cell by counting the number of cells in 1 mm of the sample (see p128). You can work out the magnification of your drawing using this formula-magnification = length of drawing of cell + real length of cell.

So here, magnification = 33 mm + O3 mm = x10.

A light microscope is better than a heavy one...

If you can use a microscope, you're halfway to ruling the world. That's what I like to think, anyway.

O1 Why might you add stain to the sample on a microscope slide?


[1 mark]

Cell Differentiation and Specialisation

Cells don't all look the same. They have different structures to suit their different functions.

Cells Differentiate to Become Specialised

- 1) Differentiation is the process by which a cell changes to become specialised for its job.
- As cells change, they develop <u>different subcellular structures</u> and turn into <u>different types of cells</u>.
 This allows them to carry out specific functions.
- 3) Most differentiation occurs as an organism develops. In most animal cells, the ability to differentiate is then lost at an early stage, after they become specialised. However, lots of plant cells don't ever lose this ability.
- 4) The cells that differentiate in mature animals are mainly used for repairing and replacing cells, such as skin or blood cells.



5) Some cells are undifferentiated cells — they're called stem cells. There's more about them on page 19.

You Need To Know These Examples of Specialised Cells

SPERM CELLS are specialised for REPRODUCTION

The function of a <u>sperm</u> is basically to get the <u>male DNA</u> to the <u>female DNA</u>. It has a <u>long tail</u> and a <u>streamlined head</u> to help it <u>swim</u> to the <u>egg</u>.

There are a lot of <u>mitochondria</u> in the cell to provide the <u>energy</u> needed. It also carries <u>enzymes</u> in its head to digest through the <u>egg</u> cell membrane.

NERVE CELLS are specialised for RAPID SIGNALLING

The function of <u>nerve cells</u> is to <u>carry electrical signals</u> from one part of the body to another. These cells are <u>long</u> (to cover more distance) and have <u>branched connections</u> at their ends to <u>connect</u> to other nerve cells and form a <u>network</u> throughout the body.

MUSCLE CELLS are specialised for CONTRACTION

The function of a <u>muscle cell</u> is to contract quickly. These cells are <u>long</u> (so that they have space to <u>contract</u>) and contain <u>lots</u> of <u>mitochondria</u> to generate the <u>energy</u> needed for contraction.

ROOT HAIR CELLS are specialised for absorbing WATER and MINERALS

Root hair cells are cells on the surface of plant roots, which grow into long "hairs" that stick out into the soil. This gives the plant a big surface area for absorbing water and mineral ions from the soil.

PHLOEM and XYLEM CELLS are specialised for TRANSPORTING SUBSTANCES

Phloem and xylem cells form phloem and xylem tubes, which transport substances such as <u>food</u> and <u>water</u> around plants. To form the tubes, the cells are long and joined <u>end to end</u>. Xylem cells are <u>hollow</u> in the centre and phloem cells have <u>very few subcellular structures</u>, so that stuff can <u>flow through</u> them.



Tadpoles and tent pegs — cells are masters of disguise...

You need to know how the structure of each of the cells on this page relates to its function. Lucky you.

- Q1 Plants transport food substances from the leaves to growing parts of the plant through phloem tubes.

 Give one feature of a phloem cell that makes it specialised for its function.

 [1 mark]
- O2 Describe how a root hair cell is specialised for its function.

[2 marks]

Chromosomes and Mitosis

In order to survive and grow, our cells have got to be able to divide. And that means our DNA as well...

Chromosomes Contain Genetic Information

- 1) Most cells in your body have a <u>nucleus</u>. The nucleus contains your genetic material in the form of chromosomes.
- 2) Chromosomes are coiled up lengths of DNA molecules.
- Each chromosome carries a large number of genes. Different genes control the development of different characteristics, e.g. hair colour.
- 4) <u>Body cells</u> normally have <u>two copies</u> of each <u>chromosome</u> one from the organism's '<u>mother</u>', and one from its 'father'. So, humans have two copies of chromosome 1, two copies of chromosome 2, etc.
- 5) The diagram shows the 23 pairs of chromosomes from a human cell.

The Cell Cycle Makes New Cells for Growth, Development and Repair

- Body cells in multicellular organisms divide to produce new cells as part of a series of stages called the cell cycle.
- 2) The stage of the cell cycle when the cell divides is called mitosis.
- 3) Multicellular organisms use mitosis to grow or replace cells that have been damaged.
- 4) The end of the cell cycle results in two new cells identical to the original cell, with the same number of chromosomes.
- 5) You need to know about these two main stages of the cell cycle:





Growth & DNA Replication

- 1) In a cell that's not dividing, the DNA is all spread out in long strings.
- 2) Before it divides, the cell has to grow and increase the amount of subcellular structures such as mitochondria and ribosomes.
- 3) It then duplicates its DNA so there's one copy for each new cell. The DNA is copied and forms

 X-shaped chromosomes. Each 'arm' of the chromosome is an exact duplicate of the other.







Once its contents and DNA have been copied, the cell is ready for mitosis...

- 4) The chromosomes <u>line up</u> at the centre of the cell and <u>cell fibres</u> pull them apart. The <u>two arms</u> of each chromosome go to <u>opposite ends</u> of the cell.
- Membranes form around each of the sets of chromosomes. These become the nuclei of the two new cells — the nucleus has divided.
- 6) Lastly, the cutoplasm and cell membrane divide.

The cell has now produced two new daughter cells. The daughter cells contain exactly the same DNA — they're identical. Their DNA is also identical to the parent cell.

A cell's favourite computer game — divide and conquer...

Mitosis can seem tricky at first. But don't worry — just go through it slowly, one step at a time.

- Q1 A student looks at cells in the tip of a plant root under a microscope.

 She counts 11 cells that are undergoing mitosis and 62 cells that are not.
 - a) Calculate the percentage of cells that are undergoing mitosis.
 - b) Suggest how the student can tell whether a cell is undergoing mitosis or not.




[1 mark]

Binary Fission

Prokaryotic cells can reproduce using a type of simple cell division called binary fission.

Prokaryotic Cells Replicate by Binary Fission

In binary fission, the cell... splits into two. There's a bit more to it than that though:



- The circular DNA and plasmid(s) replicate.
- 2) The cell gets bigger and the circular DNA strands move to opposite 'poles' (ends) of the cell.
- 3) The cutoplasm begins to divide and new cell walls begin to form.
- 4) The cytoplasm divides and two daughter cells are produced. Each daughter cell has one copy of the circular DNA, but can have a variable number of copies of the plasmid(s).

Bacteria can divide very quickly if given the right conditions (e.g. a warm environment and lots of nutrients). Some bacteria, such as E. coli, can take as little as 20 minutes to replicate in the right environment. However, if conditions become unfavourable, the cells will stop dividing and eventually begin to die.

Use Mean Division Time to Find the Number of Bacteria in a Population

The mean division time is just the average amount of time it takes for one bacterial cell to divide into two. If you know the mean division time of a cell, you can work out how many times it has divided in a certain amount of time, and so the number of cells it has produced in that time.

EXAMPLE

A bacterial cell has a mean division time of 30 minutes. How many cells will it have produced after 2.5 hours?

- 1) Make sure both times are in the same units. 2.5 hours × 60 = 150 minutes.
- 2) Divide the total time that the bacteria are producing cells by the mean division time. This gives you the number of divisions.
- 3) Multiply 2 by itself for the number of divisions to find the number of cells.

150 minutes ÷ 30 minutes = 5 divisions


 $2^5 = 2 \times 2 \times 2 \times 2 \times 2 = 32$ cells



cell divides to " produce 2 cells. so the number of cells increases by a power of 2 for each division.

Fission is division, there's no splitting hairs there...

Binary fission is the sort of subject where standard form could show up. Head back to p.12 if you're unsure.



O2 E. coli is a type of bacteria. The mean division time of an E. coli cell is 20 minutes. How many cells will a single E. coli cell produce in 2 hours and 40 minutes?

[2 marks]

Culturing Microorganisms



And now for some hands-on stuff. You can grow your own microorganisms and test how effective different antibiotics, antiseptics or disinfectants are at killing them — just watch out for those bacterial rights activists...

You Can Grow Bacteria in the Lab

- Bacteria (and some other microorganisms) are grown (cultured) in a "culture medium", which contains the carbohydrates, minerals, proteins and vitamins they need to grow.
- 2) The culture medium used can be a nutrient broth solution or solid agar jelly.
- Bacteria grown on agar 'plates' will form visible colonies on the surface of the jelly, or will spread out to give an even covering of bacteria.
 - To make an agar plate, <u>hot</u> agar jelly is poured into shallow round plastic dishes called <u>Petri dishes</u>.
 - When the jelly's cooled and set, <u>inoculating loops</u> (wire loops) can be used
 to <u>transfer</u> microorganisms to the culture medium. Alternatively, a <u>sterile</u>
 dropping pipette and <u>spreader</u> can be used to get an <u>even covering</u> of bacteria.
 - · The microorganisms then multiplu.



inoculating loop

Petri dish

4) In the <u>lab at school</u>, cultures of microorganisms are not kept <u>above 25 °C</u>, because <u>harmful pathogens</u> are more likely to grow above this temperature.

5) In industrial conditions, cultures are incubated at higher temperatures so that they can grow a lot faster.

You Can Investigate the Effect of Antibiotics on Bacterial Growth

You can test the action of antibiotics (or antiseptics) on cultures of bacteria:

- Place paper discs soaked in different types (or different concentrations) of antibiotics on an agar plate that has an even covering of bacteria. Leave some space between the discs.
- 2) The antibiotic should diffuse (soak) into the agar jelly. Antibiotic-resistant bacteria (i.e. bacteria that aren't affected by the antibiotic see p.51) will continue to grow on the agar around the paper discs, but non-resistant strains will die. A clear area will be left where the bacteria have died this is called an inhibition zone.

 (Inhibition zone)
- 3) Make sure you use a control. This is a paper disc that has not been soaked in an antibiotic. Instead, soak it in sterile water. You can then be sure that any difference between the growth of the bacteria around the control disc and around one of the antibiotic discs is due to the effect of the antibiotic alone (and not something weird in the paper, for example).
- 4) Leave the plate for 48 hours at 25 °C.
- 5) The more effective the antibiotic is against the bacteria, the larger the inhibition zone will be see next page.

You Need to Use Uncontaminated Cultures

Contamination by unwanted microorganisms will affect your results and can potentially result in the growth of pathogens. To avoid this:

- The Petri dishes and culture medium must be <u>sterilised</u> before use (e.g. by heating to a high temperature), to <u>kill</u> any <u>unwanted</u> microorganisms that may be lurking on them.
- If an inoculating loop is used to transfer the bacteria to the culture medium, it should be sterilised first by passing it through a hot flame.
- After transferring the bacteria, the lid of the Petri dish should be <u>lightly</u> taped on to stop microorganisms from the air getting in.
- The Petri dish should be stored <u>upside down</u> to <u>stop</u> drops of <u>condensation</u> falling onto the agar surface.



paper disc -

with antibiotic

bacteria


growing

paper disc

control







PRACTICAL

Culturing Microorganisms

Calculate the Sizes of the Inhibition Zones to Compare Results

You can compare the effectiveness of different antibiotics (or antiseptics) on bacteria by looking at the relative sizes of the inhibition zones.

The larger the inhibition zone around a disc, the more effective the antibiotic is against the bacteria.

You can do this <u>by eye</u> if there are large differences in size. But to get more accurate results it's a good idea to calculate the <u>area</u> of the inhibition zones using their <u>diameter</u> (the distance <u>across</u>).

Don't open the Petri dish to measure the inhibition zones they should be visible through the bottom of the dish.

To calculate the area of an inhibition zone, you need to use this equation:

This is the equation for the area of a circle. You're likely to use the units cm² or mm².



Area = πr^2 r is the radius of the inhibition zone — it's equal to half the diameter.

 $\boldsymbol{\pi}$ is just a number. You should have a button for it on your calculator. If not, just use the value 3.14.

EXAMPLE

The diagram below shows the inhibition zones produced by antibiotics A and B. Use the areas of the inhibition zones to compare the effectiveness of the antibiotics.



- Divide the diameter of zone A by two to find the radius.
- 2) Stick the radius value into the equation area = πr^2 .
- 3) Repeat steps 1 and 2 for zone B.
- 4) Compare the sizes of the areas.
 314 mm² is just over twice 154 mm², so you could say that:

Radius of A = $14 \div 2 = 7 \text{ mm}$


Area of A = $\pi \times 7^2 = 154 \text{ mm}^2$

Radius of B = $20 \div 2 = 10 \text{ mm}$ Area of B = $\pi \times 10^2 = 314 \text{ mm}^2$

The inhibition zone of antibiotic B is roughly twice the size of the inhibition zone of antibiotic A.

You Can Also Find the Area of a Colony

The equation above can also be used to calculate the <u>area</u> of a bacterial <u>colony</u>. You just need to measure the <u>diameter</u> of the colony you are interested in first.



Agar — my favourite jelly flavour after raspberry...

Microorganisms might be the perfect pets. You don't have to walk them, they won't get lonely and they hardly cost anything to feed. But whatever you do, do not feed them after midnight.


Q1 Helena investigated the effect of four different concentrations (A-D) of an antiseptic on the growth of bacteria. The diagram on the right shows the results.

- Suggest which concentration of antiseptic was the strongest.
- b) Calculate the size of the inhibition zone for Concentration C.
 Give your answer to the nearest mm².
- Suggest what Helena used as a control in this investigation.

[1 mark]

[2 marks]

[1 mark]



Stem Cells

Stem cell research has exciting possibilities, but it's also pretty controversial.

Embryonic Stem Cells Can Turn into ANY Type of Cell

- 1) Differentiation is the process by which a cell changes to become specialised for its job see p.14.
- 2) <u>Undifferentiated</u> cells, called <u>stem cells</u>, can divide to produce lots <u>more</u> undifferentiated cells. They can differentiate into different types of cell, depending on what instructions they're given.
- 3) Stem cells are found in early <u>human embryos</u>. They're <u>exciting</u> to doctors and medical researchers because they have the potential to turn into <u>any</u> kind of cell at all. This makes sense if you think about it <u>all</u> the <u>different types</u> of cell found in a human being have to come from those <u>few cells</u> in the early embryo.
- 4) Adults also have stem cells, but they're only found in certain places, like bone marrow. Unlike embryonic stem cells, they can't turn into any cell type at all, only certain ones, such as blood cells.
- 5) Stem cells from embryos and bone marrow can be grown in a lab to produce clones (genetically identical cells) and made to differentiate into specialised cells to use in medicine or research.

Stem Cells May Be Able to Cure Many Diseases

- Medicine already uses <u>adult stem cells</u> to cure <u>disease</u>. For example, <u>stem cells</u> transferred from the bone marrow of a <u>healthy person</u> can <u>replace faulty blood cells</u> in the patient who receives them.
- 2) Embryonic stem cells could also be used to replace faulty cells in sick people you could make insulin-producing cells for people with diabetes, nerve cells for people paralysed by spinal injuries, and so on.
- 3) In a type of cloning, called therapeutic cloning, an embryo could be made to have the same genetic information as the patient. This means that the stem cells produced from it would also contain the same genes and so wouldn't be rejected by the patient's body if used to replace faulty cells.
- 4) However, there are <u>risks</u> involved in using stem cells in medicine. For example, stem cells grown in the lab may become <u>contaminated</u> with a <u>virus</u> which could be <u>passed on</u> to the patient and so make them <u>sicker</u>.

Some People Are Against Stem Cell Research

- Some people are <u>against</u> stem cell research because they feel that human embryos <u>shouldn't</u> be used for experiments since each one is a <u>potential human life</u>.
- 2) Others think that curing existing patients who are suffering is more important than the rights of embryos.
- 3) They argue that the embryos used in the research are usually <u>unwanted ones</u> from <u>fertility clinics</u> which, if they weren't used for research, would probably just be <u>destroyed</u>.
- 4) However, campaigners for the rights of embryos feel that scientists should concentrate more on finding and developing <u>other sources</u> of stem cells, so people could be helped <u>without</u> having to use embryos.
- 5) In some countries stem cell research is banned. It's allowed in the UK as long as it follows strict guidelines.

Stem Cells Can Produce Identical Plants

- 1) In plants, stem cells are found in the meristems (parts of the plant where growth occurs see p.42).
- 2) Throughout the plant's entire life, cells in the meristem tissues can differentiate into any type of plant cell.
 - These stem cells can be used to produce clones (identical copies) of whole plants quickly and cheaply.
- They can be used to grow more plants of rare species (to prevent them being wiped out).
- Stem cells can also be used to grow crops of <u>identical</u> plants that have desired <u>features</u> for farmers, for example, <u>disease resistance</u>.

But florists cell stems, and nobody complains about that...

Whatever your opinion is, make sure know the uses of stem cells and the arguments for and against using them.

O1 How can stem cells be used to preserve rare plant species? [2 marks]

Diffusion


Particles move about randomly, and after a bit they end up evenly spaced. It's not rocket science, is it...

Don't Be Put Off by the Fancy Word

- "<u>Diffusion</u>" is simple. It's just the <u>gradual movement</u> of particles from places where there are <u>lots</u> of them to places where there are <u>fewer</u> of them — it's just the <u>natural tendencu</u> for stuff to <u>spread out</u>.
- 2) Unfortunately you also have to learn the fancy way of saying the same thing, which is this:

<u>DIFFUSION</u> is the <u>SPREADING OUT</u> of <u>particles</u> from an area of <u>HIGHER CONCENTRATION</u> to an area of <u>LOWER CONCENTRATION</u>.

- Diffusion happens in both solutions and gases that's because the particles in these substances are free to move about randomly.
- 4) The <u>simplest type</u> is when different <u>gases</u> diffuse through each other. This is what's happening when the smell of perfume diffuses through the air in a room:




- 5) The bigger the concentration gradient (the difference in concentration), the faster the diffusion rate.
- 6) A higher temperature will also give a faster diffusion rate because the particles have more energy, so move around faster.

Cell Membranes Are Kind of Clever...

- 1) They're clever because they hold the cell together BUT they let stuff in and out as well.
- 2) Dissolved substances can move in and out of cells by diffusion.
- Only very <u>small</u> molecules can <u>diffuse</u> through cell membranes though

 things like oxygen (needed for respiration see page 62), glucose, amino acids and water.
- 4) Big molecules like starch and proteins can't fit through the membrane:



- 5) Just like with diffusion in air, particles flow through the cell membrane from where there's a higher concentration (a lot of them) to where there's a lower-concentration (not such a lot of them).
- 6) They're only moving about <u>randomly</u> of course, so they go <u>both</u> ways but if there are a lot <u>more</u> particles on one side of the membrane, there's a <u>net</u> (overall) movement <u>from</u> that side.
- 7) The larger the surface area of the membrane, the faster the diffusion rate, because more particles can pass through at once see page 23.

Revision by diffusion - you wish...

Wouldn't it be great if all the ideas in this book would just gradually drift across into your mind...

- O1 A student adds a drop of ink to a glass of cold water.
 - a) What will the student observe to happen to the drop of ink. Explain your answer.
 - b) How might the observation differ if the ink was added to a glass of warm water?

[2 marks]

[1 mark]


Osmosis

If you've got your head round diffusion, osmosis will be a breeze. If not, have another read of the previous page.

Osmosis is a Special Case of Diffusion, That's All

OSMOSIS is the movement of water molecules across a partially permeable membrane from a region of higher water concentration to a region of lower water concentration.

- 1) A partially permeable membrane is just one with very small holes in it. So small, in fact, only tiny molecules (like water) can pass through them, and bigger molecules (e.g. sucrose) can't.
- 2) The water molecules actually pass both waus through the membrane during osmosis. This happens because water molecules move about randomly all the time.
- 3) But because there are more water molecules on one side than on the other, there's a steady net flow of water into the region with fewer water molecules. i.e. into the stronger sugar solution.
- 4) This means the strong sugar solution gets more dilute. The water acts like it's truing to "even up" the concentration either side of the membrane.



5) Osmosis is a type of diffusion — passive movement of water particles from an area of higher water concentration to an area of lower water concentration.

You can Observe How Sugar Solutions Affect Plant Tissue

PRACTICAL

By calculating the percentage change

(see p.130), you can compare the effect


of sugar concentration on cylinders I that didn't have the same initial mass.

An increase in mass will give a positive

give a negative percentage change.

There's a fairly dull experiment you can do to show osmosis at work.

1) You cut up an innocent potato into identical cylinders, and get some beakers with different sugar solutions in them. One should be <u>pure water</u> and another should be a very concentrated sugar solution (e.g. 1 mol/dm³). Then you can have a few others with concentrations in between (e.g. 0.2 mol/dm3, 0.4 mol/dm3, 0.6 mol/dm3, etc.)



- 2) You measure the mass of the cylinders, then leave one cylinder in each beaker for twenty four hours or so.
- 3) Then you take them out, dry them with a paper towel and measure their masses again.
- percentage change and a decrease will 4) If the cylinders have drawn in water by osmosis, they'll have increased in mass. If water has been drawn out, they'll have decreased in mass.

You can calculate the percentage change in mass, then plot a few graphs and things.

5) The dependent variable is the chip mass and the independent variable is the concentration of the sugar solution. All other variables (volume of solution, temperature, time, type of sugar used, etc. etc.) must be kept the same in each case or the experiment won't be a fair test.

6) Like any experiment, you need to be aware of how errors (see p.5) may arise. Sometimes they may occur when carrying out the method, e.g. if some potato cylinders were not fully dried, the excess water would give a higher mass, or if water evaporated from the beakers, the concentrations of the sugar solutions would change. You can reduce the effect of these errors by repeating the experiment and calculating a mean percentage change at each concentration.

You could also carry out this experiment using different salt solutions and see what effect they have on potato chip mass. on polaco corp

And to all you cold-hearted potato murderers...

Just remember, osmosis is really just a fancy word for the diffusion of water molecules. It's simple really.

01 Explain what will happen to the mass of a piece of potato added to a concentrated salt solution.


[2 marks]



Active Transport

Sometimes substances need to be absorbed against a concentration gradient, i.e. from a lower to a higher concentration. This process is lovingly referred to as <u>ACTIVE TRANSPORT</u>.

Root Hairs Take In Minerals and Water



- As you saw on page 14, the cells on plant roots grow into "hairs" which stick out into the soil.
- Each branch of a root will be covered in millions of these microscopic hairs.
- This gives the plant a large surface area for absorbing water and mineral ions from the soil.
- 4) Plants need these mineral ions for healthy growth.
- The concentration of minerals is usually <u>higher</u> in the <u>root hair</u> cells than in the <u>soil</u> around them.

Inside

the

6) So the root hair cells can't use diffusion to take up minerals from the soil.

Root Hairs Take in Minerals Using Active Transport

- Minerals should move <u>out</u> of the root hairs if they followed the rules of diffusion.
 The cells must use another method to draw them in.
- 2) That method is, in fact, a conveniently mysterious process called "active transport".
- Active transport allows the plant to absorb minerals from a very <u>dilute</u> solution, <u>against</u> a
 concentration gradient. This is essential for its growth. But active transport needs <u>ENERGY</u>
 from respiration to make it work.
- Active transport also happens in <u>humans</u>, for example in taking <u>glucose</u> from the <u>gut</u> (see below), and from the <u>kidney tubules</u>.

We Need Active Transport to Stop Us Starving

Active transport is used in the gut when there is a lower concentration of nutrients in the gut, but a higher concentration of nutrients in the blood.

- When there's a higher concentration of glucose and amino acids in the gut they diffuse naturally into the blood.
- <u>BUT</u> sometimes there's a <u>lower concentration</u> of nutrients in the gut than there is in the blood.
- This means that the <u>concentration gradient</u> is the wrong way.
- The same process used in plant roots is used here...
 - ... "Active transport".
- Active transport allows nutrients to be taken into the blood, despite the fact that the concentration gradient is the wrong way.
- 6) This means that <u>glucose</u> can be taken into the bloodstream when its concentration in the blood is already higher than in the <u>gut</u>. It can then be transported to cells, where it's used for respiration (see p.61).

Active transport — get on yer bike...


An important difference between active transport and diffusion is that active transport uses energy. Imagine a pen of sheep in a field. If you open the pen, the sheep will happily diffuse from the area of higher sheep concentration into the field, which has a lower sheep concentration — you won't have to do a thing. To get them back in the pen though, you'll have to put in quite a bit of energy.

Of What is the purpose of active transport in the gut?

[1 mark]

Inside

the blood



Water is taken

into root hair cells by osmosis

(see page 21).

Exchange Surfaces

How easily stuff moves between an organism and its environment depends on its surface area to volume ratio.

Organisms Exchange Substances with their Environment

- 1) Cells can use diffusion to take in substances they need and get rid of waste products. For example:
 - · Oxugen and carbon dioxide are transferred between cells and the environment during gas exchange.
 - In humans, <u>urea</u> (a waste product produced from the breakdown of proteins, see p.75) diffuses from cells into the blood plasma for removal from the body by the kidneys.
- 2) How easy it is for an organism to exchange substances with its environment depends on the organism's surface area to volume ratio (SA: V).

You Can Compare Surface Area to Volume Ratios

A ratio shows how big one value is compared to another. The larger an organism is, the smaller its surface area is compared to its volume. You can show this by calculating surface area to volume ratios:

A hippo can be represented by a 2 cm × 4 cm × 4 cm block.

The area of a surface is found by the equation: LENGTH × WIDTH

So the hippo's total <u>surface area</u> is:

(4 × 4) × 2 (text) and better surfaces of block)

 $(4 \times 4) \times 2$ (top and bottom surfaces of block) + $(4 \times 2) \times 4$ (four sides of the block)

 $+ (4 \times 2) \times 4$ (four sides of the block) $= 64 \text{ cm}^2$.

2 cm

The volume of a block is found by the equation: LENGTH \times WIDTH \times HEIGHT So the hippo's volume is $4\times4\times2=32~\text{cm}^3$.

The surface area to volume ratio of the hippo can be written as <u>64:32</u>. To <u>simplify</u> the ratio, <u>divide both sides</u> of the ratio by the <u>volume</u>.

So the surface area to volume ratio of the hippo is 2:1.

A mouse can be represented by a 1 cm \times 1 cm \times 1 cm block.

Its surface area is $(1 \times 1) \times 6 = 6$ cm².

Its volume is $1 \times 1 \times 1 = 1$ cm³.

So the surface area to volume ratio of the mouse is 6:1.

The cube mouse's surface area is six times its volume, but the cube hippo's surface area is only twice its volume. So the mouse has a larger surface area compared to its volume.

Multicellular Organisms Need Exchange Surfaces

 In <u>single-celled organisms</u>, gases and dissolved substances can diffuse <u>directly into</u> (or out of) the cell across the cell membrane. It's because they have a <u>large surface area</u> compared to their <u>volume</u>, so enough <u>substances</u> can be exchanged across the membrane to <u>supply</u> the volume of the cell.

1 cm

- 2) <u>Multicellular organisms</u> have a <u>smaller surface area</u> compared to their <u>volume</u> <u>not enough</u> substances can diffuse from their outside surface to supply their entire volume. This means they need some sort of <u>exchange surface</u> for efficient diffusion (see pages 24-25 for some examples). The exchange surface structures have to allow <u>enough</u> of the necessary substances to pass through.
- 3) Exchange surfaces are ADAPTED to maximise effectiveness:
 - . They have a thin membrane, so substances only have a short distance to diffuse.
 - They have a large surface area so lots of a substance can diffuse at once.
 - . Exchange surfaces in animals have lots of blood vessels, to get stuff into and out of the blood quickly.
 - · Gas exchange surfaces in animals (e.g. alveoli) are often ventilated too air moves in and out.

Not that I'm endorsing putting animals in boxes...

A large surface area is a key way that organisms' exchange surfaces are made more effective.

Q1 A bacterial cell can be represented by a 2 μm × 2 μm × 1 μm block.

Calculate the cell's surface area to volume ratio.

[3 marks]




Exchanging Substances

This page is about how two different parts of the human body are <u>adapted</u> so that substances can diffuse through them <u>most effectively</u>. The first bit is about how <u>gases</u> in the lungs get <u>into and out of the blood</u>. The second is about how <u>digested food</u> gets from the <u>gut to the blood</u>.

Gas Exchange Happens in the Lungs

- The job of the lungs is to transfer <u>oxygen</u> to the blood and to remove waste carbon dioxide from it.
- To do this the lungs contain millions of little air sacs called alveoli where gas exchange takes place.






Blue = blood with carbon dioxide.

- Red = blood with oxygen.
- The alveoli are specialised to maximise the <u>diffusion</u> of O₂ and CO₂. They have:
 - An enormous surface area (about 75 m² in humans).
 - A moist lining for dissolving gases.
 - Very <u>thin walls</u>.
 - A good blood supply.

The Villi Provide a Really Really Big Surface Area

- The inside of the <u>small intestine</u> is covered in millions and millions of these tiny little projections called <u>villi</u>.
- They increase the surface area in a big way so that digested food is <u>absorbed</u> much more quickly into the <u>blood</u>.
- 3) Notice they have:
 - a single layer of surface cells,
 - a very good blood supply to assist quick absorption.





Al Veoli — the Italian gas man...

Thankfully, our bodies are well adapted for efficient diffusion of substances. But the array of life's snazzy exchange surfaces doesn't stop here, oh no — just take a look at what's coming up on the next page...

Q1 Give one way in which alveoli are adapted for gas exchange.

[1 mark]

Q2 Coeliac disease causes inflammation of the small intestine, which can damage the villi. Suggest why a person with coeliac disease might have low levels of iron in their blood. [2 marks]



More on Exchanging Substances

More stuff on adaptations for diffusion now — only this time, it's plants and fish. Whoopee...

The Structure of Leaves Lets Gases Diffuse In and Out of Cells

- Carbon dioxide <u>diffuses into the air spaces</u> within the leaf, then it <u>diffuses into the cells</u> where photosynthesis happens. The leaf's structure is <u>adapted</u> so that this can happen easily.
- The underneath of the leaf is an <u>exchange surface</u>. It's covered in little holes called <u>stomata</u> which the carbon dioxide diffuses in through.
- Oxugen (produced in photosynthesis) and water vapour also diffuse out through the stomata. (Water vapour is actually lost from all over the leaf surface, but most of it is lost through the stomata.)
- 4) The size of the stomata are controlled by <u>guard cells</u> see page 44. These <u>close</u> the stomata if the plant is losing water faster than it is being replaced by the roots. Without these guard cells the plant would soon <u>wilt</u>.



mouth

gill

arteries

- 5) The <u>flattened shape</u> of the leaf increases the <u>area</u> of this exchange surface so that it's more effective.
- 6) The <u>walls of the cells</u> inside the leaf form another exchange surface. The <u>air spaces</u> inside the leaf increase the <u>area</u> of this surface so there's more chance for carbon dioxide to get into the cells.

The water vapour evaporates from the cells inside the leaf. Then it escapes by diffusion because there's a lot of it inside the leaf and less of it in the air outside.

Gills Have a Large Surface Area for Gas Exchange

- 1) The gills are the gas exchange surface in fish.
- 2) Water (containing oxygen) enters the fish through its mouth and passes out through the gills. As this happens, oxygen diffuses from the water into the blood in the gills and oarbon dioxide diffuses from the blood into the water.
- 3) Each gill is made of lots of thin plates called gill filaments, which give a big surface area for exchange of gases.
- 4) The gill filaments are covered in lots of tiny structures called <u>lamellae</u>, which <u>increase</u> the <u>surface area</u> even more.
- 5) The lamellae have lots of blood capillaries to speed up diffusion.
- They also have a thin surface layer of cells to minimise the distance that the gases have to diffuse.
- 7) <u>Blood</u> flows through the lamellae in one direction and <u>water</u> flows over in the opposite direction. This maintains a <u>large concentration gradient</u> between the water and the blood.
- 8) The <u>concentration of oxygen</u> in the <u>water</u> is always <u>higher</u> than that in the <u>blood</u>, so as much oxygen as possible diffuses from the water into the blood.




There's a theme here — multicellular organisms are really well adapted for getting the substances they need to their cells. It makes sense — if they couldn't do this well, they'd die out. If you're asked in an exam how something's adapted for exchange, think about whether surface area is important — cos it often is.

Of the Give two ways in which the structure of a gill is adapted for effective gas exchange.

[2 marks]

sill filaments



lamellae

Revision Questions for Topic 1

Well, that's $\underline{\text{Topic 1}}$ done and dusted. Now there's only one way to find out whether you've learnt anything from it. And yes, I'm afraid it involves that whole load of questions staring you in the face.

For even more practice, try the ER Retrieval Quiz for Topic 1 — just E scan this QR code!



- Try these questions and <u>tick off each one</u> when you <u>get it right</u>.
- When you're completely happy with a sub-topic, tick it off.

Ce	ells and Microscopy (p.11-13)	
1)	Name five subcellular structures that both plant and animal cells have.	$\sqrt{}$
2)	What three things do plant cells have that animal cells don't?	$\sqrt{}$
3)	Where is the genetic material found in:	
	a) animal cells,	_
	b) bacterial cells?	
4)	What type of organisms are bacteria — prokaryotes or eukaryotes?	
5)	Which gives a higher resolution — a light microscope or an electron microscope?	\checkmark
Di	fferentiation and Division (p.14-16) 🔃	
6)	What is cell differentiation?	$\sqrt{}$
7)	Give three ways that a sperm cell is adapted for swimming to an egg cell.	$\sqrt{}$
8)	Draw a diagram of a nerve cell. Why is it this shape?	\checkmark
9)	What are chromosomes?	
10)	What is the cell cycle?	
11)	What is mitosis used for by multicellular organisms?	
12)	What is the name of the process by which bacteria divide?	\checkmark
Cu	ulturing Microorganisms (p.17-18) 🔃	
13)	a) What is the maximum temperature that microorganisms should be grown at in a school lab? b) Why shouldn't a temperature above this be used?	
14)	There are ways in which you can make sure an experiment testing the effect	_
	of antibiotics on bacteria has not been contaminated. Give three of these ways.	\checkmark
Ste	em Cells (p.19)	
15)	Give two ways that embryonic stem cells could be used to cure diseases.	\checkmark
16)	Why might some people be opposed to the use of human embryos in stem cell research?	\checkmark
Ex	changing Substances (p.20-25)	
17)	What is diffusion?	$\sqrt{}$
18)	Name three substances that can diffuse through cell membranes, and two that can't.	$\sqrt{}$
19)	What type of molecules move by osmosis?	$\sqrt{}$
20)	Give the two main differences between active transport and diffusion.	
21)	Give three adaptations of exchange surfaces that increase the efficiency of diffusion.	
22)	Give two ways that the villi in the small intestine are adapted for absorbing digested food.	
221	Explain how leaves are adapted to maximize the amount of carbon dioxide that sets to their calls	

Cell Organisation

Some organisms contain loads of cells, but how, you might wonder, do all these cells end up making a working human or squirrel... the answer's organisation. Without it, they'd just make a meaty splodge.

Large Multicellular Organisms are Made Up of Organ Systems

- 1) Cells are the basic building blocks that make up all living organisms.
- 2) As you know from page 14, specialised cells carry out a particular function.
- 3) The process by which cells become specialised for a particular job is called differentiation. Differentiation occurs during the development of a multicellular organism.
- 4) These specialised cells form tissues, which form organs, which form organ sustems (see below).
- 5) Large multicellular organisms (e.g. squirrels) have different systems inside them for exchanging and transporting materials.

Epithelial cell

Epithelial tissue

0.1 mm

Similar Cells are Organised into Tissues

A tissue is a group of similar cells that work together to carry out a particular function. It can include more than one tupe of cell.

In mammals (like humans), examples of tissues include:

- 1) Muscular tissue, which contracts (shortens) to move whatever it's attached to.
- 2) Glandular tissue, which makes and secretes chemicals like enzymes and hormones.
- 3) Epithelial tissue, which covers some parts of the body, e.g. the inside of the gut.

Tissues are Organised into Organs

An organ is a group of different tissues that work together to perform a certain function.

For example, the stomach is an organ made of these tissues:

- 1) Muscular tissue, which moves the stomach wall to churn up the food.
- 2) Glandular tissue, which makes digestive juices to digest food.
- 3) Epithelial tissue, which covers the outside and inside of the stomach.

Organs are Organised into Organ Systems

Salivary glands

An organ system is a group of organs working together to perform a particular function.

For example, the digestive system (found in humans and other mammals) breaks down and absorbs food. It's made up of these organs:

- 1) Glands (e.g. the pancreas and salivary glands), which produce digestive juices.
- 2) The stomach and small intestine, which digest food.
- 3) The liver, which produces bile.
- 4) The small intestine, which absorbs soluble food molecules.
- 5) The large intestine, which absorbs water from undigested food, leaving faeces.

Liver Large intestine

Digestive sustem

Stomach

about 10 cm

(over 1000 times longer than an

epithelial cell)

Stomach

Pancreas 4 1

Small intestine

You need to know where these organs are on a

Organ systems work together to make entire organisms.

Soft and guilted — the best kind of tissues...

So in summary, an organism consists of organ systems, which are groups of organs, which are made of tissues, which are groups of cells working together. Now just for the thrill of it, here's a practice question.

The bladder is an organ. Explain what this means. 01

[2 marks]

diagram — see page 31 too.

Enzymes

Chemical reactions are what make you work. And enzymes are what make them work.

Enzymes Are Catalysts Produced by Living Things

- Living things have thousands of different chemical reactions going on inside them all the time.
 These reactions need to be carefully controlled to get the right amounts of substances.
- 2) You can usually make a reaction happen more quickly by raising the temperature. This would speed up the useful reactions but also the unwanted ones too... not good. There's also a limit to how far you can raise the temperature inside a living creature before its cells start getting damaged.
- So... living things produce enzymes that act as biological catalysts. Enzymes reduce the need for high temperatures and we only have enzymes to speed up the useful chemical reactions in the body.

A <u>CATALYST</u> is a substance which <u>INCREASES</u> the speed of a reaction, without being <u>CHANGED</u> or <u>USED UP</u> in the reaction.


4) Enzymes are all large proteins and all proteins are made up of chains of amino acids. These chains are folded into unique shapes, which enzymes need to do their jobs (see below).

Enzymes Have Special Shapes So They Can Catalyse Reactions

- 1) Chemical reactions usually involve things either being split apart or joined together.
- 2) Every enzyme has an active site with a unique shape that fits onto the substance involved in a reaction.
- 3) Enzymes are really picky they usually only catalyse one specific reaction.
- 4) This is because, for the enzyme to work, the substrate has to fit into its active site. If the substrate doesn't match the enzyme's active site, then the reaction won't be catalysed.

The substance that an enzyme acts on is called the substrate.


5) This diagram shows the 'lock and key' model of enzyme action. This is simpler than how enzymes actually work. In reality, the active site changes shape a little as the substrate binds to it to get a tighter fit. This is called the 'induced fit' model of enzyme action.



Rate of

reaction

Enzymes Need the Right Temperature and pH



- 1) Changing the temperature changes the rate of an enzyme-catalysed reaction.
- 2) Like with any reaction, a higher temperature increases the rate at first. But if it gets too hot, some of the bonds holding the enzyme together break. This changes the shape of the enzyme's active site, so the substrate won't fit any more. The enzyme is said to be denatured.
- 45 °C Temp. 3) All enzymes have an optimum temperature that they work best at.
- 4) The <u>pH</u> also affects enzymes. If it's too high or too low, the <u>pH</u> interferes with the <u>bonds</u> holding the enzyme together. This changes the <u>shape</u> of the <u>active site</u> and <u>denatures</u> the enzyme.



5) All enzymes have an optimum pH that they work best at. It's often neutral pH 7, but not always — e.g. pepsin is an enzyme used to break down proteins in the stomach. It works best at pH 2, which means it's well-suited to the acidic conditions there.

If only enzymes could speed up revision...

Make sure you use the special terms like 'active site' and 'denatured' — the examiners will love it.

1 Explain why enzymes have an optimum pH.

[2 marks]

Investigating Enzymatic Reactions

You'll soon know how to investigate the effect of pH on the rate of enzyme activity... I bet you're thrilled.

You Can Investigate the Effect of pH on Enzyme Activity

PRACTICAL

mixture sampled

every 30 seconds

_dropping

pipette

water bath, instead of

a Bunsen and a beaker

of water, to control the temperature.

drop of iodine

solution 嶤 spotting tile

The enzume amulase cataluses the breakdown of starch to maltose. It's easy to detect starch using jodine solution — if starch is present, the iodine solution will change from browny-orange You could use an electric to blue-black. This is how you can investigate how pH affects amulase activity:

amylase,

starch and

buffer

solution

- Put a drop of iodine solution into every well of a spotting tile.
- 2) Place a Bunsen burner on a heat-proof mat, and a tripod and gauze over the Bunsen burner. Put a beaker of water on top of the tripod and heat the water until it is 35 °C (use a thermometer to measure the temperature). Try to keep the temperature of
- the water constant throughout the experiment. 3) Use a syringe to add 1 cm3 of amylase solution and 1 cm3 of a buffer solution with a pH of 5 to a boiling tube. Using test tube holders, put the tube into the beaker of water and wait for five minutes.
- 4) Next, use a different suringe to add 5 cm³ of a starch solution to the boiling tube.
- 5) Immediately mix the contents of the boiling tube and start a stop clock.
- 6) Use continuous sampling to record how long it takes for the amylase to break down all of the starch. To do this, use a dropping pipette to take a fresh sample from the boiling tube every 30 seconds and put a drop into a well. When the iodine solution remains browny-orange, starch is no longer present.
- 7) Repeat the whole experiment with buffer solutions of different pH values to see how pH affects the time taken for the starch to be broken down.
- 8) Remember to control any variables each time (e.g. concentration and volume of amulase solution) to make it a fair test.

= to accurately measure the = pH of your solutions.

Here's How to Calculate the Rate of Reaction

- 1) It's often useful to calculate the rate of reaction after an experiment. Rate is a measure of how much something changes over time.
- 2) For the experiment above, you can calculate the rate of reaction using this formula: E.g. At pH 6, the time taken for amulase to break down all of the starch in a solution

was 90 seconds. So the rate of the reaction = $1000 \div 90 = 11 \text{ s}^{-1}$ (2 s.f.)



per unit time. per unit time.

1000

3) If an experiment measures how much something changes over time, you calculate the rate of reaction by dividing the amount that it has changed by the time taken.

EXAMPLE

The enzyme catalase catalyses the breakdown of hydrogen peroxide into water and oxygen. During an investigation into the activity of catalase, 24 cm3 of oxygen was released in 50 seconds (s). Calculate the rate of the reaction. Write your answer in cm³/s.

Amount of product formed = change = 24 cm³ Rate of reaction = change \div time = 24 cm³ \div 50 s = 0.48 cm³/s

Mad scientists — they're experi-mental...

You could adapt this experiment to investigate how factors other than pH affect the rate of amylase activity. E.g. you could use a water bath set to different temperatures to investigate the effect of temperature.

O1 An enzyme-controlled reaction was carried out at pH 4. After 2 minutes, 36 cm³ of product had been released. Calculate the rate of reaction in cm³/s.




[1 mark]

Enzymes and Digestion

The enzymes used in digestion are produced by cells and then released into the gut to mix with food.

Digestive Enzymes Break Down Big Molecules

 Starch, proteins and fats are BIG molecules. They're too big to pass through the walls of the digestive system, so digestive enzymes break these BIG molecules down into smaller ones like sugars (e.g. glucose and maltose), amino acids, glycerol and fatty acids. These smaller, soluble molecules can pass easily through the walls of the digestive system, allowing them to be absorbed into the bloodstream.



2) The body makes good use of the <u>products</u> of digestion. They can be used to make <u>new carbohydrates</u>, <u>proteins</u> and <u>lipids</u>. Some of the <u>glucose</u> (a carbohydrate) that's made is used in <u>respiration</u> (see p.61).

and oils.

The small intestine

Glucerol & fatty acids

Bile Neutralises the Stomach Acid and Emulsifies Fats

- 1) Bile is produced in the liver. It's stored in the gall bladder before it's released into the small intestine.
- 2) The <u>hydrochloric acid</u> in the stomach makes the pH <u>too acidic</u> for enzymes in the small intestine to work properly. Bile is <u>alkaline</u> it <u>neutralises</u> the acid and makes conditions <u>alkaline</u>. The enzymes in the small intestine <u>work best</u> in these alkaline conditions.

enzymes

Lipases are made in two places: 1) The pancreas

3) It emulsifies fats. In other words it breaks the fat into tinu droplets. This gives a much bigger surface area of fat for the enzyme lipase to work on — which makes its digestion faster.

What do you call an acid that's eaten all the pies...

Make sure you know the examples of amylase, protease and lipase, and the reactions that they catalyse.


D1 Bile is a product of the liver. Describe and explain its role in digestion. [4 marks]

More on Enzymes and Digestion

So now you know what the enzymes do, here's a nice big picture of the whole of the digestive system.

The Breakdown of Food is Catalysed by Enzymes

- 1) Enzymes used in the digestive system are produced by specialised cells in glands and in the gut lining.
- 2) Different enzymes catalyse the breakdown of different food molecules.



Mmmm — so who's for a chocolate digestive...

Did you know that the whole of your digestive system is actually a hole that goes right through your body. Think about it. It just gets loads of food, digestive juices and enzymes piled into it. Most of it's then absorbed into the body and the rest is politely stored for removal.

O1 Name the three parts of the digestive system that produce protease enzymes.

[3 marks]

PRACTICAL

Food Tests

There are some clever ways to identify what type of food molecule a sample contains. For each of the tests, you need to prepare a food sample. It's the same each time though — here's what you'd do:

- 1) Get a piece of food and break it up using a pestle and mortar.
- 2) Transfer the ground up food to a beaker and add some distilled water.
- 3) Give the mixture a good stir with a glass rod to dissolve some of the food.
- 4) Filter the solution using a funnel lined with filter paper to get rid of the solid bits of food.



Use the Benedict's Test to Test for Sugars

Sugars are found in all sorts of foods such as biscuits, cereal and bread. There are two types of sugars - non-reducing and reducing. You can test for reducing sugars in foods using the Benedict's test:

- 1) Prepare a food sample and transfer 5 cm³ to a test tube.
- 2) Prepare a water bath so that it's set to 75 °C.
- 3) Add some Benedict's solution to the test tube (about 10 drops) using a pipette.
- 4) Place the test tube in the water bath using a test tube holder and leave it in there for 5 minutes. Make sure the tube is pointing away from you.
- 5) If the food sample contains a reducing sugar, the solution in the test tube will change from the normal blue colour to green, yellow or brick-red — it depends on how much sugar is in the food.

Use Iodine Solution to Test for Starch

You can also check food samples for the presence of starch. Foods like pasta, rice and potatoes contain a lot of starch. Here's how to do the test:

- 1) Make a food sample and transfer 5 cm3 of your sample to a test tube.
- 2) Then add a few drops of iodine solution and gently shake the tube to mix the contents. If the sample contains starch, the colour of the solution will change from browny-orange to black or blue-black.

Use the Biuret Test to Test for Proteins

You can use the biuret test to see if a type of food contains protein. Meat and cheese are protein rich and good foods to use in this test. Here's how it's done:

- 1) Prepare a sample of your food and transfer 2 cm³ of your sample to a test tube.
- 2) Add 2 cm³ of biuret solution to the sample and mix the contents of the tube by gently shaking it.
- 3) If the food sample contains protein, the solution will change from blue to purple. If no protein is present, the solution will stay blue.

Use the Sudan III Test to Test for Lipids

Lipids are found in foods such as olive oil, margarine and milk.


You can test for the presence of lipids in a food using Sudan III stain solution.

- 1) Prepare a sample of the food you're testing (but you don't need to filter it). Transfer about 5 cm3 into a test tube.
- 2) Use a pipette to add 3 drops of Sudan III stain solution to the test tube and gently shake the tube.
- 3) Sudan III stain solution stains lipids. If the sample contains lipids, the mixture will separate out into two layers. The top layer will be bright red. If no lipids are present, no separate red layer will form at the top of the liquid.

All this talk of food is making me hungry...

Make sure you do a risk assessment before starting these tests — there are a lot of chemicals to use here.

Name the chemical that you would use to test a sample for the presence of starch.














The Lungs

You need to get oxygen into your bloodstream to supply your cells for respiration. You also need to get rid of carbon dioxide from your blood. This all happens in your lungs when you breathe air in and out.

The Lungs Are in the Thorax

- 1) The thorax is the top part of your body.
- 2) It's separated from the lower part of the body by the diaphragm.
- The <u>lungs</u> are like big pink <u>sponges</u> and are protected by the <u>ribcage</u>.
 They're surrounded by the pleural membranes.
- 4) The air that you breathe in goes through the <u>trachea</u>. This splits into two tubes called <u>bronchi</u> (each one is a bronchus), one going to each lung.
- The bronchi split into progressively smaller tubes called <u>bronchioles</u>.
- 6) The bronchioles finally end at small bags called <u>alveolis</u> where the gas exchange takes place (see below).



Alveoli Carry Out Gas Exchange in the Body

- The <u>lungs</u> contain millions and millions of little air sacs called <u>alveoli</u>, surrounded by a <u>network</u> of <u>blood capillaries</u>. This is where <u>gas exchange</u> happens.
- 2) The <u>blood</u> passing next to the alveoli has just returned to the lungs from the rest of the body, so it contains <u>lots</u> of <u>carbon dioxide</u> and <u>very little oxygen</u>. <u>Oxygen diffuses out of the alveolus</u> (high concentration) into the <u>blood</u> (low concentration). <u>Carbon dioxide</u> diffuses <u>out of the blood</u> (high concentration) into the <u>alveolus</u> (low concentration) to be breathed out.



- When the blood reaches body cells <u>oxugen</u> is released from the <u>red blood cells</u> (where there's a high concentration) and diffuses into the body cells (where the concentration is low).
- 4) At the same time, <u>carbon dioxide</u> diffuses out of the <u>body cells</u> (where there's a high concentration) into the <u>blood</u> (where there's a low concentration). It's then carried back to the <u>lungs</u>.

You Can Calculate the Breathing Rate in Breaths Per Minute

Rate calculations pop up all the time in biology, and you're expected to know how to do them — thankfully they're pretty easy. Breathing rate is the sort of thing that you could get asked to work out in your exam.

EXAMPLE

Kerry takes 91 breaths in 7 minutes. Calculate her average breathing rate in breaths per minute.

breaths per minute = number of breaths ÷ number of minutes

 $= 91 \div 7$

= 13 breaths per minute

Stop huffing and puffing and just learn it...

Alveoli are really well adapted for carrying out gas exchange. It could be a wise move to learn all about exactly how they're adapted. You met them back on page 24, so head back there if you need a reminder.

Q1 During a 12 minute run, Aaqib took 495 breaths.

Calculate his average breathing rate in breaths per minute.

Circulatory System — The Heart

The circulatory system carries food and oxygen to every cell in the body. As well as being a delivery service, it's also a waste collection service — it carries waste products to where they can be removed from the body.

The DOUBLE Circulatory System, Actually

The circulatory system is made up of the heart, blood vessels and blood. Humans have a double circulatory system — two circuits joined together:

- 1) In the first one, the right ventricle (see below) pumps deoxugenated blood (blood without oxygen) to the lungs to take in oxygen. The blood then returns to the heart.
- 2) In the second one, the left ventricle (see below) pumps oxugenated blood around all the other organs of the bodu. The blood gives up its oxugen at the bodu cells and the deoxugenated blood returns to the heart to be pumped out to the lungs again.



The Heart Contracts to Pump Blood Around The Body

- 1) The heart is a pumping organ that keeps the blood flowing around the body. The walls of the heart are mostly made of muscle tissue.
- 2) The heart has valves to make sure that blood flows in the right direction — they prevent it flowing backwards.
- 3) This is how the heart uses its four chambers (right atrium. right ventricle, left atrium and left ventricle) to pump blood around:
 - 1) Blood flows into the two atria from the vena cava and the pulmonary vein.
 - 2) The atria contract, pushing the blood into the ventricles.
 - 3) The ventricles contract, forcing the blood into the pulmonary artery and the aorta, and out of the heart.
 - 4) The blood then flows to the organs through arteries, and returns through veins (see next page).

5) The atria fill again and the whole cucle starts over.

The heart also needs its own supply of oxugenated blood. Arteries called coronary arteries branch off the aorta and surround the heart, making sure that it gets all the oxygenated blood it needs.

right left. atrium (No, we haven't right left entricle ventricle whose heart it is.)

made a mistake — this is the right and left side of the person

Left

Qide

Atrium is when there is just one. Atria is plural. ริยมัยนี้นักเกิดเกรา

The Heart Has a Pacemaker

1) Your resting heart rate is controlled by a group of cells in the right atrium wall that act as a pacemaker.

Right

Side

- 2) These cells produce a small electric impulse which spreads to the surrounding muscle cells, causing them to contract.
- 3) An artificial pacemaker is often used to control heartbeat if the natural pacemaker cells don't work properly (e.g. if the patient has an irregular heartbeat). It's a little device that's implanted under the skin and has a wire going to the heart. It produces an electric current to keep the heart beating regularly.

Okay — let's get to the heart of the matter...

Interesting fact — when doctors use a stethoscope to listen to your heart, it's the valves closing that they hear.

- Which chamber of the heart pumps deoxygenated blood to the lungs? Q1 [1 mark]
- O2 What is the function of the coronary arteries?

Circulatory System — Blood Vessels

Want to know more about the circulatory system... Good. Because here's a whole extra page.


Blood Vessels are Designed for Their Function

There are three different types of blood vessel:


- 1) ARTERIES these carry the blood away from the heart.
- 2) CAPILLARIES these are involved in the exchange of materials at the tissues.
- 3) VEINS these carry the blood to the heart.

Arteries Carry Blood Under Pressure

- The heart pumps the blood out at <u>high pressure</u> so the arteru walls are strong and elastic.
- 2) The walls are thick compared to the size of the hole down the middle (the "lumen" silly name!).
- They contain thick layers of <u>muscle</u> to make them <u>strong</u>, and <u>elastic fibres</u> to allow them to stretch and <u>spring back</u>.



Capillaries are Really Small




- Arteries branch into <u>capillaries</u>.
- 2) Capillaries are really tiny too small to see.
- 3) They carry the blood really close to every cell in the
- body to <u>exchange substances</u> with them.

 4) They have permeable walls, so substances can diffuse in and out.
- 5) They supply food and oxugen, and take away waste like CO.
- 6) Their walls are usually <u>only one cell thick</u>. This <u>increases</u> the rate of diffusion by <u>decreasing</u> the <u>distance</u> over which it occurs.

Veins Take Blood Back to the Heart

- Capillaries eventually join up to form veins.
 The blood is at lower pressure in the veins so the walls don't need to be as thick as artery walls.
- They have a bigger lumen than arteries to help the blood flow despite the lower pressure.
- They also have <u>valves</u> to help keep the blood flowing in the <u>right direction</u>.



You Can Calculate the Rate of Blood Flow

You might get asked to calculate the <u>rate of blood flow</u> in your exam. Thankfully, it's not too tricky. Take a look at this example:

EXAMPLE

1464 ml of blood passed through an artery in 4.5 minutes.

Calculate the rate of blood flow through the artery in ml/min.

rate of blood flow = volume of blood \div number of minutes = 1464 \div 4.5 = 325 ml/min

Learn this page — don't struggle in vein...

Here's an interesting fact for you — your body contains about 60 000 miles of blood vessels.

Q1 2.175 litres of blood passed through a vein in 8.7 minutes. Calculate the rate of blood flow through the vein in ml/min.

[2 marks]



Circulatory System — Blood

Blood is a tissue. One of its jobs is to act as a huge transport system. There are four main things in blood...

Red Blood Cells Carry Oxygen

- The job of red blood cells is to carry <u>oxygen</u> from the lungs to all the cells in the body.
- Their shape is a <u>biconcave disc</u> (like a doughnut) this gives a <u>large surface area</u> for absorbing <u>oxygen</u>.
- They don't have a nucleus this allows more room to carry oxygen.
- 4) They contain a red pigment called haemoglobin.
- 5) In the <u>lungs</u>, haemoglobin binds to <u>oxygen</u> to become <u>oxyhaemoglobin</u>. In body tissues, the reverse happens oxyhaemoglobin splits up into haemoglobin and oxygen, to <u>release oxygen</u> to the <u>cells</u>.



The more red blood cells you've got, the more oxygen can get to your cells. At high altitudes there's less oxygen in the air — so people who live there produce more red blood cells to compensate.

White Blood Cells Defend Against Infection







- Some can change shape to gobble up unwelcome microorganisms, in a process called <u>phagocytosis</u>.
- Others produce <u>antibodies</u> to fight microorganisms, as well as <u>antitoxins</u> to neutralise any toxins produced by the microorganisms.
 - Unlike red blood cells, they do have a nucleus.

Platelets Help Blood Clot

- These are <u>small fragments</u> of <u>cells</u>. They have <u>no nucleus</u>.
- They help the blood to <u>clot</u> at a wound to stop all your <u>blood pouring out</u> and to stop <u>microorganisms</u> getting in.
 (So basically platelets just float about waiting for accidents to happen.)
- 3) Lack of platelets can cause excessive bleeding and bruising.



Plasma is the Liquid That Carries Everything in Blood

This is a pale straw-coloured liquid which carries just about everything:

- 1) Red and white blood cells and platelets.
- Nutrients like <u>glucose</u> and <u>amino acids</u>.
 These are the soluble products of digestion which are absorbed from the gut and taken to the cells of the body.
- 3) Carbon dioxide from the organs to the lungs.
- 4) Urea from the liver to the kidneys.
- 5) Hormones.
- 6) Proteins.
- 7) Antibodies and antitoxins produced by the white blood cells.

Platelets — ideal for small dinners...

Blood tests can be used to diagnose loads of things — not just disorders of the blood. This is because the blood transports so many chemicals produced by so many organs... and it's easier to take blood than, say, a piece of muscle.



O2 State the function of the cell labelled X in the image on the right.





[1 mark]

Cardiovascular Disease

<u>Cardiovascular disease</u> is a term used to describe diseases of the heart or blood vessels, for example coronary heart disease. This page tells you all about how <u>stents</u> and <u>statins</u> are used to combat coronary heart disease.

Stents Keep Arteries Open

Coronary heart disease is when the coronary arteries that supply the blood to
the muscle of the heart get blocked by layers of fatty material building up.
This causes the arteries to become narrow, so blood flow is restricted and
there's a lack of oxugen to the heart muscle — this can result in a heart attack.

outside of heart coronary artery

 Stents are tubes that are inserted inside arteries. They keep them open, making sure blood can pass through to the heart muscles. This keeps the person's heart beating (and the person alive).



- 3) Stents are a way of lowering the risk of a heart attack in people with coronary heart disease. They are effective for a long time and the recovery time from the surgery is relatively quick.
- 4) On the down side, there is a risk of <u>complications</u> during the operation (e.g. heart attack) and a risk of <u>infection</u> from surgery. There is also the risk of patients developing a <u>blood clot</u> near the stent this is called thrombosis.

Statins Reduce Cholesterol in the Blood

- Cholesterol is an essential lipid that your body produces and needs to function properly. However, too
 much of a certain type of cholesterol (known as 'bad' or LDL cholesterol) can cause health problems.
- Having too much of this 'bad' cholesterol in the bloodstream can cause <u>fatty deposits</u> to form inside <u>arteries</u>, which can lead to <u>coronary heart disease</u>.
- Statins are drugs that can reduce the amount of 'bad' cholesterol present in the bloodstream. This slows down the rate of fatty deposits forming.

Statins Have Advantages and Disadvantages

Advantages

- By reducing the amount of 'bad' cholesterol in the blood, statins can <u>reduce</u> the risk of <u>strokes</u>, <u>coronary heart disease</u> and <u>heart attacks</u>.
- 2) As well as reducing the amount of 'bad' cholesterol, statins can increase the amount of a beneficial type of cholesterol (known as 'good' or HDL cholesterol) in your bloodstream. This type can remove 'bad' cholesterol from the blood.
- Some studies suggest that statins may also help prevent some other diseases.

Disadvantages

- Statins are a long-term drug that must be taken regularly. There's the risk that someone could forget to take them.
- Statins can sometimes cause <u>negative side effects</u>, e.g. headaches. Some of these side effects can be <u>serious</u>, e.g. kidney failure, liver damage and memory loss.
- The effect of statins <u>isn't instant</u>.
 It takes <u>time</u> for their <u>effect</u> to kick in.

Unlike stents and statins, using CGP books only has advantages...

Stents and statins might be good treatments for coronary heart disease, but they're not perfect. Make sure you're aware of the drawbacks as well as the advantages of each. That way, you'll be covered if they come up in the exam.

Q1 a) How can stents be used to reduce the risk of heart attacks in people with coronary heart disease?

[2 marks]

b) Suggest two disadvantages of treating patients using stents.

[2 marks]

More on Cardiovascular Disease

With more fakery than a 'Rollecks' watch, this page is about artificial hearts, artificial blood and replacing heart valves. All a bit gruesome, I'll admit — but it's life-saving stuff.

An Artificial Heart Can Pump Blood Around the Body

If a patient has heart failure, doctors may perform a heart transplant (or heart and lungs transplant if the lungs are also diseased) using donor organs from people who have recently died. However, if donor organs aren't available right away or they're not the best option, doctors may fit an artificial heart.

- Artificial hearts are mechanical devices that <u>pump blood</u> for a person whose own heart has <u>failed</u>.
 They're <u>usually</u> only used as a <u>temporary</u> fix, to keep a person <u>alive</u> until a <u>donor heart</u> can be found or to help a person <u>recover</u> by allowing the heart to <u>rest</u> and <u>heal</u>. In some cases though they're used as a <u>permanent</u> fix, which <u>reduces</u> the <u>need</u> for a donor heart.
- 2) The main <u>advantage</u> of artificial hearts is that they're <u>less likely to be rejected</u> by the body's immune system than a donor heart. This is because they're made from <u>metals</u> or <u>plastics</u>, so the body doesn't recognise them as 'foreign' and attack in the same way as it does with living tissue.
- 3) But surgery to fit an artificial heart (as with transplant surgery) can lead to bleeding and infection. Also, artificial hearts don't work as well as healthy natural ones parts of the heart could wear out or the electrical motor could fail. Blood doesn't flow through artificial hearts as smoothly, which can cause blood clots and lead to strokes. The patient has to take drugs to thin their blood and make sure this doesn't happen, which can cause problems with bleeding if theu're hurt in an accident.

Faulty Heart Valves Can Be Replaced With Biological or Mechanical Valves

- 1) The valves in the heart can be damaged or weakened by heart attacks, infection or old age.
- 2) The damage may cause the <u>valve tissue</u> to <u>stiffen</u>, so it <u>won't open properly</u>. Or a valve may become <u>leaky</u>, allowing blood to flow in <u>both directions</u> rather than just forward. This means that blood <u>doesn't circulate</u> as <u>effectively</u> as normal.
- 3) Severe valve damage can be treated by <u>replacing</u> the valve. Replacement valves can be ones taken from <u>humans</u> or <u>other mammals</u> (e.g. cows or pigs) these are <u>biological valves</u>. Or they can be <u>man-made</u> these are <u>mechanical valves</u>.
- Replacing a valve is a much less drastic procedure than a whole heart transplant.
 But fitting artificial valves is still major surgery and there can still be problems with blood clots.

Artificial Blood Can Keep You Alive In An Emergency

- 1) When someone loses a lot of blood, e.g. in an accident, their heart can still pump the remaining red blood cells around (to get oxygen to their organs), as long as the volume of their blood can be topped up.
- 2) Artificial blood is a blood substitute, e.g. a salt solution ("saline"), which is used to replace the lost volume of blood. It's safe (if no air bubbles get into the blood) and can keep people alive even if they lose % of their red blood cells. This may give the patient enough time to produce new blood cells. If not, the patient will need a blood transfusion.
- 3) Ideally, an artificial blood product would <u>replace</u> the function of the lost <u>red blood cells</u>, so that there's <u>no need</u> for a blood transfusion. Scientists are currently working on products that can do this.

Pity they can't fit me an artificial brain before the exam...

Make sure you know about the consequences of faulty heart valves or heart failure, as well as the advantages and disadvantages of the treatments on this page. Obviously if someone is really ill, it's unlikely that they'd turn down an artificial heart, artificial blood or a valve replacement — but these treatments aren't perfect.

Q1 a) Describe how faulty heart valves can lead to poor blood circulation. [2 marks] b) Suggest how severe damage to a heart valve can be treated. [1 mark]

O2 Suggest one disadvantage of treating coronary heart disease with an artificial heart.

Health and Disease

There's not a great deal about <u>diseases</u> and <u>health problems</u> that you can laugh about, so excuse me if this page is a bit dull. It's important stuff to know about though, so you'd best get cracking.

Diseases are a Major Cause of Ill Health

Health is the state of physical and mental wellbeing. Diseases are often responsible for causing ill health.

Diseases Can be Communicable or Non-Communicable

- Communicable diseases are those that can spread from person to person or between animals and people. They can be caused by things like bacteria, viruses, parasites and fungi. They're sometimes described as contagious or infectious diseases. Measles and malaria are examples of communicable diseases. There's more about them on pages 46-48.
- 2) Non-communicable diseases are those that <u>cannot spread</u> between people or between animals and people. They generally last for a <u>long time</u> and <u>get worse slowly</u>. <u>Asthma, cancer</u> and <u>coronary heart disease</u> (see page 37) are examples of non-communicable diseases.

Different Types of Disease Sometimes Interact

Sometimes diseases can interact and cause other physical and mental health issues that don't immediately seem related. Here are a few examples:

Pathogen is just the fancy term for a microorganism that can cause a disease when it infects its host.

- 1) People who have problems with their immune system (the system that your body uses to help fight off infection see p.49) have an increased chance of suffering from communicable diseases such as influenza (flu), because their body is less likely to be able to defend itself against the pathogen that causes the disease.
- 2) Some types of <u>cancer</u> can be triggered by <u>infection</u> by certain <u>viruses</u>. For example, infection with some types of <u>hepatitis virus</u> can cause long-term infections in the liver, where the virus lives in the cells. This can lead to an <u>increased chance</u> of developing <u>liver cancer</u>. Another example is infection with <u>HPV</u> (human papilloma virus), which can cause <u>cervical cancer</u> in women.
- 3) Immune system reactions in the body caused by infection by a pathogen can sometimes trigger allergic reactions such as skin rashes or worsen the symptoms of asthma for asthma sufferers.
- 4) Mental health issues such as depression can be triggered when someone is suffering from severe physical health problems, particularly if they have an impact on the person's ability to carry out everyday activities or if they affect the person's life expectancy.

Other Factors Can Also Affect Your Health

There are plenty of factors other than diseases that can also affect your health. For example:

- Whether or not you have a good, <u>balanced diet</u> that provides your body with <u>everything</u> it needs, and in the <u>right amounts</u>. A poor diet can affect your physical and mental health.
- 2) The stress you are under being constantly under lots of stress can lead to health issues.
- 3) Your <u>life situation</u> for example, whether you have easy <u>access</u> to <u>medicines</u> to treat illness, or whether you have <u>access</u> to things that can <u>prevent</u> you from <u>getting ill</u> in the first place, e.g. being able to buy <u>healthy food</u> or access <u>condoms</u> to prevent the <u>transmission</u> of some sexually transmitted diseases.

If stress can affect your health, why do we have exams...

You really need to get the terms communicable and non-communicable disease into your head. They could come up in the exam and you'd be really sad if you didn't understand the question.

Q1 What is meant by 'health'?

[1 mark]

O2 Why is influenza classed as a communicable disease?

Risk Factors for Non-Communicable Diseases

You've probably heard the term 'risk factor' before. This page has all the info you need to know about them.

Risk Factors Increase Your Chance of Getting a Disease

- Risk factors are things that are linked to an increase in the likelihood that a person will develop a
 certain disease during their lifetime. They don't guarantee that someone will get the disease.
- 2) Risk factors are often aspects of a person's <u>lifestyle</u> (e.g. how much exercise they do). They can also be the presence of certain substances in the <u>environment</u> (e.g. air pollution) or <u>substances</u> in your <u>body</u> (e.g. asbestos fibres asbestos was a material used in buildings until it was realised that the fibres could build up in your airways and cause diseases such as cancer later in life).
- Many non-communicable diseases are caused by several different risk factors interacting with each other rather than one factor alone.
- 4) Lifestyle factors can have different impacts locally, nationally and globally.

E.g. in <u>developed countries</u>, non-communicable diseases are <u>more common</u> as people generally have a <u>higher income</u> and can buy <u>high-fat</u> food. <u>Nationally</u>, people from <u>deprived areas</u> are <u>more likely</u> to smoke, have a poor diet and not exercise. This means the incidence of <u>cardiovascular disease</u>, <u>obesity</u> and <u>Type 2 diabetes is higher</u> in those areas. Your <u>individual choices affect the local incidence of disease</u>.

Some Risk Factors Can Cause a Disease Directly

- 1) Some risk factors are able to directly cause a disease. For example:
 - Smoking has been proven to directly cause <u>cardiovascular disease</u>, <u>lung disease</u> and <u>lung cancer</u>.
 It damages the walls of arteries and the cells in the lining of the lungs.
 - 2) It's thought that obesity can directly cause <u>Type 2 diabetes</u> by making the body <u>less sensitive</u> or <u>resistant</u> to insulin, meaning that it struggles to <u>control</u> the <u>concentration</u> of <u>glucose</u> in the blood.
 - 3) Drinking too much alcohol has been shown to cause liver disease. Too much alcohol can affect brain function too. It can damage the nerve cells in the brain, causing the brain to lose volume.
 - Smoking when pregnant can cause <u>lots</u> of <u>health problems</u> for the <u>unborn baby</u>.
 Drinking <u>alcohol</u> has similar effects.
 - 5) Cancer can be directly caused by <u>exposure</u> to certain <u>substances</u> or <u>radiation</u>. Things that cause cancer are known as <u>carcinogens</u>. <u>lonising radiation</u> (e.g. from X-rays) is an example of a <u>carcinogen</u>.
- 2) However, risk factors are identified by scientists looking for <u>correlations</u> in data, and <u>correlation doesn't always equal cause</u> (see p.9). Some risk factors <u>aren't</u> capable of directly causing a disease. For example, a <u>lack of exercise</u> and a <u>high fat diet</u> are heavily linked to an increased chance of <u>cardiovascular disease</u>, but they can't cause the disease directly. It's the resulting <u>high blood pressure</u> and <u>high 'bad' cholesterol levels</u> (see p.37) that can <u>actually cause</u> it.

Non-Communicable Diseases Can Be Costly

- The <u>HUMAN</u> cost of non-communicable diseases is obvious. <u>Tens of millions</u> of people around the world die from non-communicable diseases per year. People with these diseases may have a <u>lower quality of</u> <u>life</u> or a <u>shorter lifespan</u>. This not only affects the <u>sufferers</u> themselves, but their <u>loved ones</u> too.
- 2) It's also important to think about the <u>FINANCIAL</u> cost. The cost to the NH9 of <u>researching</u> and <u>treating</u> these diseases is <u>huge</u> and it's the same for other <u>health services</u> and <u>organisations</u> around the world. <u>Families</u> may have to <u>move</u> or <u>adapt their home</u> to help a family member with a disease, which can be <u>costly</u>. Also, if the family member with the disease has to <u>give up work</u> or <u>dies</u>, the family's <u>income</u> will be <u>reduced</u>. A <u>reduction</u> in the number of people <u>able</u> to <u>work</u> can also affect a <u>country's economy</u>.

Best put down that cake and go for a run...

You might be asked to interpret data about risk factors. See p.9 for a few tips on what you can and can't say.

Of the control of the

Cancer

Cancer's <u>not</u> a pleasant topic, but the more we understand about it, the better our chances of <u>avoiding</u> and <u>beating</u> it (and getting good <u>marks</u> in the exam). You're a good way through the topic, so keep going.

Cancer is Caused by Uncontrolled Cell Growth and Division

This <u>uncontrolled</u> growth and division is a result of <u>changes</u> that occur to the <u>cells</u> and results in the formation of a <u>tumour</u> (a mass of cells). Not all tumours are cancerous. They can be <u>benign</u> or <u>malignant</u>:

- Benign This is where the tumour grows until there's no more room. The tumour stays
 in one place (usually within a membrane) rather than invading other tissues in the body.
 This type isn't normally dangerous, and the tumour isn't cancerous.
- 2) <u>Malignant</u> This is where the tumour grows and <u>spreads</u> to neighbouring healthy tissues. Cells can <u>break off</u> and spread to other parts of the body by travelling in the <u>bloodstream</u>. The malignant cells then <u>invade</u> healthy tissues elsewhere in the body and form <u>secondary tumours</u>. Malignant tumours are <u>dangerous</u> and can be fatal <u>they are cancers</u>.

Risk Factors Can Increase the Chance of Some Cancers

Anyone can develop cancer. Having risk factors doesn't mean that you'll definitely get cancer. It just means that you're at an increased risk of developing the disease. Cancer survival rates have increased due to medical advances such as improved treatment, being able to diagnose cancer earlier and increased screening for the disease.

Risk Factors Can Be Associated With Lifestyle

Scientists have identified lots of lifestyle risk factors for various types of cancer. For example:

 Smoking — It's a well known fact that smoking is linked to <u>lung cancer</u>, but research has also linked it to <u>other types</u> of cancer too, including mouth, bowel, stomach and cervical cancer.



- Obesity Obesity has been linked to many different cancers, including bowel, liver
 and kidney cancer. It's the second biggest preventable cause of cancer after smoking.
- 3) <u>UV exposure</u> People who are often exposed to <u>UV radiation</u> from the Sun have an increased chance of developing <u>skin cancer</u>. People who live in <u>sunny climates</u> and people who spend a lot of time <u>outside</u> are at <u>higher risk</u> of the disease. People who frequently use sun beds are also putting themselves at higher risk of developing skin cancer.
- 4) <u>Viral infection</u> Infection with some viruses has been shown to <u>increase</u> the chances of developing <u>certain types</u> of cancer. For example, infection with <u>hepatitis B</u> and <u>hepatitis C</u> viruses can increase the risk of developing <u>liver cancer</u>. The likelihood of becoming infected with these viruses sometimes depends on lifestyle e.g. they can be spread between people through <u>unprotected sex</u> or <u>sharing needles</u>.

Risk Factors Can Also Be Associated With Genetics

- 1) Sometimes you can inherit faulty genes that make you more susceptible to cancer.
- For example, mutations (changes) in the <u>BRCA</u> genes have been linked to an <u>increased likelihood</u> of developing <u>breast</u> and <u>ovarian cancer</u>.

At least our rubbish summers reduce our UV exposure...

Joking aside, UV radiation can still reach us through the clouds, and like many other lifestyle risk factors, we can take steps to reduce the risk, e.g. by keeping covered up outside and wearing sun block.

O1 What are tumours the result of? [1 mark]

O2 List three lifestyle factors that can increase the risk of developing cancer.

[3 marks]

Plant Cell Organisation

You saw on page 27 how animals keep their specialised cells neat and tidy — plants are in on the act too.

Plant Cells Are Organised Into Tissues And Organs

<u>Plants</u> are made of <u>organs</u> like <u>stems</u>, <u>roots</u> and <u>leaves</u>. Plant organs work together to make <u>organ systems</u>. These can perform the various tasks that a plant needs to carry out to survive and grow — for example, <u>transporting substances</u> around the plant. Plant organs are made of <u>tissues</u>. Examples of plant tissues are:


- 1) Epidermal tissue this covers the whole plant.
- 2) Palisade mesophyll tissue this is the part of the leaf where most photosynthesis happens.
- Spongy mesophyll tissue this is also in the leaf, and contains big <u>air spaces</u> to allow gases to <u>diffuse</u> in and out of cells.
- 4) <u>Xylem</u> and <u>phloem</u> they <u>transport</u> things like <u>water</u>, <u>mineral ions</u> and <u>food</u> around the plant (through the roots, stems and leaves see next page for more).
- 5) Meristem tissue this is found at the growing tips of shoots and roots and is able to differentiate (change) into lots of different tupes of plant cell, allowing the plant to grow.





The Leaf is an Organ Made Up of Several Types of Tissue

Leaves contain epidermal, mesophyll, xylem and phloem tissues.



You need to know how the structures of the tissues that make up the leaf are related to their function:

- 1) The epidermal tissues are covered with a waxy cuticle, which helps to reduce water loss by evaporation.
- 2) The upper epidermis is transparent so that light can pass through it to the palisade layer.
- 3) The <u>palisade layer</u> has lots of <u>chloroplasts</u> (the little structures where photosynthesis takes place). This means that they're near the top of the leaf where they can get the most light.
- 4) The xylem and phloem form a network of vascular bundles, which deliver water and other <u>nutrients</u> to the entire leaf and take away the <u>glucose</u> produced by photosynthesis. They also help <u>support</u> the structure.
- 5) The tissues of leaves are also adapted for efficient gas exchange (see page 25). E.g. the lower epidermis is full of little holes called stomata, which let CO₂ diffuse directly into the leaf. The opening and closing of stomata is controlled by guard cells in response to environmental conditions. The air spaces in the spongu mesophull tissue increase the rate of diffusion of gases.

Plant cell organisation — millions of members worldwide...

There are a lot of weird names here, so make sure you spend plenty of time on this page. Maybe you could draw your own leaf diagram and label it with descriptions of the different tissue types. It would make an excellent Christmas present for someone, or an art collector might even want it.

Describe the characteristics of meristem tissue.

[2 marks]

Transpiration and Translocation

You might be surprised to learn that there aren't tiny trucks that transport substances around plants. Then again, you might not be — either way, you need to learn the stuff on this page...



Phloem Tubes Transport Food:

- Made of columns of <u>elongated</u> living cells with small <u>pores</u> in the end walls to allow cell sap to flow through.
- They transport food substances (mainly dissolved sugars)
 made in the leaves to the rest of the plant for immediate
 use (e.g. in growing regions) or for storage.
- 3) The transport goes in both directions.
- 4) This process is called <u>translocation</u>.






Xylem Tubes Take Water Up:

- Made of dead cells joined end to end with no end walls between them and a hole down the middle. They're strengthened with a material called lignin.
- They carry <u>water</u> and <u>mineral</u> ions from the <u>roots</u> to the <u>stem</u> and <u>leaves</u>.
- The movement of water from the roots, through the xylem and out of the leaves is called the transpiration stream (see below).

Transpiration is the Loss of Water from the Plant



- Transpiration is caused by the <u>evaporation</u> and <u>diffusion</u> (see page 20) of water from a plant's surface. Most transpiration happens at the <u>leaves</u>.
- This evaporation creates a slight <u>shortage</u> of water in the leaf, and so more water is drawn up from the rest of the plant through the <u>xulem vessels</u> to replace it.
- This in turn means more water is drawn up from the <u>roots</u>, and so there's a constant <u>transpiration</u> stream of water through the plant.

Head back to page 22 = to see how root hair = cells are adapted for taking up water.

Transpiration is just a side-effect of the way leaves are adapted for photosynthesis.

They have to have stomata in them so that gases can be exchanged easily (see page 25). Because there's more water inside the plant than in the air outside, the water escapes from the leaves through the stomata by diffusion.

Don't let revision stress you out — just go with the phloem...

Phloem transports substances in both directions, but xylem only transports things upwards — xy to the sky.

O1 Describe the structure of xvlem.

[3 marks]

Transpiration and Stomata

Sorry, more on transpiration. But then it's a quick dash through stomata and out of the other end of the topic.

Transpiration Rate is Affected by Four Main Things

- LIGHT INTENSITY the brighter the light, the greater the transpiration rate. Stomata begin to close as it gets darker. Photosynthesis can't happen in the dark, so they don't need to be open to let co. in. When the stomata are closed, very little water can escape.
- 2) TEMPERATURE the warmer it is, the faster transpiration happens.


When it's warm the water particles have more energy to evaporate and diffuse out of the stomata.

- 3) AIR FLOW the better the air flow around a leaf (e.g. stronger wind), the greater the transpiration rate. If air flow around a leaf is poor, the water vapour just surrounds the leaf and doesn't move away. This means there's a high concentration of water particles outside the leaf as well as inside it, so diffusion doesn't happen as quicklu. If there's good air flow, the water vapour is swept away, maintaining a low concentration of water in the air outside the leaf. Diffusion then happens quickly, from an area of higher concentration to an area of lower concentration.
- 4) HUMIDITY the drier the air around a leaf, the faster transpiration happens.

This is like what happens with air flow. If the air is humid there's a lot of water in it already, so there's not much of a difference between the inside and the outside of the leaf. Diffusion happens fastest if there's a really high concentration in one place, and a really low concentration in the other.

You can estimate the rate of transpiration by measuring the uptake of water by a plant. This is because you can assume that water uptake by the plant is directly related to water loss by the leaves (transpiration).

Set up the apparatus as in the diagram, and then record the starting position of the air bubble. Start a stopwatch and record the distance moved by the bubble per unit time, e.g. per hour. Keep the conditions constant throughout the experiment, e.g. the temperature and air humidity.



Guard Cells Are Adapted to Open and Close Stomata



- They have a kidney shape which opens and closes the stomata (page 25) in a leaf.
- 2) When the plant has lots of water the guard cells fill with it and go plump and turgid. This makes the stomata open so gases can be exchanged for photosunthesis.
- 3) When the plant is short of water, the guard cells lose water and become flaccid, making the stomata close. This helps stop too much water vapour escaping.
- 4) Thin outer walls and thickened inner walls make the opening and closing work.
- 5) They're also sensitive to light and close at night to save water without losing out on photosynthesis.
- 6) You usually find more stomata on the undersides of leaves than on the top. The lower surface is shaded and cooler — so less water is lost through the stomata than if they were on the upper surface.
- 7) Guard cells are therefore adapted for gas exchange and controlling water loss within a leaf.

I say stomaaarta, you say stomaaayta...

Different leaves will have different distributions of stomata. You can peel the epidermal tissue off some leaves and mount them on microscope slides (see page 13) to compare them. It's thrilling stuff.



01 Aloe vera plants grow in hot, dry areas. Primroses grow in cool, wet areas. Predict which plant will have fewer stomata per cm² on the underside of its leaves. Explain your answer. [2 marks]

Revision Questions for Topic 2

Well, that's Topic 2 finished. Now it's time for the greatest quiz ever.

Try these questions and tick off each one when you get it right.

When you're completely happy with a sub-topic, tick it off.

Cell Organisation (p.27)

For even more practice, try the
Retrieval Quiz for Topic 2 — just
scan this QR code!



What is a tissue? 2) Explain what is meant by the term 'organ system'. The Role of Enzymes and Food Tests (p.28-32) 3) Why can enzymes be described as biological catalysts? 4) Why do enzymes only usually catalyse one reaction? 5) What does it mean when an enzyme has been 'denatured'? 6) Describe how you could investigate the effect of pH on the rate of amulase activity. 7) List the three places where amylase is made in the human body. 8) What is the role of lipases? 9) Where is bile stored? 10) Name the solution that you would use to test for the presence of lipids in a food sample. The Lungs and Circulatory System (p.33-36) 11) Name the tubes that split off the trachea. 12) Explain the role that alveoli play in gas exchange. 13) Explain why the circulatory system in humans is described as a 'double circulatory system'. 14) Why does the heart have valves? 15) Name the four chambers of the heart. 16) How is the resting heart rate controlled in a healthy heart? 17) How are arteries adapted to carry blood away from the heart? 18) Why do red blood cells not have a nucleus? Diseases and Risk Factors (p.37-41) 19) Give two advantages and two disadvantages of statins. 20) What is the difference between biological and mechanical replacement heart valves? 21) What is meant by a non-communicable disease? 22) Give an example of where different types of disease might interact in the body. 23) What is meant by a risk factor of a disease? 24) Which type of tumour is cancerous? Plant Cell Organisation and Transport (p.42-44) 25) List the tissues that make up a leaf. 26) Explain how the structure of the upper epidermal tissue in a leaf is related to its function. 27) What is the function of phloem? 28) What is transpiration? 29) List the four main things that affect transpiration. 30) How could you measure the rate of transpiration? 31) Name the type of cell that helps open and close stomata.

Communicable Disease

If you're hoping I'll ease you gently into this new topic... no such luck. Straight on to the baddies of biology.

There Are Several Types of Pathogen

- 1) Pathogens are microorganisms that enter the body and cause disease.
- 2) They cause communicable (infectious) diseases diseases that can easily spread (see p.39).
- 3) Both plants and animals can be infected by pathogens.

1. Bacteria Are Very Small Living Cells

- Bacteria are very small cells (about 1/100th the size of your body cells), which can reproduce rapidly inside your body.
- 2) They can make you feel ill by producing toxins (poisons) that damage your cells and tissues.

2. Viruses Are Not Cells — They're Much Smaller

- 1) Viruses are not cells. They're tiny, about 1/100th the size of a bacterium.
- 2) Like bacteria, they can reproduce rapidly inside your body.
- They live inside your cells and replicate themselves using the cells' machinery to produce many copies of themselves.
 The cell will usually then burst, releasing all the new viruses.
- 4) This cell damage is what makes you feel ill.

3. Protists are Single-Celled Eukaryotes

- There are lots of different types of protists. But they're all <u>eukaryotes</u> (see page 11) and most of them are <u>single-celled</u>.
- Some protists are <u>parasites</u>. Parasites live <u>on</u> or <u>inside</u> other organisms and can cause them <u>damage</u>.
 They are often transferred to the organism by a <u>vector</u>, which doesn't get the disease itself
 — e.g. an insect that carries the protist.

4. Fungi Come in Different Shapes

- 1) Some fungi are single-celled. Others have a body which is made up of hyphae (thread-like structures).
- 2) These hyphae can grow and penetrate human skin and the surface of plants, causing diseases.
- 3) The huphae can produce spores, which can be spread to other plants and animals.

Pathogens Can Be Spread in Different Ways

Pathogens can be spread in many ways. Here are a few that you need to know about.

- WATER Some pathogens can be picked up by drinking or bathing in <u>dirty water</u>. E.g. <u>cholera</u> is a
 <u>bacterial infection</u> that's spread by <u>drinking</u> water <u>contaminated</u> with the diarrhoea of other sufferers.
- 2) AIR Pathogens can be carried in the air and can then be breathed in. Some airborne pathogens are carried in the air in droplets produced when you cough or sneeze e.g. the influenza virus that causes flu is spread this way.
- 3) <u>DIRECT CONTACT</u> Some pathogens can be picked up by <u>touching</u> contaminated surfaces, including the <u>skin</u>. E.g. <u>athlete's foot</u> is a <u>fungus</u> which makes skin itch and flake off. It's most commonly spread by touching the same things as an infected person, e.g. <u>shower floors</u> and <u>towels</u>.


Hooray, I've avoided the classic 'he was a fungi to be with' joke...

Yuck, lots of nasties out there that can cause disease. Plants need to be worried too, as you'll find out.

Describe how viruses cause cell damage.

[2 marks]

A body cel



Viral, Fungal and Protist Diseases

There are heaps of diseases caused by viruses, fungi and protists, but you just need to know about these ones.

You Need to Know About Three Viral Diseases...

- 1) Measles is a viral disease. It is spread by droplets from an infected person's sneeze or cough.
- 2) People with measles develop a red skin rash, and they'll show signs of a fever (a high temperature).
- 3) Measles can be very serious, or even fatal, if there are complications. For example, measles can sometimes lead to pneumonia (a lung infection) or inflammation of the brain (encephalitis).
- 4) Most people are vaccinated against measles when they're young.
- HIV is a <u>virus</u> spread by <u>sexual contact</u>, or by exchanging <u>bodily fluids</u> such as blood.
 This can happen when people <u>share needles</u> when taking drugs.
- 2) HIV initially causes <u>flu-like symptoms</u> for a few weeks. Usually, the person doesn't then experience any symptoms for several years. During this time, HIV can be controlled with <u>antiretroviral drugs</u>. These stop the virus <u>replicating</u> in the body.
- 3) The virus attacks the immune cells (see page 49).
- 4) If the body's immune system is badly damaged, it <u>can't cope</u> with <u>other infections</u> or <u>cancers</u>. At this stage, the virus is known as <u>late stage HIV infection</u>, or <u>AIDS</u>.
- 1) Tobacco mosaic virus (IMV) is a virus that affects many species of plants, e.g. tomatoes.
- 2) It causes a mosaic pattern on the leaves of the plants parts of the leaves become discoloured.
- 3) The discolouration means the plant can't carry out photosynthesis as well, so the virus affects growth.

...a Fungal Disease...

1) Rose black spot is a <u>fungus</u> that causes <u>purple or black spots</u> to develop on the <u>leaves</u> of <u>rose plants</u>. Who'd have guessed. The leaves can then turn <u>uellow</u> and <u>drop off</u>.

2) This means that less photosynthesis can happen, so the plant doesn't grow very well.

3) It spreads through the environment in water or by the wind.

4) Gardeners can treat the disease using <u>fungicides</u> and by <u>stripping</u> the plant of its <u>affected leaves</u>. These leaves then need to be <u>destroyed</u> so that the fungus can't spread to other rose plants.

Photosynthesis is important for plant growth because it produces glucose — see page 57.



...and a Disease Caused by a Protist

- 1) Malaria is caused by a profist (see the previous page).
- Part of the malarial protist's life cycle takes place inside the mosquito. The mosquitoes are vectors
 (see the previous page) they pick up the malarial protist when they feed on an infected animal.
- Every time the mosquito feeds on another animal, it <u>infects it</u> by inserting the protist into the animal's blood vessels.
- 4) Malaria causes repeating episodes of fever. It can be fatal.
- 5) The spread of malaria can be reduced by stopping the mosquitoes from breeding.
- 6) People can be protected from mosquitoes using insecticides and mosquito nets.

I've heard this page has gone viral...

The examiner could grill you on any one of these diseases, so make sure you know them all inside out.

Q1 What symptom of measles is shown on the skin?

[1 mark]

O2 How can rose black spot be treated so that it doesn't spread to other plants?

[2 marks]

Bacterial Diseases and Preventing Disease

Sorry — I'm afraid there are some more diseases to learn about here. This time, they're diseases caused by bacteria. I don't know about you, but I'm starting to feel a bit itchy all over...

You Need to Know About Two Bacterial Diseases

- 1) Salmonella is a type of bacteria that causes food poisoning.
- 2) Infected people can suffer from fever, stomach cramps, vomiting and diarrhoea. Pleasant.
- 3) These symptoms are caused by the toxins that the bacteria produce (see page 46).
- 4) You can get Salmonella food poisoning by eating food that's been contaminated with Salmonella bacteria, e.g. eating chicken that caught the disease whilst it was alive, or eating food that has been contaminated by being prepared in unhygienic conditions.
- In the UK, most poultry (e.g. chickens and turkeys) is given a vaccination against Salmonella. This is to control the spread of the disease.
- 1) Gonorrhoea is a sexually transmitted disease (STD).
- 2) STDs are passed on by sexual contact, e.g. having unprotected sex.
- 3) Gonorrhoea is caused by bacteria.
- 4) A person with gonorrhoea will get <u>pain</u> when they <u>urinate</u>. Another symptom is a thick yellow or green <u>discharge</u> from the <u>vagina</u> or the <u>penis</u>.
- 5) Gonorrhoea was originally treated with an antibiotic called penicillin, but this has become trickier now because strains of the bacteria have become resistant to it (see page 51).
- 6) To prevent the <u>spread</u> of gonorrhoea, people can be treated with <u>antibiotics</u> and should use barrier methods of contraception (see page 78), such as condoms.

The Spread of Disease Can Be Reduced or Prevented

There are things that we can do to reduce, and even prevent, the spread of disease. For example:

- Being hygienic Using simple hygiene measures can prevent the spread of disease.
 For example, doing things like washing your hands thoroughly before preparing food or after you've sneezed can stop you infecting another person.
- Destroying vectors By getting rid of the organisms that spread disease, you can
 prevent the disease from being passed on. Vectors that are insects can be killed
 using insecticides or by destroying their habitat so that they can no longer breed.
- 3) <u>Isolating infected individuals</u> If you isolate someone who has a communicable disease, it <u>prevents</u> them from <u>passing it on</u> to anyone else.
- 4) <u>Vaccination</u> Vaccinating people and animals against communicable diseases means that they are <u>less likely</u> to develop the infection and then <u>pass it on</u> to someone else. There's more about how vaccination works on page 50.

The spread of disease — mouldy margarine...

OK, I promise, that's it. No more diseases to learn about in this Topic. You may be sick of them already (geddit?) but don't turn this page until you've got all the facts firmly attached to your cranial material.

Q1 What has made it harder to treat gonorrhoea? [1 mark]

O2 It is important for chefs to wash their hands thoroughly before cooking. Suggest why.

Fighting Disease

The human body has some pretty neat features when it comes to fighting disease.

Your Body Has a Pretty Sophisticated Defence System


- 1) The human body has got features that stop a lot of nasties getting inside in the first place.
- 2) The skin acts as a barrier to pathogens. It also secretes antimicrobial substances which kill pathogens.
- 3) Hairs and mucus in your nose trap particles that could contain pathogens.
- 4) The trachea and bronchi (breathing pipework see page 33) secrete mucus to trap pathogens.
- 5) The trachea and bronchi are lined with <u>cilia</u>. These are hair-like structures, which <u>waft the mucus</u> up to the back of the throat where it can be <u>swallowed</u>.
- 6) The stomach produces hydrochloric acid. This kills pathogens that make it that far from the mouth.

Your Immune System Can Attack Pathogens

- 1) If pathogens do make it into your body, your immune system kicks in to destroy them.
- 2) The most important part of your immune system is the <u>white blood cells</u>. They travel around in your blood and crawl into every part of you, constantly patrolling for microbes. When they come across an invading microbe they have three lines of attack.


1. Consuming Them

White blood cells can engulf foreign cells and digest them. This is called phagocytosis.



2. Producing Antibodies

- 1) Every invading pathogen has unique molecules (called antigens) on its surface.
- 2) When some types of white blood cell come across a foreign antigen (i.e. one they don't recognise), they will start to produce proteins called antibodies to lock onto the invading cells so that they can be found and destroyed by other white blood cells. The antibodies produced are specific to that type of antigen they won't lock on to any others.
- 3) Antibodies are then produced rapidly and carried around the body to find all similar bacteria or viruses.
- 4) If the person is infected with the same pathogen again the white blood cells will rapidly produce the antibodies to kill it — the person is <u>naturally</u> immune to that pathogen and won't get ill.



3. Producing Antitoxins

These counteract toxins produced by the invading bacteria.

Fight disease — blow your nose with boxing gloves...

If you have a low level of white blood cells, you'll be more susceptible to infections. HIV attacks white blood cells and weakens the immune system, making it easier for other pathogens to invade.

Q1 What is phagocytosis?

[1 mark]

O2 How are the trachea and the bronchi adapted to defend against the entry of pathogens?


[3 marks]

Fighting Disease — Vaccination

Vaccinations have changed the way we fight disease. We don't always have to deal with the problem once it's happened — we can prevent it happening in the first place.

Vaccination can Protect from Future Infections

- When you're infected with a new <u>pathogen</u>, it takes your white blood cells a few days to <u>learn</u> how to deal with it. But by that time, you can be pretty ill.
- 2) <u>Vaccinations</u> usually involve injecting small amounts of <u>dead</u> or <u>inactive</u> pathogens. These carry <u>antigens</u>, which cause your body to produce <u>antibodies</u> to attack them even though the pathogen is <u>harmless</u> (since it's dead or inactive). For example, the MMR vaccine contains <u>weakened</u> versions of the viruses that cause <u>measles</u>, <u>mumps</u> and <u>rubella</u> (German measles) all in one vaccine.
- But if live pathogens of the same type appear after that, the white blood cells can rapidly
 mass-produce antibodies to kill off the pathogen. Cool.



There are Pros and Cons of Vaccination

PROS

- Vaccines have helped <u>control</u> lots of communicable diseases that were once <u>common</u> in the UK (e.g. polio, measles, whooping cough, rubella, mumps, tetanus...). <u>Smallbox</u> no longer occurs at all, and <u>polio</u> infections have fallen by 99%.
- 2) Big outbreaks of disease called epidemics can be prevented if a large percentage of the population is vaccinated. That way, even the people who aren't vaccinated are unlikely to catch the disease because there are fewer people able to pass it on. But if a significant number of people aren't vaccinated, the disease can spread quickly through them and lots of people will be ill at the same time.

CONS

- 1) Vaccines don't always work sometimes they don't give you immunitu.
- You can sometimes have a <u>bad reaction</u> to a vaccine (e.g. swelling, or maybe something more serious like a fever or seizures). But bad reactions are very rare.

Prevention is better than cure...

Deciding whether to have a vaccination means balancing risks — the risk of catching the disease if you don't have a vaccine, against the risk of having a bad reaction if you do. As always, you need to look at the evidence. For example, if you get measles (the disease), there's about a 1 in 15 chance that you'll get complications (e.g. pneumonia) — and about 1 in 500 people who get measles actually die. However, the number of people who have a problem with the vaccine is more like 1 in 1 000 000.

Q1 Basia is vaccinated against flu and Cassian isn't. They are both exposed to a flu virus.

Cassian falls ill whereas Basia doesn't. Explain why.



Fighting Disease — Drugs

...a biscuit, nurse? Thanks very much. Sorry, couldn't face that last page — I'm squeamish about needles.*

Some Drugs Relieve Symptoms — Others Cure the Problem

- Painkillers (e.g. aspirin) are drugs that relieve pain (no, really). However, they don't actually
 tackle the <u>cause</u> of the disease or <u>kill</u> pathogens, they just help to reduce the <u>symptoms</u>.
- Other drugs do a similar kind of thing reduce the <u>symptoms</u> without tackling the underlying <u>cause</u>. For example, lots of "cold remedies" don't actually <u>cure</u> colds.
- 3) Antibiotics (e.g. penicillin) work differently they actually kill (or prevent the growth of) the bacteria causing the problem without killing your own body cells. <u>Different antibiotics</u> kill <u>different types</u> of bacteria, so it's important to be treated with the <u>right one</u>.
- 4) But antibiotics don't destroy viruses (e.g. flu or cold viruses). Viruses reproduce using your body cells, which makes it very difficult to develop drugs that destroy just the virus without killing the body's cells.
- The use of antibiotics has greatly reduced the number of deaths from communicable diseases caused by bacteria.

Bacteria Can Become Resistant to Antibiotics

- 1) Bacteria can mutate sometimes the mutations cause them to be resistant to (not killed by) an antibiotic.
- 2) If you have an infection, some of the bacteria might be resistant to antibiotics.
- 3) This means that when you treat the infection, only the non-resistant strains of bacteria will be killed.
- 4) The individual resistant bacteria will survive and reproduce, and the population of the resistant strain will increase. This is an example of natural selection (see page 96).
- 5) This resistant strain could cause a <u>serious infection</u> that <u>can't</u> be treated by antibiotics. E.g. <u>MRSA</u> (meticillin-resistant <u>Staphylococcus aureus</u>) causes serious wound infections and is resistant to the powerful antibiotic <u>meticillin</u>.
- 6) To slow down the <u>rate</u> of development of <u>resistant strains</u>, it's important for doctors to <u>avoid over-prescribing</u> antibiotics. So you <u>won't</u> get them for a <u>sore throat</u>, only for something more serious.
- 7) It's also important that you finish the whole course of antibiotics and don't just stop once you feel better.

Many Drugs Originally Came From Plants

- 1) Plants produce a variety of chemicals to defend themselves against pests and pathogens (see page 55).
- Some of these chemicals can be used as drugs to treat human diseases or relieve symptoms.
 A lot of our current medicines were discovered by studying plants used in traditional cures. For example:
 - Aspirin is used as a painkiller and to lower fever. It was developed from a chemical found in willow.
 - <u>Digitalis</u> is used to treat <u>heart conditions</u>. It was developed from a chemical found in <u>foxgloves</u>.
- 3) Some drugs were extracted from microorganisms. For example:
 - Alexander Fleming was clearing out some Petri dishes containing bacteria. He noticed that one of
 the dishes of bacteria also had mould on it and the area around the mould was free of the bacteria.
 - He found that the mould (called Penicillium notatum) on the Petri dish was
 producing a <u>substance</u> that <u>killed the bacteria</u> this substance was <u>penicillin</u>.
- 4) These days, drugs are made on a large scale in the <u>pharmaceutical industry</u> they're synthesised by chemists in labs. However, the process still might start with a chemical <u>extracted</u> from a <u>plant</u>.

Ahh...Ahh... Ahhhhh Choooooooo — urghh, this page is catching...

Drug development is a big industry. And guess what — you're about to find out some more about it.

O1 Which type of pathogen can antibiotics be used to kill?

Developing Drugs

New drugs are constantly being developed. But before they can be given to the general public, they have to go through a thorough testing procedure. This is what usually happens...

There are Three Main Stages in Drug Testing

- 1
- 1) In preclinical testing, drugs are tested on human cells and tissues in the lab.
- 2) However, you can't use human cells and tissues to test drugs that affect whole or multiple body systems, e.g. testing a drug for blood pressure must be done on a whole animal because it has an intact circulatory system.



The next step in preclinical testing is to test the drug on live animals.
 This is to test efficacy (whether the drug works and produces the effect you're looking for), to find out about its toxicity (how harmful it is) and to find the best dosage (the concentration that should be given, and how often it should be given).



2) The law in Britain states that any new drug must be tested on two different live mammals. Some people think it's <u>cruel</u> to test on animals, but others believe this is the <u>safest</u> way to make sure a drug isn't dangerous before it's given to humans.





- 1) If the drug passes the tests on animals then it's tested on human volunteers in a clinical trial.
- First, the drug is tested on healthy volunteers. This is to make sure that it doesn't have any harmful side effects when the body is working normally. At the start of the trial, a very low dose of the drug is given and this is gradually increased.
- 3) If the results of the tests on healthy volunteers are good, the drugs can be tested on people suffering from the <u>illness</u>. The <u>optimum dose</u> is found this is the dose of drug that is the most effective and has few side effects.
- 4) To test how well the drug works, patients are <u>randomly</u> put into <u>two groups</u>. One is given the <u>new drug</u>, the other is given a <u>placebo</u> (a substance that's like the drug being tested but doesn't do anything). This is so the <u>doctor</u> can see the actual difference the drug makes it allows for the <u>placebo effect</u> (when the patient expects the treatment to work and so <u>feels better</u>, even though the treatment isn't doing anything).
- 5) Clinical trials are <u>blind</u>—the patient in the study <u>doesn't know</u> whether they're getting the drug or the placebo. In fact, they're often <u>double-blind</u>—neither the patient nor the <u>doctor</u> knows until all the <u>results</u> have been gathered. This is so the doctors <u>monitoring</u> the patients and <u>analysing</u> the results aren't <u>subconsciously influenced</u> by their knowledge.
- 6) The results of drug testing and drug trials aren't published until they've been through <u>peer review</u>. This helps to prevent false claims.

Peer review is when other scientists check that the work is valid and has been carried out rigorously — see page 1.

The placebo effect doesn't work with revision...

... you can't just expect to get a good mark and then magically get it. I know, I know, there's a lot of information to take in on this page, but just read it through slowly. There's nothing too tricky here — it's just a case of going over it again and again until you've got it all firmly lodged in your memory.

Q1 What is meant by the efficacy of a drug? [1 mark]

Q2 Why do clinical trials of a new drug begin with healthy volunteers? [1 mark]


Why must the results from drug testing be assessed by peer review? [1 mark]

Monoclonal Antibodies

Antibodies aren't only used by the immune system — scientists have engineered them for lots of new uses.


Monoclonal Antibodies are Identical Antibodies

- 1) Antibodies are produced by B-lymphocytes a type of white blood cell (see page 49).
- 2) Monoclonal antibodies are produced from lots of clones of a single white blood cell. This means all the antibodies are identical and will only target one specific protein antigen.
- 3) However, you can't just grab the lumphocute that made the antibody and grow more - lymphocytes don't divide very easily.
- 4) Tumour cells, on the other hand, don't produce antibodies but divide lots — so they can be grown really easily.
- 5) It's possible to fuse a mouse B-lumphocute with a tumour cell to create a cell called a hubridoma.
- 6) Hybridoma cells can be cloned to get lots of identical cells. These cells all produce the same antibodies (monoclonal antibodies). The antibodies can be collected and purified.
- 7) You can make monoclonal antibodies that bind to anything you want, e.g. an antigen that's only found on the surface of one type of cell. Monoclonal antibodies are really useful because they will only bind to (target) this molecule — this means you can use them to target a specific cell or chemical in the body.




It divides quickly to produce lots of clones that produce the monoclonal antibodies

Monoclonal Antibodies Are Used In Pregnancy Tests



If you're pregnant:



...and stick to strip

If you're not pregnant:



...but don't stick

A hormone called HCG is found in the urine of women only when they are pregnant. Pregnancy testing sticks detect this hormone. Here's how they work:

- 1) The bit of the stick you wee on has some antibodies to the hormone, with blue beads attached.
- 2) The test strip (the bit of the stick that turns blue if you're pregnant) has some more antibodies to the hormone stuck onto it (so that they can't move).

If you're pregnant and you wee on the stick:

- The hormone binds to the antibodies on the blue beads.
- The urine moves up the stick, carrying the hormone and the beads.
- The beads and hormone bind to the antibodies on the strip.
- So the blue beads get stuck on the strip, turning it blue.

4) If you're not pregnant and you wee on the stick, the urine still moves up the stick, carruing the blue beads. But there's nothing to stick the blue beads onto the test strip, so it doesn't go blue.

The one time when you can write "wee on a stick" in an exam...

There's more on monoclonal antibodies coming next, but don't move on until you understand this page.


01 What is a hybridoma cell made from? [2 marks]

More on Monoclonal Antibodies

Because monoclonal antibodies can be produced to target a specific chemical or cell, they have loads of uses.

Monoclonal Antibodies Can be Used to Treat Diseases...

- Different cells in the body have different antigens on their cell surface. So you can make
 monoclonal antibodies that will bind to specific cells in the body (e.g. just liver cells).
- Cancer cells have <u>antigens</u> on their <u>cell membranes</u> that <u>aren't</u> found on normal body cells.
 Theu're called tumour markers.
- In the lab, you can make monoclonal antibodies that will bind to these tumour markers.
- An <u>anti-cancer drug</u> can be attached to these monoclonal antibodies. This might be a <u>radioactive</u> substance, a <u>toxic drug</u> or a chemical which stops cancer cells <u>growing</u> and <u>dividing</u>.
- 5) The antibodies are given to the patient through a drip.
- 6) The antibodies target specific cells (the cancer cells) because they only bind to the tumour markers.
- The drug kills the cancer cells but doesn't kill any normal body cells near the tumour.



...and in Laboratories and Research to Find Specific Substances

Monoclonal antibodies can be used to:

- 1) Bind to hormones and other chemicals in blood to measure their levels.
- 2) Test blood samples in laboratories for certain pathogens.
- 3) Locate specific molecules on a cell or in a tissue:
 - 1) First, monoclonal antibodies are made that will bind to the specific molecules you're looking for.
 - 2) The antibodies are then bound to a fluorescent due.
 - If the molecules are present in the sample you're analysing, the monoclonal antibodies will attach to them, and they can be detected using the due.

Monoclonal Antibodies Do Have Some Problems Though

- 1) There are some obvious advantages of monoclonal antibodies. One big one is in cancer treatment. Other cancer treatments (like standard chemotherapy and radiotherapy) can affect normal body cells as well as killing cancer cells, whereas monoclonal antibodies target specific cells. This means the side effects of an antibody-based drug are lower than for standard chemotherapy or radiotherapy.
- 2) However, monoclonal antibodies do cause more side effects than were originally expected, e.g. they can cause fever, vomiting and low blood pressure. When they were first developed, scientists thought that because they targeted a very specific cell or molecule, they wouldn't create a lot of side effects.
- This means that they are <u>not as widely used</u> as treatments as scientists had originally thought they might be.

Bonoclonal antibodies — used to detect Irish rock bands...

As you've just found out, monoclonal antibodies are really useful for finding stuff and making it glow.

Q1 A scientist has a sample of tissue from a patient with a suspected bacterial infection. He sticks it to a microscope slide. Then he adds a solution of monoclonal antibodies against the bacteria that have been joined to a fluorescent dye. Finally, he washes off any unbound antibodies and looks at the slide under a microscope. Explain how the test works to confirm that the bacteria are present. [2 marks]



Plant Diseases and Defences

It's not just us humans that can get diseases - plants get their fair share too.

Plants Need Mineral Ions

Plants need mineral ions from the soil. If there aren't enough, plants suffer deficiency symptoms.

- 1) Nitrates are needed to make proteins and therefore for growth. A lack of nitrates causes stunted growth.
- 2) Magnesium ions are needed for making chlorophyll, which is needed for photosynthesis. Plants without enough magnesium suffer from chlorosis and have yellow leaves.

Plants Can Get Diseases

- 1) Plants can be infected by viral, bacterial and fungal pathogens see page 47 for some examples. They can also be infested and damaged by insects. For example, aphids are an insect that can cause huge damage to plants.
- 2) It's usually pretty clear that a plant has a disease. The common signs are:
 - 1) Stunted growth

- 2) Spots on the leaves
- Patches of decay (rot)

- 4) Abnormal growths, e.g. lumps 5) Malformed stems or leaves 6) Discolouration
- 3) Infestations of pests are easy to spot too you should be able to see them on the plants.
- 4) Different plant diseases have different signs. They can be identified by:
 - 1) Looking up the signs in a gardening manual or on a gardening website.
 - 2) Taking the infected plant to a laboratory, where scientists can identify the pathogen.
 - 3) Using testing kits that identify the pathogen using monoclonal antibodies (see page 53).

Plants Have Physical, Chemical and Mechanical Defences...

PHYSICAL DEFENCES

- 1) Most plant leaves and stems have a waxy cuticle, which provides a barrier to stop pathogens entering.
- 2) Plant cells themselves are surrounded by cell walls made from cellulose. These form a physical barrier against pathogens that make it past the waxy cuticle.
- 3) Plants have layers of dead cells around their stems, for example, the outer part of the bark on trees. These act as a barrier to stop pathogens entering.

CHEMICAL DEFENCES

- 1) Some can produce antibacterial chemicals which kill bacteria e.g. the mint plant and witch hazel.
- 2) Other plants produce poisons which can deter herbivores (organisms that eat plants) e.g. tobacco plants, foxgloves and deadly nightshade.

MECHANICAL DEFENCES

- 1) Some plants have adapted to have thorns and hairs. These stop animals from touching and eating them.
- 2) Other plants have leaves that droop or curl when something touches them. This means that they can prevent themselves from being eaten by knocking insects off themselves and moving away from things.
- 3) Some plants can cleverly mimic other organisms. E.g. the passion flower has bright yellow spots on its leaves which look like butterflu eggs. This stops other butterflies laying their eggs there. Several species of plant in the 'ice plant family' in southern Africa look like stones and pebbles. This tricks other organisms into not eating them.

Symptoms of revision deficiency include nagging parents...

You've made it to the end of the Topic. But don't run off just yet — have a go at this question first.

01 How could you tell that a plant has a nitrate deficiency?

Revision Questions for Topic 3

Well, that wraps up $\underline{\text{Topic 3}}$ — time to put your knowledge to the test.

- Try these questions and tick off each one when you get it right.
- When you're completely happy with a sub-topic, tick it off.

For even more practice, try the Retrieval Quiz for Topic 3 — just Scan this QR code!



Types of Disease (p.46-48)		
1)	How can bacteria make us feel ill?	\checkmark
2)	Give one way that fungi can cause disease.	
3)	How does tobacco mosaic virus affect a plant's growth?	
4)	How are mosquitoes involved in the spread of malaria?	
5)	What are the symptoms of gonorrhoea?	
6)	How can destroying vectors help to prevent the spread of disease?	\checkmark
Fighting Disease (p.49-52)		
7)	What does the stomach produce that can kill pathogens?	\checkmark
8)	Give three ways that the white blood cells can defend against pathogens.	
9)	Give one pro and one con of vaccination.	
10)	Why is it difficult to develop drugs that kill viruses without also damaging body tissues?	
11)	Which plant does the painkiller aspirin originate from?	
12)	What two things are drugs tested on in preclinical testing?	\checkmark
13)	What is meant by a drug's toxicity?	\checkmark
Antibodies (p.53-54)		
14)	What is a monoclonal antibody?	\checkmark
15)	Where are the B-lymphocytes taken from when making monoclonal antibodies?	\checkmark
16)	Why might a fluorescent dye be added to a monoclonal antibody?	\checkmark
17)	Why aren't monoclonal antibodies used as drugs as much as scientists hoped they would be?	\checkmark
Plant Diseases and Defences (p.55)		
18)	What do plants use magnesium ions for?	\checkmark
19)	Give one way of identifying a plant disease.	
20	Give one mechanical defence that a plant may have to defend itself.	V

Photosynthesis and Limiting Factors

First, photosunthesis equations. Then there are some more bits 'n' bobs you should know...

Photosynthesis Produces Glucose Using Light

- 1) Photosynthesis uses energy to change carbon dioxide and water into glucose and oxygen.
- 2) It takes place in chloroplasts in green plant cells they contain pigments like chlorophyll that absorb light.
- 3) Energy is transferred to the chloroplasts from the environment by light.
- 4) Photosynthesis is endothermic this means energy is transferred from the environment in the process.
- 5) The word equation for photosynthesis is:

6) Here's the symbol equation too:





Plants Use Glucose in Five Main Ways...

- For respiration This transfers energy from glucose (see p.61) which enables the plants to convert the rest of the glucose into various other useful substances.
- 2) Making cellulose Glucose is converted into cellulose for making strong plant cell walls (see p.11).
- Making amino acids <u>Glucose</u> is combined with <u>nitrate ions</u> (absorbed from the soil) to make <u>amino acids</u>, which are then made into <u>proteins</u>.
- 4) Stored as oils or fats Glucose is turned into lipids (fats and oils) for storing in seeds.
- 5) Stored as starch Glucose is turned into starch and stored in roots, stems and leaves, ready for use when photosynthesis isn't happening, like in the winter. Starch is insoluble, which makes it much better for storing than glucose a cell with lots of glucose in would draw in loads of water and swell up.

Limiting Factors Affect the Rate of Photosynthesis

- 1) The rate of photosynthesis is affected by intensity of light, concentration of CO, and temperature.
- Any of these three factors can become the <u>limiting factor</u> this just means that it's stopping photosynthesis from happening any <u>faster</u>.
- 3) These factors have a combined effect on the rate of photosynthesis, but which factor is limiting at a particular time depends on the environmental conditions:
 - · at night it's pretty obvious that light is the limiting factor,
 - in winter it's often the temperature.
 - if it's warm enough and bright enough, the amount of CO is usually limiting.
- 4) Chlorophull can also be a limiting factor of photosynthesis.

The <u>amount of chlorophyll</u> in a plant can be affected by <u>disease</u> (e.g. infection with the tobacco mosaic virus) or <u>environmental stress</u>, such as a <u>lack of nutrients</u>. These factors can cause <u>chloroplasts</u> to become <u>damaged</u> or to <u>not</u> make <u>enough chlorophyll</u>. This means the rate of photosynthesis is <u>reduced</u> because they <u>can't absorb</u> as much <u>light</u>.

Now you'll have something to bore the great-grandkids with...

You'll be able to tell them how, in your day, all you needed was a bit of carbon dioxide and some water and you could make your own entertainment. But at the moment you need to learn this page...

Q1 Name the products of photosynthesis.

[2 marks]

O2 Apart from temperature, name three other limiting factors of photosynthesis.


[3 marks]

The Rate of Photosynthesis

Now that you know light, CO₂ and temperature all <u>affect</u> the <u>rate of photosynthesis</u>, you also need to know <u>how</u> they affect the rate, so you can take a gander at a load of lovely pictures... well, graphs. I've also thrown an experiment and an equation in for good measure. I can tell these pages are going to be your favourites...

Three Important Graphs for Rate of Photosynthesis


1) Not Enough Light Slows Down the Rate of Photosynthesis




- 1) Light provides the energy needed for photosynthesis.
- As the light level is raised, the rate of photosynthesis increases steadily — but only up to a certain point.
- Beyond that, it won't make any difference as light intensity increases, the rate will no longer increase.
 This is because it'll be either the temperature or the CO₂ level which is now the limiting factor, not light.
- In the lab you can change the light intensity by moving a lamp closer to or further away from your plant (see the next page for this experiment).
- 5) But if you just plot the rate of photosynthesis against "distance of lamp from the plant", you get a weird-shaped graph. To get a graph like the one above you either need to measure the light intensity at the plant using a light meter or do a bit of nifty maths with your results.

2) Too Little Carbon Dioxide Also Slows it Down

- CO₂ is one of the <u>raw materials</u> needed for photosynthesis.
- 2) As with light intensity, the amount of CO₂ will only increase the rate of photosynthesis up to a point. After this the graph <u>flattens out</u>—as the amount of CO₂ increases, the rate no longer increases. This shows that CO₂ is no longer the <u>limiting factor</u>.
- As long as <u>light</u> and <u>CO</u>₂ are in plentiful supply then the factor limiting photosynthesis must be <u>temperature</u>.



3) The Temperature has to be Just Right



- Usually, if the temperature is the <u>limiting factor</u> it's because it's too <u>low</u> — the <u>enzymes</u> needed for photosynthesis work more <u>slowly</u> at low temperatures.
- But if the plant gets too hot, the enzymes it needs for photosynthesis and its other reactions will be damaged.
- This happens at about 45 °C (which is pretty hot for outdoors, although greenhouses can get that hot if you're not careful).


The Rate of Photosynthesis

One Graph May Show the Effect of Many Limiting Factors


You could get a graph that shows more than one limiting factor on the rate of photosynthesis, for example:

 The graph on the right shows how the rate of photosynthesis is affected by <u>light intensity</u> and <u>temperature</u>.

- At the start, both of the lines show that as the light intensity <u>increases</u>, the rate of photosynthesis <u>increases steadily</u>.
- 3) But the lines level off when light is no longer the limiting factor. The line at 25 °C levels off at a higher point than the one at 15 °C, showing that temperature must have been a limiting factor at 15 °C.



- 1) The graph on the right shows how the rate of photosynthesis is affected by light intensity and CO, concentration.
- 2) Again, both the lines <u>level off</u> when <u>light</u> is <u>no longer</u> the limiting factor.
- 3) The line at the higher CO₂ concentration of 0.4% levels off at a higher point than the one at 0.04%. This means CO₂ concentration must have been a <u>limiting factor</u> at 0.04% CO₂. The limiting factor here isn't temperature because it's the <u>same</u> for both lines (25 °C).



Oxygen Production Shows the Rate of Photosynthesis

PRACTICAL

<u>Canadian pondweed</u> can be used to measure the effect of <u>light intensity</u> on the <u>rate of photosynthesis</u>. The rate at which the pondweed produces <u>oxugen</u> corresponds to the rate at which it's photosynthesising — the <u>faster</u> the rate of oxygen production, the <u>faster</u> the rate of photosynthesis.

Here's how the experiment works:

- A source of <u>white light</u> is placed at a <u>specific distance</u> from the pondweed.
- The pondweed is left to photosynthesise for a <u>set amount of time</u>.
 As it photosynthesises, the oxygen released will collect in the capillary tube.
- 3) At the end of the experiment, the <u>suringe</u> is used to draw the gas bubble in the tube up allongside a ruler and the <u>length</u> of the <u>gas bubble</u> is <u>measured</u>. This is <u>proportional</u> to the <u>volume of O</u>, produced.



- 4) For this experiment, any <u>variables</u> that could affect the results should be <u>controlled</u>, e.g. the <u>temperature</u> and <u>time</u> the pondweed is left to photosynthesise.
- 5) The experiment is <u>repeated</u> twice with the <u>light source</u> at the <u>same</u> distance and the <u>mean</u> volume of O₂ produced is calculated.
- 6) Then the whole experiment is repeated with the light source at different distances from the pondweed.

The apparatus above can be altered to measure the effect of temperature or \underline{CO}_2 on photosynthesis. E.g. the test tube of pondweed can be put into a <u>water bath</u> at a <u>set temperature</u>, or a measured amount of <u>sodium hydrogencarbonate</u> can be dissolved in the water (which gives off \underline{CO}_2). The experiment can then be repeated with different temperatures of water / concentrations of sodium hydrogencarbonate.


The Rate of Photosynthesis

The Inverse Square Law Links Light Intensity and Distance

- In the experiment on the previous page, when the lamp is moved away from the pondweed, the amount of light that reaches the pondweed decreases.
- 2) You can say that as the distance increases, the light intensity decreases.
- In other words, distance and light intensity are inversely proportional to each other.

 3) However, it's not quite as simple as that. It turns out that light intensity decreases in proportion to the square of the distance. This is called the inverse square law and is written out like this:

 | This is the 'proportional to' symbol. This is the 'proportional to' symbol. The distance inverse square law and is written out like this:
- 4) The inverse square law means that if you halve the distance, the light intensity will be four times greater and if you third the distance, the light intensity will be nine times greater. Likewise, if you double the distance, the light intensity will be four times smaller and if you treble the distance, the light intensity will be nine times smaller.
- 5) You can use 1/d2 as a measure of light intensity.



You can Artificially Create the Ideal Conditions for Farming


- 1) The most common way to artificially create the ideal environment for plants is to grow them in a greenhouse.
- 2) Greenhouses help to <u>trap</u> the Sun's <u>heat</u>, and make sure that the <u>temperature</u> doesn't become <u>limiting</u>. In winter a farmer or gardener might use a <u>heater</u> as well to keep the temperature at the ideal level. In summer it could get <u>too hot</u>, so they might use <u>shades</u> and <u>ventilation</u> to cool things down.



- Light is always needed for photosynthesis, so commercial farmers often supply artificial light after the Sun goes down to give their plants more quality photosynthesis time.
- 4) Farmers and gardeners can also increase the level of <u>carbon dioxide</u> in the greenhouse. E.g. by using a <u>paraffin heater</u> to heat the greenhouse. As the paraffin burns, it makes carbon dioxide as a <u>bu-product</u>.
- 5) Keeping plants enclosed in a greenhouse also makes it easier to keep them free from <u>pests</u> and <u>diseases</u>. The farmer can add <u>fertilisers</u> to the soil as well, to provide all the <u>minerals</u> needed for healthy growth.
- 6) Sorting all this out <u>costs money</u> but if the farmer can keep the conditions just right for photosynthesis, the plants will grow much <u>faster</u> and a <u>decent crop</u> can be harvested much more <u>often</u>, which can then be <u>sold</u>. It's important that a farmer supplies just the <u>right amount</u> of heat, light, etc. enough to make the plants grow well, but <u>not</u> more than the plants need, as this would just be <u>wasting money</u>.

With enough light, photographers can also photosynthesise...

Now don't let the inverse square law put you off learning everything on these past three pages.





Q2 A plant is moved from 15 cm away from its light source to 5 cm away from its light source.
Using the inverse square law, show that the light intensity becomes nine times greater. [3 marks]

Respiration and Metabolism

You need energy to keep your body going. Energy comes from food, and it's transferred by respiration.

Respiration is NOT "Breathing In and Out"

Respiration involves many reactions. These are really important reactions, as respiration transfers the energy that the cell needs to do just about everything — this energy is used for all living processes.

- 1) Respiration is not breathing in and breathing out, as you might think.
- Respiration is the process of transferring energy from the breakdown of glucose (sugar)
 — and it goes on in every cell in your body continuously.
- It happens in plants too. All living things respire. It's how they transfer energy from their food
 to their cells.

<u>RESPIRATION</u> is the process of <u>IRANSFERRING ENERGY FROM GLUCOSE</u>, which goes on <u>IN EVERY CELL</u>.

4) Respiration is exothermic — it transfers energy to the environment.

Respiration Transfers Energy for All Kinds of Things

Here are three examples of how organisms use the energy transferred by respiration:

- 1) To build up larger molecules from smaller ones (like proteins from amino acids see below).
- 2) In animals it's used to allow the muscles to contract (so they can move about).
- In <u>mammals</u> and <u>birds</u> the energy is used to keep their <u>body temperature</u> steady in colder surroundings. (Unlike other animals, mammals and birds keep their bodies constantly warm.)

Metabolism is ALL the Chemical Reactions in an Organism

- 1) In a <u>cell</u> there are <u>lots</u> of <u>chemical reactions</u> happening <u>all the time</u>, which are controlled by <u>enzymes</u>.
- 2) Many of these reactions are linked together to form bigger reactions:



- 3) In some of these reactions, larger molecules are made from smaller ones. For example:
 - Lots of small <u>glucose</u> molecules are <u>joined together</u> in reactions to form <u>starch</u>
 (a storage molecule in plant cells), <u>glucogen</u> (a storage molecule in animal cells)
 and <u>cellulose</u> (a component of plant cell walls).
 - · Lipid molecules are each made from one molecule of glucerol and three fatty acids.
 - . Glucose is combined with nitrate ions to make amino acids, which are then made into proteins.
- 4) In other reactions, larger molecules are broken down into smaller ones. For example:
 - Glucose is broken down in <u>respiration</u>. Respiration transfers energy to power all the reactions in the body that make molecules.
 - Excess protein is broken down in a reaction to produce urea. Urea is then excreted in urine.
- 5) The sum (total) of all of the reactions that happen in a cell or the body is called its metabolism.

Respiration transfers energy — but this page has worn me out...

Isn't it strange to think that each individual living cell in your body is respiring every second of every day, transferring energy from the food you eat. This energy is used to make molecules that our cells need.

Q1 Give two examples of how animals use the energy transferred by respiration. [2 marks]

Q2 What is metabolism? [1 mark]

Aerobic and Anaerobic Respiration

There are two types of respiration, don't cha know...

Aerobic Respiration Needs Plenty of Oxygen

- Aerobic respiration is respiration using oxygen.
 It's the most efficient way to transfer energy from glucose.
- 2) Aerobic respiration goes on all the time in plants and animals.
- 3) Most of the reactions in aerobic respiration happen inside mitochondria (see page 11).
- 4) Here are the word and symbol equations for aerobic respiration:

glucose + oxygen
$$\longrightarrow$$
 carbon dioxide + water
$$C_6H_{12}O_6 + 6O_2 \longrightarrow 6CO_2 + 6H_2O$$

Anaerobic Respiration is Used if There's Not Enough Oxygen

When you do vigorous exercise and your body can't supply enough <u>oxygen</u> to your muscles, they start doing <u>anaerobic respiration</u> as well as aerobic respiration.

- 1) "Anaerobic" just means "without oxygen". It's the incomplete breakdown of glucose, making lactic acid.
- 2) Here's the word equation for anaerobic respiration in muscle cells:

- Anaerobic respiration does not transfer nearly as much energy as aerobic respiration.
 This is because glucose isn't fully oxidised (because it doesn't combine with oxygen).
- 4) So, anaerobic respiration is only useful in emergencies, e.g. during exercise when it allows you to keep on using your muscles for a while longer.

Anaerobic Respiration in Plants and Yeast is Slightly Different

- Plants and yeast cells can respire without oxygen too, but they produce ethanol (alcohol) and carbon dioxide instead of lactic acid.
- 2) Here is the word equation for anaerobic respiration in plants and yeast cells:





- 3) Anaerobic respiration in yeast cells is called fermentation.
- In the food and drinks industry, fermentation by yeast is of great value because it's used to make bread and alcoholic drinks, e.g. beer and wine.
- 5) In bread-making, it's the carbon dioxide from fermentation that makes bread rise.
- 6) In beer and wine-making, it's the fermentation process that produces alcohol.

I'd like a ham and fermentation sandwich please... yum

Fermentation is a really important process because of its use in making alcoholic drinks and bread. We drink and eat so much of these that making them is big bucks. And it's all down to tiny yeast cells.

Q1 What are the reactants of aerobic respiration?

[2 marks]

O2 What is the process of anaerobic respiration in yeast called?



Exercise

When you exercise, your body responds in different ways to get enough energy to your cells.

When You Exercise You Respire More

- Muscles need energy from respiration to contract. When you exercise, some of your muscles contract
 more frequently than normal so you need more energy. This energy comes from increased respiration.
- The increase in respiration in your cells means you need to get more oxygen into them.
- 3) Your breathing rate and breath volume increase to get more oxygen into the blood, and your heart rate increases to get this oxygenated blood around the body faster. This removes CO₂ more quickly at the same time.
- When you do really vigorous exercise (like sprinting) your body can't supply oxygen to your muscles quickly enough, so they start respiring anaerobically (see the previous page).
- This is NOT the best way to transfer energy from glucose because lactic acid builds up in the muscles, which gets painful.
- Long periods of exercise also cause muscle fatigue the muscles get <u>tired</u> and then <u>stop contracting efficiently</u>.



Anaerobic Respiration Leads to an Oxygen Debt

- 1) After resorting to anaerobic respiration, when you stop exercising you'll have an "oxusen debt".
- An oxygen debt is the amount of extra oxygen your body needs to react with the build up of lactic acid and remove it from the cells. Oxygen reacts with the lactic acid to form harmless CO_o and water.
- In other words you have to "repay" the oxygen that you didn't get to your muscles in time, because your lungs, heart and blood couldn't keep up with the demand earlier on.
- This means you have to keep breathing hard for a while <u>after you stop</u>, to get <u>more oxygen</u> into your blood, which is transported to the muscle cells.
- 5) The <u>pulse</u> and <u>breathing rate</u> stay high whilst there are <u>high levels</u> of <u>lactic acid</u> and <u>CO.</u>.
- 6) Your body also has another way of coping with the high level of lactic acid the <u>blood</u> that enters your muscles <u>transports</u> the <u>lactic acid</u> to the <u>liver</u>. In the liver, the lactic acid is <u>converted</u> back to <u>glucose</u>.


You Can Investigate The Effect of Exercise on The Body

1) You can measure breathing rate by counting breaths, and heart rate by taking the pulse.

2) E.g. you could take your pulse after:

- sitting down for 5 minutes,
- then after 5 minutes of gentle walking.
- then again after 5 minutes of slow jogging.
- then again after <u>running</u> for 5 minutes.

and plot your results in a bar chart.



You put two fingers on the interest of your wrist or your energy of the count the number of pulses in 1 minute.

- 3) Your pulse rate will increase the more intense the exercise is, as your body needs to get more oxygen to the muscles and take more carbon dioxide away from the muscles.
- To reduce the effect of any random errors on your results, do it as a group and plot the average pulse rate for each exercise.

There's more about = random error on page 5. =

Oxygen debt — cheap to pay back...

At the end of a sprinting race you often see athletes breathing hard — now you know this is to get rid of the lactic acid that's built up in the muscles. But remember, the liver plays a role in breaking it down too.

Q1 Look at the graph above. Predict which type of exercise would lead to the highest concentration of lactic acid in the blood after 10 minutes. Explain your answer.



Revision Questions for Topic 4

Well, it's all over for Topic 4 folks. So here are some questions on it...

- Try these questions and tick off each one when you get it right.
- When you're completely happy with a sub-topic, tick it off.

23) What happens to muscles when they become fatigued? 24) In what organ is lactic acid converted back to glucose?

For even more practice, try the Retrieval Quiz for Topic 4 - just scan this QR code!



Photosynthesis (p.57-60) Where in a plant cell does photosynthesis take place? 2) How is energy transferred to a plant from its environment for photosynthesis? 3) What is an endothermic reaction? 4) What is the word equation for photosunthesis? 5) What tupe of molecule is formed from combining glucose molecules with nitrate ions? 6) Why do plants store glucose as starch? 7) What is meant by a 'limiting factor' of photosynthesis? 8) What effect would a low carbon dioxide concentration have on the rate of photosynthesis? 9) What effect would a temperature above 45 °C have on the rate of photosynthesis? 10) Describe a method that could be used to measure the effect of light intensity on the rate of photosynthesis. 11) In the inverse square law, how are light intensity and distance linked? 12) Give one way that the temperature can be decreased in a greenhouse. 13) Give one way that the level of carbon dioxide can be increased in a greenhouse. Respiration and Metabolism (p.61-63) 14) What is respiration? 15) What is an exothermic reaction? 16) What process transfers energy to make new molecules in cells? 17) Give three examples of metabolic reactions. 18) Name the products of aerobic respiration. 19) What is produced by anaerobic respiration in muscle cells? 20) What is the word equation for anaerobic respiration in yeast cells? 21) Name two products of the food and drink industry that fermentation is needed for. 22) Give three things that increase to supply the muscles with more oxygenated blood during exercise.

I'm not really a doctor — this clipboard isn't holding anything. But take it from

ne homenotorio io one important tonio

Homeostasis


<u>Homeostasis</u> — a word that strikes fear into the heart of many a GCSE student. But it's really not that bad at all. This page is a brief introduction to the topic, so you need to nail all of this before you can move on.

Homeostasis — Maintaining a Stable Internal Environment

- The conditions inside your body need to be kept steady, even when the external environment changes. This is really important because your cells need the right conditions in order to function properly, including the right conditions for enzume action (see p.28).
- Homeostasis is all about the <u>regulation</u> of the conditions inside your body (and cells) to <u>maintain a stable internal environment</u>, in response to <u>changes</u> in both internal and external conditions.
- 3) You have loads of <u>automatic control systems</u> in your body that regulate your internal environment — these include both <u>nervous</u> and <u>hormonal</u> communication systems. For example, there are control systems that maintain your <u>body temperature</u> (see page 72), your <u>blood plucose</u> level (see page 74) and your <u>water content</u> (see page 75).
- 4) All your automatic control systems are made up of <u>three main components</u> which work together to maintain a steady condition — cells called <u>receptors</u>, <u>coordination centres</u> (including the brain, spinal cord and pancreas) and effectors.

Negative Feedback Counteracts Changes

Your automatic control systems keep your internal environment stable using a mechanism called negative-feedback. When the level of something (e.g. water or temperature) gets too low, your body uses negative feedback to bring it back to normal.



This process happens without you thinking about it — it's all automatic.

If you do enough revision, you can avoid negative feedback...

making the level change too much (away from the ideal). Luckily the receptor detects if the level becomes too different and negative feedback starts again.

Negative feedback is a fancy-sounding name for a not-very-complicated idea. It's common sense really. For example, if you looked sad, I'd try and cheer you up. And if you looked really happy, I'd probably start to annoy you by flicking the backs of your ears. It stops things getting out of balance, I think.

Q1 Why do the internal conditions of your body need to be regulated?

[1 mark]


O2 Name the component of a control system that detects stimuli.

The Nervous System

Organisms need to respond to stimuli (changes in the environment) in order to <u>survive</u>. A <u>single-celled</u> organism can just respond to its environment, but the cells of <u>multicellular</u> organisms need to <u>communicate</u> with each other first. So as multicellular organisms evolved, they developed nervous and hormonal communication systems.

The Nervous System Detects and Reacts to Stimuli

The nervous system means that humans can react to their surroundings and coordinate their behaviour.



Receptors and Effectors can form part of Complex Organs


- Receptors are the cells that detect stimuli.
- There are many <u>different types</u>
 of receptors, such as <u>taste</u>
 receptors on the tongue and
 sound receptors in the ears.
- Receptors can form part
 of larger, complex organs,
 e.g. the retina of the eye is
 covered in light receptor cells.
- Effectors respond to nervous impulses and bring about a change.
- 5) Muscles and glands are known as <u>effectors</u>—they respond in different ways. <u>Muscles contract</u> in response to a nervous impulse, whereas <u>glands secrete hormones</u>.

The Central Nervous System (CNS) Coordinates the Response

The CNS is a <u>coordination centre</u> — it receives information from the <u>receptors</u> and then <u>coordinates a response</u> (decides what to do about it). The response is carried out by <u>effectors</u>.

For example, a small bird is eating some seed...

- 1) ...when, out of the corner of its eye, it spots a cat skulking towards it (this is the stimulus).
- The receptors in the bird's eye are stimulated. Sensory neurones carry the information from the receptors to the CNS.
- 3) The CNS decides what to do about it.
- The CNS sends information to the muscles in the bird's wings (the effectors) along motor neurones. The muscles contract and the bird flies away to safety.



Don't let the thought of exams play on your nerves...

Don't forget that it's only large animals like mammals and birds that have complex nervous systems. Simple animals like jellyfish don't — everything they do is a reflex response (see next page).

Name two types of effector.


[2 marks]

Synapses and Reflexes

<u>Neurones</u> transmit information <u>very quickly</u> to and from the brain, and your brain <u>quickly decides</u> how to respond to a stimulus. But reflexes are even quicker...

Synapses Connect Neurones


- 1) The connection between two neurones is called a sunapse.
- 2) The nerve signal is transferred by <u>chemicals</u> which <u>diffuse</u> (move) across the gap.
- 3) These chemicals then set off a new electrical signal in the next neurone.



Reflexes Help Prevent Injury

- Reflexes are rapid, automatic responses to certain stimuli that don't involve the conscious part of the brain — they can reduce the chances of being injured.
- For example, if someone shines a <u>bright light</u> in your eyes, your <u>pupils</u> automatically get smaller so that less light gets into the eye — this stops it getting <u>damaged</u>.
- Or if you get a shock, your body releases the <u>hormone</u> adrenaline automatically
 — it doesn't wait for you to <u>decide</u> that you're shocked.
- 4) The passage of information in a reflex (from receptor to effector) is called a reflex arc.

The Reflex Arc Goes Through the Central Nervous System



- The neurones in reflex arcs go through the <u>spinal</u> cord or through an unconscious part of the brain.
- 2) When a <u>stimulus</u> (e.g. a painful bee sting) is detected by receptors, <u>impulses</u> are sent along a sensoru neurone to a relau neurone in the CNS.
- B) When the impulses reach a <u>synapse</u> between the sensory neurone and the relay neurone, they trigger chemicals to be released (see above).
 - These chemicals cause impulses to be sent along the relay neurone.
- Relay neurones connect sensory
- 4) When the impulses reach a <u>sunapse</u> between the relay neurone and a motor neurone, the same thing happens. Chemicals are released and cause impulses to be sent along the <u>motor neurone</u>.
- The impulses then travel along the motor neurone to the effector (which is usually a <u>muscle</u>, like in this example).
- 6) The muscle then contracts and moves your hand away from the bee.
- 7) Because you don't have to think about the response (which takes time) it's quicker than normal responses.

Don't get all twitchy — just learn it...

Reflexes bypass your conscious brain completely when a quick response is essential — your body just gets on with things. If you had to stop and think first, you'd end up a lot more sore (or worse).



Q1 What is a reflex action?

- [1 mark]
- O2 A chef touches a hot pan. A reflex reaction causes him to immediately move his hand away.
 - a) State the effector in this reflex reaction.
 - b) Describe the pathway of the reflex from stimulus to effector.

[1 mark] [4 marks]



Investigating Reaction Time

On your marks... get set... read this page.

Reaction Time is How Quickly You Respond

Reaction time is the time it takes to <u>respond to a stimulus</u> — it's often <u>less</u> than a <u>second</u>. It can be <u>affected</u> by factors such as <u>age, gender</u> or <u>drugs</u>.

You Can Measure Reaction Time

Caffeine is a drug that can speed up a person's reaction time.

The effect of caffeine on reaction time can be measured like this...

- The person being tested should sit with their arm resting on the edge of a table (this should stop them moving their arm up or down during the test).
- Hold a <u>ruler</u> vertically between their thumb and forefinger. Make sure that the <u>zero end</u> of the ruler is <u>level</u> with their thumb and finger. Then <u>let go</u> without giving any warning.
- The person being tested should try to catch the ruler as quickly as they can — as soon as they see it fall.
- 4) Reaction time is measured by the <u>number</u> on the ruler where it's caught. The number should be read from the <u>top</u> of the <u>thumb</u>. The further down the ruler it's caught (i.e. the higher the number), the slower their reaction time.
- 5) Repeat the test several times then calculate the mean distance that the ruler fell.
- The person being tested should then have a <u>caffeinated drink</u> (e.g. 300 ml of cola). After <u>ten minutes</u>, repeat steps 1 to 5.
- 7) You need to control any variables to make sure that this is a fair test.

 For example, you should use the same person to catch the ruler each time, and that person should always use the same hand to catch the ruler. Also, the ruler should always be dropped from the same height, and you should make sure that the person being tested has not had any caffeine (or anything else that may affect their reaction time) before the start of the experiment.
- 8) Too much caffeine can cause <u>unpleasant side-effects</u>, so the person being tested should avoid drinking <u>any more caffeine</u> for the rest of the day after the experiment is completed.

Reaction Time Can Be Measured Using a Computer



- Simple computer tests can also be used to measure reaction time.
 For example, the person being tested has to click the mouse (or press a key) as soon as they see a stimulus on the screen, e.g. a box change colour.
- Computers can give a <u>more precise</u> reaction time because they remove the possibility of <u>human error</u> from the measurement.
- As the computer can record the reaction time in <u>milliseconds</u>, it can also give a more <u>accurate</u> measurement.
- 4) Using a computer can also remove the possibility that the person can predict when to respond — using the ruler test, the catcher may learn to anticipate the drop by reading the tester's body language.

Ready... Steady...

... Ah, too slow.

- Q1 Some students investigated the effect of an energy drink on reaction time. They measured their reaction times using a computer test. They had to click the mouse when the screen changed from red to green.
 Each student repeated the test five times before having an energy drink, and five times afterwards.
 a) The results for one of the students before having the energy drink were as follows:
 - 242 ms, 256 ms, 253 ms, 249 ms, 235 ms. Calculate the mean reaction time. [2 marks]
 - b) Suggest two variables that the students needed to control during their investigation. [2 marks]



30 cm

0 cm

14 cm

With a little bit of maths, it's possible


to work out the reaction time in

The Brain

Scientists know a bit about the <u>brain</u> but <u>not as much</u> as they'd like. Their knowledge is improving with the invention of new gadgetry that helps them study the brain. Read on, it's pretty amazing stuff.

The Brain is Responsible for Complex Behaviours

- 1) Along with the spinal cord, the brain is part of the central nervous system.
- 2) It's made up of billions of interconnected neurones (neurones that are connected together).
- The brain is in charge of all of our <u>complex behaviours</u>. It controls and coordinates everything you do — running, breathing, sleeping, remembering your gym kit...
- 4) We know that different regions of the brain carry out different functions:



Scientists Use a Range of Methods to Study the Brain

Scientists use a few different methods to study the brain and map out which bits do what:

Studying patients with brain damage If a <u>small</u> part of the brain has been <u>damaged</u>, the <u>effect</u> this has on the patient can tell you a lot about what the damaged part of the brain does. E.g. if an area at the back of the brain was damaged by a stroke and the patient went blind, you know that that area has something to do with vision.

Scientists that study the brain are called neuroscientists.

Electrically stimulating the brain The brain can be <u>stimulated electrically</u> by pushing a tiny <u>electrode</u> into the tissue and giving it a small zap of electricity. By observing what stimulating <u>different parts</u> of the brain does, it's possible to get an idea of what those parts do. E.g. when a certain part of the brain (known as the motor area) is stimulated, it causes muscle contraction and movement.

MRI Scans A <u>magnetic resonance imaging (MRI) scanner</u> is a big fancy tube-like machine that can produce a very <u>detailed picture</u> of the brain's structures. Scientists use it to find out what areas of the brain are <u>active</u> when people are doing things like listening to music or truing to recall a memory.



Messing With the Brain Can Have Consequences

- Knowledge of how the brain works has led to the development of <u>treatments</u> for <u>disorders</u> of the nervous system. For example, <u>electrical stimulation</u> of the brain can help reduce <u>muscle tremors</u> caused by nervous system disorders such as Parkinson's disease.
- 2) However, the brain is incredibly complex and delicate the investigation of brain function and any treatment of brain damage or disease is difficult. It also carries <u>risks</u>, such as <u>physical damage</u> to the brain or increased problems with brain function (e.g. difficulties with speech).

A whole page dedicated to that squidgy thing in your head...

...lucky you. But actually it's really fascinating stuff. It'll make it that bit easier to learn...

Q1 Name the region of the brain that controls coordinated movement.


[1 mark]

The Eye

The eye is a sense organ. There are several parts you need to learn about, so get focused.

Learn the Eye with All Its Labels


- 1) The SCLERA is the tough, supporting wall of the eye.
- 2) The CORNEA is the transparent outer layer found at the front of the eye. It refracts (bends) light into the eye.
- 3) The IRIS contains muscles that allow it to control the diameter of the PUPIL (the hole in the middle) and therefore how much light enters the eye.
- 4) The LENS focuses the light onto the RETINA (which contains receptor cells sensitive to light intensity and colour).
- 5) The shape of the lens is controlled by the CILIARY MUSCLES and SUSPENSORY LIGAMENTS.
- 6) The OPTIC NERVE carries impulses from the receptors on the retina to the brain.



The Iris Reflex — Adjusting for Bright Light

Very bright light can damage the retina — so you have a reflex to protect it.

- 1) When light receptors in the eue detect very bright light, a reflex is triggered that makes the pupil smaller. The circular muscles in the iris contract and the radial muscles relax. This reduces the amount of light that can enter the eye.
- 2) The opposite process happens in dim light. This time, the radial muscles contract and the circular muscles relax, which makes the pupil wider.





Focusing on Near and Distant Objects — Another Reflex


The eye focuses light on the retina by changing the shape of the lens — this is known as accommodation.

To look at near objects:



- 1) The ciliary muscles contract, which slackens the suspensory ligaments.
- 2) The lens becomes fat (more curved).
- 3) This increases the amount by which it refracts light.

To look at distant objects:



- The ciliary muscles relax, which allows the suspensory ligaments to pull tight.
- 2) This makes the lens go thin (less curved).
- 3) So it refracts light by a smaller amount.

As you get older, your eye's lens loses <u>flexibility</u>, so it can't easily spring back to a round shape. This means light can't be focused well for near viewing, so older people often have to use reading glasses.

If the lens cannot refract the light by the right amount (so that it focuses on the retina), the person will be short- or long-sighted — see the next page for more.

Eye eye, Captain...

It doesn't matter how good you are at blagging in the exam — you need to learn those diagrams of the eye.

01 What name is given to the transparent layer at the front of the eye? [1 mark]

O2 Explain how the eye focuses on an object that's close to it.

[4 marks]

image

image

Corrected

Corrected

Correcting Vision Defects

As you can see on the previous page, the way the eye works is quite <u>complex</u>. It's not surprising really that sometimes it <u>doesn't work so well</u>. Thankfully, we have ways of <u>correcting</u> it...

Some People are Long- or Short-Sighted

Long-sighted people are unable to focus on near objects:

- This occurs when the <u>lens</u> is the wrong shape and doesn't <u>refract</u> (bend) the light enough or the <u>eyeball</u> is too <u>short</u>.
- The images of near objects are brought into focus <u>behind</u> the <u>retina</u>.
- 3) You can use glasses with a <u>convex lens</u> (a lens which <u>curves outwards</u>) to correct it. The lens <u>refracts</u> the light rays so they focus on the <u>retina</u>.

Uncorrected

Uncorrected

image

4) The medical term for long-sightedness is huperopia.

Short-sighted people are unable to focus on distant objects:

- This occurs when the <u>lens</u> is the wrong shape and refracts the light too much or the eyeball is too long.
- The images of distant objects are brought into focus in front of the retina.



4) The medical term for short-sightedness is muopia.

There are Several Treatments for Vision Defects

Wearing glasses isn't for everyone. You need to know about these alternatives:

<u>CONTACT LENSES</u>: <u>Contact lenses</u> are thin lenses that sit on the <u>surface</u> of the eye and are shaped to <u>compensate</u> for the fault in focusing. They're popular because they are <u>lightweight</u> and almost <u>invisible</u>. They're also more <u>convenient</u> than glasses for activities like <u>sports</u>. The two main types of contact lenses are <u>hard lenses</u> and <u>soft lenses</u>. Soft lenses are generally more comfortable, but carry a higher risk of eye infections than hard lenses.



LASER EYE SURGERY: Bad eyesight can sometimes be corrected with laser eye surgery. A laser can be used to vaporise tissue, changing the shape of the cornea (and so changing how strongly it refracts light into the eye). Slimming it down makes it less powerful and can improve short sight. Changing the shape so that it's more powerful will improve long sight. The surgeon can precisely control how much tissue the laser takes off, completely correcting the vision. However, like all surgical procedures, there is a risk of complications, such as infection or the eye reacting in a way that makes your vision worse than before.

REPLACEMENT LENS SURGERY: Sometimes long-sightedness may be more effectively treated by replacing the lens of the eye (rather than altering the shape of the cornea with laser eye surgery). In replacement lens surgery, the natural lens of the eye is removed and an artificial lens, made of clear plastic, is inserted in its place. As it involves work inside the eye, replacing a lens carries higher risks than laser eye surgery, including possible damage to the retina (which could lead to loss of sight).



I think I'm a little long-sighted...

You won't have to draw those diagrams for different lenses in the exam, but you may have to interpret them, so it's probably for the best that you make sure you totally understand what's going on in each one.

Q1 Explain the cause of short-sightedness.

O2 How can glasses help a long-sighted person focus on something nearby?

[2 marks] [1 mark]

Controlling Body Temperature

The body has to keep its insides at around 37 °C — the optimum temperature for enzymes in the body.

Body Temperature Must be Kept Constant

1) The body has to balance the amount of energy gained (e.g. through


sensitive to the temperature of the blood flowing through the brain.

- temperature inside your body, where your internal organs are. respiration) and lost to keep the core body temperature constant. 2) There is a thermoregulatory centre in the brain, which contains receptors that are
- 3) The thermoregulatory centre also receives impulses from temperature receptors in the skin, giving information about skin temperature.
 - 1) Temperature receptors detect that 1) Temperature receptors detect that core body temperature is too high. core body temperature is too low. 2) The thermoregulatory centre acts as 2) The thermoregulatory centre acts as a coordination centre - it receives a coordination centre - it receives information from the temperature receptors information from the temperature receptors and triggers the effectors automatically. and triggers the effectors automatically. 3) Effectors, e.g. sweat glands, 3) Effectors, e.g. muscles, produce a response (see below) produce a response (see below) and counteract the change. and counteract the change.


Some effectors work antagonistically, e.g. one effector heats and another cools — they'll work at the same time to achieve a very precise temperature. This mechanism allows a more sensitive response.

The Body has Some Nifty Tricks for Altering its Temperature

<u>Different responses</u> are produced by <u>effectors</u> to <u>counteract</u> an increase or decrease in <u>body temperature</u>.



- 1) Sweat is produced by sweat glands and evaporates from the skin. This transfers energy to the environment.
- 2) The blood vessels supplying the skin dilate so more blood flows close to the surface. of the skin. This is called vasodilation. This helps transfer energy from the skin to the environment.



Core body temperature is the

- 1) Hairs stand up to trap an insulating layer of air.
- 2) No sweat is produced.
- 3) Blood vessels supplying skin capillaries constrict to close off the skin's blood supply. This is called vasoconstriction.
- 4) When you're cold you shiver too (your muscles contract automatically). This needs respiration, which transfers some energy to warm the body.

Shiver me timbers — it's a wee bit nippy in here...

People who are exposed to extreme cold for a long time without protection can get frostbite — the blood supply to the fingers and toes is cut off to reduce the amount of energy lost (but this kills the cells, and they go black)... yuk.

Name the area of the brain that controls body temperature.

[1 mark]




The Endocrine System

The other way to send information around the body (apart from along nerves) is by using hormones.

Hormones Are Chemical Messengers Sent in the Blood

- Hormones are chemical molecules released directly into the blood. They are carried in the blood to
 other parts of the body, but only affect particular cells in particular organs (called <u>target organs</u>).
 Hormones control things in organs and cells that need <u>constant adjustment</u>.
- Hormones are produced in (and secreted by) various glands, called endocrine glands.
 These glands make up your endocrine system.
- 3) Hormones tend to have relatively long-lasting effects.



Hormones and Nerves Have Differences

NERVES: Very FAST action.

Act for a very SHORT TIME.

Act on a very PRECISE AREA.

HORMONES:

SLOWER action.
Act for a LONG TIME.
Act in a more GENERAL way.

So if you're not sure whether a response is nervous or hormonal, have a think...

- If the response is <u>really quick</u>, it's <u>probably nervous</u>. Some information needs to be passed to
 effectors really quickly (e.g. pain signals, or information from your eyes telling you about the lion
 heading your way), so it's no good using hormones to carry the message they're too slow.
 - 2) But if a response lasts for a long time, it's probably hormonal. For example, when you get a shock, a hormone called adrenaline is released into the body (causing the fight or flight response, where your body is hyped up ready for action). You can tell it's a hormonal response (even though it kicks in pretty quickly) because you feel a bit wobbly for a while afterwards.

Nerves, hormones — no wonder revision makes me tense...

Hormones control various organs and cells in the body, though they tend to control things that aren't immediately life-threatening (so things like sexual development, blood sugar level, water content, etc.).

Q1 Why is the pituitary gland referred to as the 'master gland'?


[1 mark]

Controlling Blood Glucose

Blood glucose is also controlled as part of homeostasis. Insulin and glucagon are the two hormones involved.

Insulin and Glucagon Control Blood Glucose Level

- 1) Eating foods containing carbohydrate puts glucose (a type of sugar) into the blood from the gut.
- 2) The normal metabolism of cells removes glucose from the blood.
- 3) Vigorous exercise removes much more glucose from the blood.
- 4) Excess glucose can be stored as glucogen in the liver and in the muscles.
- 5) The <u>level of glucose</u> in the <u>blood</u> must be kept <u>steady</u>. <u>Changes</u> are monitored and controlled by the <u>pancreas</u>, using the hormones <u>insulin</u> and <u>glucagon</u>, in a negative feedback cycle:





With Diabetes, You Can't Control Your Blood Sugar Level

Diabetes is a condition that affects your ability to control your blood sugar level. There are two types:

- 1) Tupe 1 diabetes is where the pancreas doesn't produce enough insulin. It may not produce any insulin at all. This means a person's blood glucose level can rise to a level that can kill them. People with Tupe 1 diabetes need insulin therapy this usually involves several injections of insulin throughout the day, most likely at mealtimes. This makes sure that glucose is removed from the blood quickly once the food has been digested, stopping the level getting too high. It's a very effective treatment. The amount of insulin that needs to be injected depends on the person's diet and how active they are.
- 2) Tupe 2 diabetes is where a person becomes resistant to their own insulin (they still produce insulin, but their body's cells don't respond properly to the hormone). This can also cause a person's blood sugar level to rise to a dangerous level. Being overweight can increase your chance of developing Tupe 2 diabetes, as obesity is a major risk factor in the development of the disease. Tupe 2 diabetes can be controlled by eating a carbohudrate-controlled diet and getting regular exercise.

And people used to think the pancreas was just a cushion...

This stuff can seem a bit confusing at first, but if you learn those two diagrams, it should get a bit easier.

Q1 The graph shows the relative secretion rates of insulin and glucagon as the blood glucose level increases. Which curve represents glucagon? Explain your answer. [2 marks]





unregulated amount of

urea is also

lost from the

The Kidneys

The kidneys are really important in this whole homeostasis thing.

Kidneys Basically Act as Filters to "Clean the Blood"

The kidneys make urine by taking <u>waste products</u> (and other unwanted substances) out of your blood. Substances are <u>filtered out</u> of the blood as it passes through the kidneys. This process is called <u>filtration</u>. Useful substances like <u>glucose</u>, some <u>ions</u> and the right amount of <u>water</u> are then absorbed back into the blood. This process is called <u>selective reabsorption</u>. The substances that are removed from the body in <u>urine</u> include:

UREA

- Proteins (and the <u>amino acids</u> that they are broken down into) can't be <u>stored</u> by the body —
 so any <u>excess</u> amino acids are converted into <u>fats</u> and <u>carbohydrates</u>, which can be stored.
 This occurs in the <u>liver</u> and involves a process called <u>deamination</u>.
- 2) Ammonia is produced as a waste product from this process.
- 3) Ammonia is toxic so it's converted to urea in the liver. Urea is then transported to the kidneys, where it's filtered out of the blood and excreted from the body in urine.

2 IONS


- 1) lons such as sodium are taken into the body in food, and then absorbed into the blood.
- 2) If the ion (or water) content of the body is <u>wrong</u>, this could upset the <u>balance</u> between ions and water, meaning too much or too little <u>water</u> is drawn into cells by <u>osmosis</u> (see page 21). Having the wrong amount of water can <u>damage</u> cells or mean they <u>don't work</u> as well as normal.
- 3) Some ions are lost in <u>sweat</u> (which tastes salty, you may have noticed). However, this amount is <u>not regulated</u>, so the right <u>balance</u> of ions in the body must be maintained by the <u>kidneys</u>. The right amount of ions is <u>reabsorbed</u> into the blood after filtration and the rest is removed from the body in urine.

3 WATER

- 1) The body has to constantly balance the water coming in against the water going out.
- 2) We lose water from the skin in sweat and from the lungs when breathing out.
- 3) We can't control how much we lose in these ways, so the amount of water is balanced by the amount we <u>consume</u> and the amount <u>removed by the kidneys</u> in <u>urine</u>.

The Concentration of Urine is Controlled by a Hormone

- 1) The concentration of urine is controlled by a hormone called <u>anti-diuretic hormone</u> (ADH). This is released into the <u>bloodstream</u> by the <u>pituitary gland</u>.
- The brain monitors the water content of the blood and instructs the pituitary gland to release ADH into the blood according to how much is needed.
- 3) The whole process of water content regulation is controlled by <u>negative feedback</u> (see page 65). This means that if the water content gets <u>too high</u> or <u>too low</u> a mechanism will be triggered that brings it back to normal.



Adjusting water content — blood, sweat and, erm, wee...

The kidneys excrete about 547.5 litres of urine a year... that's five baths full. (Not that you should put it there.)

O1 Alcohol suppresses the production of ADH. Suggest how this can lead to dehydration. [2 marks]



Kidney Failure

If someone's kidneys stop working, there are basically two treatments — regular dialysis or a transplant.


The Kidneys Remove Waste Substances from the Blood

- If the kidneys don't work properly, <u>waste substances build up</u> in the blood and you lose your ability to <u>control</u> the <u>levels of ions and water</u> in your body. Eventually, this results in death.
- People with kidney failure can be kept alive by having <u>dialysis treatment</u> —
 where <u>machines</u> do the job of the kidneys. Or they can have a <u>kidney transplant</u>.

The kidneys are incredibly important — if they don't work as they should, you can get problems in the heart, bones, nervous system, stomach, mouth, etc.

Dialysis Machines Filter the Blood

- Dialysis has to be done <u>regularly</u> to keep the concentrations of <u>dissolved substances</u> in the blood at normal levels, and to remove waste substances.
- In a <u>dialysis machine</u> the person's blood flows between <u>partially permeable membranes</u>, surrounded by dialysis fluid. It's permeable to things like <u>ions</u> and <u>waste substances</u>, but not <u>big molecules</u> like proteins (just like the membranes in the kidney).



- The dialysis fluid has the <u>same concentration</u> of dissolved ions and glucose as <u>healthy blood</u>.
- This means that useful <u>dissolved ions</u> and <u>glucose</u> won't be lost from the blood during dialysis.
- 5) Only <u>waste substances</u> (such as <u>urea</u>) and excess ions and water diffuse across the barrier.

■ ← from person ■ ⇒ back to person

- Many patients with kidney failure have to have a dialysis session three times a week.
 Each session takes 3-4 hours — not much fun.
- 7) Plus, dialysis may cause blood clots or infections.
- 8) Being on a dialysis machine is <u>not a pleasant experience</u> and it is <u>expensive</u> for the NHS to run.
- 9) However, dialusis can buy a patient with kidney failure valuable time until a donor organ is found.

Kidney Transplants are a Cure, but can be Rejected

- 1) At the moment, the only <u>cure</u> for kidney failure is to have a <u>kidney transplant</u>.
- 2) Healthy kidneys are usually transplanted from people who have died suddenly.
- The person who died has to be on the <u>organ donor register</u> or carry a <u>donor card</u> (provided their relatives agree too).
- Kidneys can also be transplanted from people who are still alive (as we all have two of them) but there is a small <u>risk</u> to the person donating the kidney.
- 5) There is also a risk that the donor kidney can be rejected by the patient's immune system. The patient is treated with drugs to prevent this but it can still happen.
- 6) Transplants are <u>cheaper</u> (in the long run) than dialysis and they can put an end to the hours patients have to spend on dialysis, but there are long <u>waiting lists</u> for kidneys.



Dialysis or transplant? Both have their downsides...

Donor kidneys are ideally matched by blood type (and a few other things) to the recipient, which make them less likely to be rejected. However, it means a potentially long waiting time for a suitable kidney.

O1 What is the purpose of the partially permeable membrane in a dialysis machine?

[2 marks]

Puberty and the Menstrual Cycle

The monthly release of an egg from a woman's ovaries is part of the menstrual cycle.


Hormones Promote Sexual Characteristics at Puberty

At <u>puberty</u>, your body starts releasing <u>sex hormones</u> that trigger off <u>secondary sexual characteristics</u> (such as the development of <u>facial hair</u> in men and <u>breasts</u> in women) and cause <u>eggs</u> to <u>mature</u> in women.

- In men, the main reproductive hormone is <u>testosterone</u>.
 It's produced by the testes and stimulates sperm production.
- In women, the main reproductive hormone is <u>oestrogen</u>. It's produced by the <u>ovaries</u>.
 As well as bringing about <u>physical changes</u>, oestrogen is also involved in the <u>menstrual cucle</u>.




The Menstrual Cycle Has Four Stages



- Stage 1 Day 1 menstruation starts. The uterus lining breaks down for about four days.
- <u>Stage 2</u> The uterus lining builds up again, from day 4 to day 14, into a thick spongy layer full of blood vessels, ready to receive a fertilised egg.
- Stage 3 An egg develops and is released from the ovary at day 14 this is called ovulation.
- Stage 4 The wall is then maintained for about 14 days until day 28. If no fertilised egg has landed on the uterus wall bu day 28, the spongu lining starts to break down and the whole cucle starts again.

It's Controlled by Four Hormones



- Oestrogen
 - 1) Produced in the ovaries.
 - 2) Causes the lining of the uterus to grow.
 - Stimulates the release of LH (which causes the release of an egg) and inhibits release of FSH.

- FSH (Follicle-Stimulating Hormone)
 - 1) Produced in the pituitary gland.
 - Causes an egg to mature in one of the ovaries, in a structure called a follicle.
 - 3) Stimulates the ovaries to produce oestrogen.
 - 3 LH (Luteinising Hormone)
 - 1) Produced by the pituitary gland.
 - 2) Stimulates the <u>release of an egg</u> at day 14 (<u>ovulation</u>).

- 4 Progesterone
 - 1) Produced in the ovaries by the remains of the follicle after ovulation.
 - Maintains the lining of the uterus during the second half of the cycle.
 When the level of progesterone falls, the lining breaks down.
 - 3) Inhibits the release of LH and FSH.

Which came first — the chicken or the luteinising hormone...

Learn this page until you know which hormone does what, and you understand the graphs.

- Q1 Name the hormone that stimulates an egg to mature in the ovary.
- Q2 Where is testosterone produced in the male body?

[1 mark] [1 mark]

Controlling Fertility

Pregnancy can happen if sperm reaches the ovulated egg. Contraception tries to stop this happening.

Hormones Can Be Used to Reduce Fertility

- 1) Oestrogen can be used to prevent the release of an egg so it can be used as a method of contraception.
- 2) This may seem kind of strange (since naturally oestrogen helps stimulate the release of eggs). But if oestrogen is taken every day to keep the level of it permanently high, it inhibits the production of FSH, and after a while egg development and production stop and stay stopped.
- 3) <u>Progesterone</u> also reduces fertility, e.g. by stimulating the production of thick mucus which prevents any sperm getting through and reaching an egg.
- 4) The pill is an oral contraceptive containing oestrogen and progesterone (known as the combined oral contraceptive pill).
- 5) It's over 99% effective at preventing pregnancy, but it can cause side effects like headaches and nausea and it doesn't protect against sexually transmitted diseases.
- 6) There's also a progesterone-only pill it has fewer side effects than the pill, and is just as effective.
- 7) There are other methods of contraception that use hormones:
 - The contraceptive patch contains oestrogen and progesterone (the same as the combined pill).
 It's a small (5 cm × 5 cm) patch that's stuck to the skin. Each patch lasts one week.
 - The <u>contraceptive implant</u> is inserted <u>under the skin</u> of the arm. It releases a <u>continuous</u> amount of <u>progesterone</u>, which stops the ovaries releasing eggs, makes it hard for sperm to swim to the egg, and stops any fertilised egg implanting in the uterus. An implant can last for <u>three years</u>.
 - The contraceptive injection also contains progesterone. Each dose lasts 2 to 3 months.
 - An intrauterine device (IUD) is a <u>T-shaped</u> device that is inserted into the <u>uterus</u> to <u>kill sperm</u> and <u>prevent implantation</u> of a fertilised egg. There are <u>two</u> main types <u>plastic</u> IUDs that release <u>progesterone</u> and <u>copper</u> IUDs that prevent the sperm <u>surviving</u> in the uterus.

this barrier thing sorted

Barriers Stop Egg and Sperm Meeting

- 1) Non-hormonal forms of contraception are designed to stop the sperm from getting to the egg.
- 2) <u>Condoms</u> are worn over the <u>penis</u> during intercourse to prevent the sperm entering the vagina. There are also <u>female condoms</u> that are worn inside the <u>vagina</u>. Condoms are the only form of contraception that will protect against <u>sexually transmitted diseases</u>.
- 3) A <u>diaphragm</u> is a shallow plastic cup that fits over the <u>cervix</u> (the entrance to the uterus) to form a barrier. It has to be used with <u>spermicide</u> (a substance that <u>disables</u> or <u>kills</u> the sperm).
- 4) Spermicide can be used alone as a form of contraception, but it is not as effective (only about 70-80%).

There are Other Ways to Avoid Pregnancy

STERILISATION — Sterilisation involves cutting or tying the fallopian tubes (which connect the ovaries to the uterus) in a female, or the sperm duct (the tube between the testes and penis) in a male. This is a permanent procedure. However, there is a very small chance that the tubes can rejoin.

'NATURAL' METHODS — Pregnancy may be avoided by finding out <u>when</u> in the menstrual cycle the woman is most <u>fertile</u> and <u>avoiding sexual intercourse</u> on those days. It's popular with people who think that hormonal and barrier methods are <u>unnatural</u>, but it's <u>not very effective</u>.

ABSTINENCE — The only way to be completely sure that sperm and egg don't meet is to not have intercourse.

The winner of best contraceptive ever — just not doing it...

You might be asked to evaluate the different hormonal and non-hormonal methods of contraception in your exam. If you do, make sure you weigh up and write about both the pros and the cons of each method. Exciting stuff.

Name two forms of contraception that reduce fertility by releasing oestrogen. [2 marks]

More on Controlling Fertility

Scientific advances in understanding fertility have led to many infertile women being helped to have babies.

Hormones Can Be Used to Increase Fertility

- 1) Some women have levels of FSH (follicle-stimulating hormone) that are too low to cause their eggs to mature. This means that no eggs are released and the women can't get pregnant.
- 2) The hormones FSH and LH can be given to women in a fertility drug to stimulate ovulation.

It helps a lot of women to get pregnant when previously they couldn't... pretty obvious.

It doesn't always work - some women may have to do it many times, which can be expensive. Too manu eggs could be stimulated, resulting in unexpected multiple pregnancies (twins, triplets, etc.).

IVF Can Also Help Couples to Have Children

If a woman cannot get pregnant using medication, she may chose to try WF ("in vitro fertilisation").

- 1) IVF involves collecting eggs from the woman's ovaries and fertilising them in a lab using the man's sperm.
- 2) IVF treatment can also involve a technique called Intra-Cytoplasmic Sperm Injection (ICSI), where the sperm is injected directly into an egg. It's useful if the man has a very low sperm count.
- 3) The fertilised eggs are then grown into embryos in a laboratory incubator.
- 4) Once the embruos are tinu balls of cells, one or two of them are transferred to the woman's uterus to improve the chance of pregnancu.
- 5) FSH and LH are given before egg collection to stimulate several eggs to mature (so more than one egg can be collected).

PRO Fertility treatment can give an infertile couple a child — a pretty obvious benefit.



Multiple births can happen if more than one embryo grows into a baby — these are risky for the mother and babies (there's a higher risk of miscarriage, stillbirth...).

The success rate of IVF is low — the average success rate in the UK is about 26%. This makes the process incredibly stressful and often upsetting, especially if it ends in multiple failures. As well as being emotionally stressful, the process is also physically stressful for the woman. Some women have a strong reaction to the hormones — e.g. abdominal pain, vomiting, dehudration.

Advances in microscope techniques have helped to improve the techniques (and therefore the success rate) of IVF. Specialised micro-tools have been developed to use on the eggs and sperm under the microscope. They're also used to remove single cells from the embryo for genetic testing (to check that it is healthu see page 93). More recently, the development of time-lapse imaging (using a microscope and camera built into the incubator) means that the growth of the embryos can be continuously monitored to help identify those that are more likely to result in a successful pregnancy.

Some People Are Against IVF

- 1) The process of IVF often results in unused embryos that are eventually destroyed. Because of this, some people think it is unethical because each embryo is a potential human life.
- 2) The genetic testing of embryos before implantation also raises ethical issues as some people think it could lead to the selection of preferred characteristics, such as gender or eye colour.

Nothing funny here, sorry...

Fertility treatment can help to increase the chance of pregnancy, but it can be hard on those involved.

What is the role of FSH and LH during IVF? Q1

O2 Give one drawback to using hormones to increase fertility. [1 mark] [1 mark]

Adrenaline and Thyroxine

You've met a lot of human hormones so far, but two more won't hurt. Then that's it, I promise...

Adrenaline Prepares You for "Fight or Flight"

- 1) Adrenaline is a hormone released by the adrenal glands, which are just above the kidneys (see p.73).
- Adrenaline is released in response to <u>stressful or scary situations</u> —
 your brain detects fear or stress and sends <u>nervous impulses</u> to the
 adrenal glands, which respond by secreting adrenaline.
- 3) It gets the body ready for 'fight or flight' by triggering mechanisms that increase the supply of oxygen and glucose to cells in the brain and muscles. For example, adrenaline increases heart rate.




Hormone Release can be Affected by Negative Feedback

Your body can control the levels of hormones (and other substances) in the blood using negative feedback systems. When the body detects that the level of a substance has gone above or below the normal level, it triggers a response to bring the level back to normal again. Here's an example of just that:

Thyroxine Regulates Metabolism

- Thyroxine is a hormone released by the thyroid gland, which is in the neck (see p.73).
- Thyroxine is made in the thyroid gland E from iodine and amino acids.
- 2) It plays an important role in regulating the <u>basal metabolic rate</u> the speed at which chemical reactions in the body occur while the body is at <u>rest</u>. Thyroxine is also important for loads of processes in the body, such as stimulating <u>protein synthesis</u> for growth and <u>development</u>.
- Thyroxine is released in response to thyroid stimulating hormone (TSH), which is released from the pituitary gland.
- 4) A <u>negative feedback system</u> keeps the amount of thyroxine in the blood at the right level when the level of thyroxine in the blood is <u>higher than normal</u>, the secretion of <u>TSH</u> from the pituitary gland is <u>inhibited</u> (stopped). This <u>reduces</u> the amount of <u>thyroxine</u> released from the thyroid gland, so the level in the blood <u>falls</u> back towards <u>normal</u>.



Negative feedback sucks, especially from your science teacher...

You can think about negative feedback working like a thermostat — if the temperature gets too low, the thermostat will turn the heating on, then if the temperature gets too high, it'll turn the heating off again.

- Q1 Name the gland that releases thyroxine.
- Describe the response if the level of thyroxine in the blood gets too high.

[1 mark] [3 marks]

Plant Hormones

Plants don't just grow randomly. Plant hormones make sure they grow in the right direction.

Auxin is a Plant Growth Hormone

- 1) Auxin is a plant hormone that controls growth near the tips of shoots and roots.
- 2) It controls the growth of a plant in response to <u>light</u> (<u>phototropism</u>) and <u>gravity</u> (<u>gravitropism</u> or <u>geotropism</u>).
- Auxin is produced in the tips and moves backwards to stimulate the cell elongation (enlargement) process which occurs in the cells just behind the tips.
- 4) If the tip of a shoot is removed, no auxin is available and the shoot may stop growing.
- 5) Extra auxin promotes growth in the shoot but inhibits growth in the root producing the desired result...


Shoots grow towards light

- When a <u>shoot tip</u> is exposed to <u>light</u>, <u>more auxin</u> accumulates on the side that's in the <u>shade</u> than the side that's in the light.
- This makes the cells grow (elongate) <u>faster</u> on the <u>shaded side</u>, so the shoot bends <u>towards</u> the light.



Shoots grow away from gravity and roots grow towards gravity

- When a <u>shoot</u> is growing sideways, <u>gravity</u> produces an unequal distribution of auxin in the tip, with <u>more auxin</u> on the <u>lower side</u>.
- 2) This causes the lower side to grow faster, bending the shoot upwards.
- 3) A root growing sideways will also have more auxin on its lower side.
- 4) But in a root the <u>extra</u> auxin <u>inhibits</u> growth. This means the cells on top elongate faster, and the root bends downwards.



You can Investigate Plant Growth Responses

PRACTICAL

You can investigate the effect of <u>light</u> on the <u>growth</u> of cress seeds like this...

1) Put 10 cress seeds into three different Petri dishes, each lined with <u>moist filter pape</u>

(Remember to label your dishes, e.g. A, B, C.)
2) Shine a <u>light</u> onto one of the dishes from <u>above</u>





4) You know that the growth response of the cress seeds is due to light only, if you control all other variables.

Examples of variables

to control are:

You can also investigate the effect of gravity

You can also investigate the effect of gravity on plant growth. Just place four seedlings or damp cotton wool in a Petri dish, each with their roots pointing in a different direction, and store the Petri dish vertically for a few days in the dark. You should find that the roots of each seedling grow downwards.

٠	VARIABLE	HOW TO CONTROL IT
	number of seeds	use the same number of seeds in each dish
	type of seed	use seeds that all come from the same packet
	temperature	keep your Petri dishes in a place where the temperature is stable (i.e. away from heat sources and draughts)
	water	use a measuring cylinder to add the same amount of water
	light intensity	keep the distance between the bulb and dish the same

A plant auxin to a bar — 'ouch'...

You need to be able to draw labelled diagrams to show the results of your experiment. Make sure you do it neatly.

Q1 Explain what causes plant shoots to grow towards light.

[2 marks]

Commercial Uses of Plant Hormones

Plant hormones can be extracted, or artificial copies can be made. They can then be used to do all kinds of useful things, including killing weeds, growing cuttings and ripening fruit.

Auxins Have Many Uses

Auxins are useful for controlling plant growth. Here are some of the ways they come in handy:

1) KILLING WEEDS — Most weeds growing in fields of crops or in a lawn are broad-leaved, in contrast to grasses and gereals which have very narrow leaves. Selective weedkillers have been developed using auxins, which only affect the broad-leaved plants. They totally disrupt their normal growth patterns, which soon kills them, whilst leaving the grass and crops untouched.



- 2) GROWING FROM CUTTINGS WITH ROOTING POWDER A <u>cutting</u> is part of a plant that has been <u>cut off it</u>, like the end of a branch with a few leaves on it. Normally, if you stick cuttings in the soil they <u>won't grow</u>, but if you add <u>rooting powder</u>, which contains <u>auxins</u>, they will <u>produce roots</u> rapidly and start growing as <u>new plants</u>. This enables growers to produce lots of <u>clones</u> (exact copies) of a really good plant <u>yery quickly</u>.
- 3) GROWING CELLS IN TISSUE CULTURE <u>Tissue culture</u> can be used to grow <u>clones</u> of a plant from a few of its cells. To do this, hormones such as auxins need to be added to the <u>growth medium</u> (along with nutrients) to stimulate the cells to divide to form both roots and shoots.

Gibberellin Stimulates Plant Stems to Grow

Gibberellin is another type of plant growth hormone. It stimulates seed germination, stem growth and flowering. Its uses include:

Seed germination is when a seed = starts to grow into a plant.

- CONTROLLING DORMANCY Lots of seeds won't germinate until they've been through certain
 conditions (e.g. a period of cold or or druness). This is called dormancy. Seeds can be treated
 with gibberellin to alter dormancy and make them germinate at times of year that they wouldn't
 normally. It also helps to make sure all the seeds in a batch germinate at the same time.
- 2) INDUCING FLOWERING Some plants require certain conditions to flower, such as longer days or low temperatures. If these plants are treated with gibberellin, they will flower without any change in their environment. Gibberellin can also be used to grow bigger flowers.
- 3) GROWING LARGER FRUIT <u>Geedless varieties</u> of fruit (e.g. seedless grapes) often do not grow as large as seeded fruit. However, if <u>gibberellin</u> is added to these fruits, they will grow larger to match the normal tubes.

Ethene Stimulates Ripening of Fruit

 Ethene is a gas produced by aging parts of a plant. It influences the growth of the plant by controlling cell division. It also <u>stimulates enzymes</u> that cause <u>fruit to ripen</u>.



- Commercially, it can be used to speed up the ripening of fruits either while they are still on the plant, or during transport to the shops.
- 3) This means that fruit can be picked while it's still <u>unripe</u> (and therefore firmer and <u>less easily damaged</u>). The gas is then added to the fruit on the way to the supermarket so that it will be <u>perfect</u> just as it reaches the shelves.
- 4) Ripening can also be delayed while the fruit is in storage by adding chemicals that block etheno's effect on the fruit or reduce the amount of ethene that the fruit can produce. Alternatively, some chemicals can be used that react with ethene to remove it from the air.

You will ripen when I SAY you can ripen — and NOT BEFORE...

Three different hormones, many different uses. Bet you didn't know plant hormones could be so useful...

Of the one way that gibberellin is used commercially.

[1 mark]

Revision Questions for Topic 5

Congratulations, you've finished Topic 5 — now for some questions...

- Try these questions and tick off each one when you get it right.
- When you're completely happy with a sub-topic, tick it off.

30) Which plant hormone is responsible for fruit ripening?

For even more practice, try the Retrieval Quiz for Topic 5 — just Scan this QR code!




Homeostasis and the Nervous System (p.65-68)			
1) What is a stimulus?	\checkmark		
2) Explain how negative feedback helps to maintain a stable internal environment.	\checkmark		
3) What makes up the central nervous system and what does it do?	\checkmark		
4) What is a synapse?	\checkmark		
5) What is the purpose of a reflex action?	√		
6) What is a reaction time?			
The Brain, the Eye and Body Temperature (p.69-72)			
7) Give one thing the cerebral cortex is responsible for.	\checkmark		
8) Give three methods used by scientists to study the brain.	<u></u>		
9) Explain the roles of the following parts of the eye:	_		
a) cornea b) iris c) lens	√		
10) Describe the iris reflex. Why is this needed?	<u> </u>		
11) What is the medical term for short-sightedness?	\checkmark		
12) Give three things that the body can do to reduce heat loss if it gets too cold.	\checkmark		
Hormones in Humans (p.73-80)			
13) What is a hormone?	\checkmark		
14) Give two differences between nervous and hormonal responses.			
15) Where is excess glucose stored in the body?	\checkmark		
16) What effect does the hormone glucagon have on blood glucose level?	\checkmark		
17) What is the difference between how type 1 and type 2 diabetes are usually controlled?	\checkmark		
18) Name three things that are reabsorbed by kidneys.	\checkmark		
19) What hormone controls the amount of water reabsorbed by the kidneys?	\checkmark		
20) What are the advantages and disadvantages of a kidney transplant over dialysis?	✓		
21) Draw a timeline of the 28 day menstrual cycle.	_		
Label the four stages of the cycle and label when the egg is released.			
22) Describe two effects of FSH on the body.	<u>√</u>		
23) Which of the following is a hormonal contraceptive — condom, plastic IUD or diaphragm?	<u>√</u>		
24) Briefly describe how IVF is carried out.	\checkmark		
25) How does adrenaline prepare the body for 'fight or flight'?	\checkmark		
Plant Hormones (p.81-82)	_		
26) What is auxin?	✓		
27) What is: a) phototropism? b) gravitropism?	\checkmark		
28) Explain how auxin causes plant roots to grow downwards.	\checkmark		
29) Give three ways that auxin can be used commercially.	\checkmark		

DNA

The first step in understanding genetics is getting to grips with DNA and genes.

Chromosomes Are Really Long Molecules of DNA

- 1) DNA stands for deoxyribonucleic acid. It's the chemical that all of the genetic material in a cell is made up from.
- 2) It contains coded information basically all the instructions to put an organism together and make it work.
- Head to the next page for more on the structure of DNA. STITLE OF THE STREET OF STREET
- 3) So it's what's in your DNA that determines what inherited characteristics you have.
- 4) DNA is found in the nucleus of animal and plant cells, in really long structures called chromosomes.
- 5) Chromosomes normally come in pairs.
- 6) DNA is a polumer. It's made up of two strands coiled together in the shape of a double helix.



A Gene Codes for a Specific Protein

- 1) A gene is a small section of DNA found on a chromosome.
- 2) Each gene codes for (tells the cells to make) a particular sequence of amino acids which are put together to make a specific protein.
- 3) Only 20 amino acids are used, but they make up thousands of different proteins.
- 4) Genes simply tell cells in what order to put the amino acids together (more on this on the next page).
- 5) DNA also determines what proteins the cell produces, e.g. haemoglobin, keratin.
- 6) That in turn determines what type of cell it is, e.g. red blood cell, skin cell.

Every Organism Has a Genome

- 1) Genome is just the fancy term for the entire set of genetic material in an organism.
- 2) Scientists have worked out the complete human genome.
- 3) Understanding the human genome is a really important tool for science and medicine for many reasons.
 - 1) It allows scientists to identify genes in the genome that are linked to different types of disease.
 - 2) Knowing which genes are linked to inherited diseases could help us to understand them better and could help us to develop effective treatments for them.
 - 3) Scientists can look at genomes to trace the migration of certain populations of people around the world. All modern humans are descended from a common ancestor who lived in Africa, but humans can now be found all over the planet. The human genome is mostly identical in all individuals, but as different populations of people migrated away from Africa, they gradually developed tiny differences in their genomes. By investigating these differences, scientists can work out when new populations split off in a different direction and what route they took.

Insert joke about genes and jeans here...

There are so many, I thought you could come up with your own as a bit of light relief. Make sure that you're clued up on this stuff about DNA, genes and proteins before you move on.

Q1 What is a gene?

What is an organism's genome?

[1 mark]

Topic 6 — Inheritance, Variation and Evolution

[3 marks]



A DNA molecule

with a double

helix structure

(a doublestranded spiral).

Part of a DNA strand sugar-phosphate backbone

nucleotide

base on one strand is joined

to a base on

the other

etrand

base

Part of a

DNA molecule

The Structure of DNA and Protein Synthesis

So here's how life works — DNA molecules contain a genetic code that determines which proteins are built.

DNA is Made Up of Nucleotides

- DNA strands are <u>polymers</u> made up of lots of repeating units called <u>nucleotides</u>.
- Each nucleotide consists of a sugar, a phosphate group and one 'base'.
- The <u>sugar</u> and <u>phosphate</u> groups in the nucleotides form a '<u>backbone</u>' to the DNA strands. The sugar and phosphate groups <u>alternate</u>. One of <u>four</u> different bases — A, T, C or G — joins to each sugar.
- 4) Each base links to a base on the opposite strand in the helix.
- 5) A always pairs up with T, and C always pairs up with G. This is called complementary base pairing.
- 6) It's the <u>order</u> of <u>bases</u> in a <u>gene</u> that decides the order of <u>amino acids</u> in a <u>protein</u>.
- 7) Each amino acid is coded for by a sequence of three bases in the gene.
- The amino acids are joined together to make various proteins, depending on the order of the gene's bases.
- 9) There are parts of DNA that don't code for proteins. Some of these non-coding parts switch genes on and off, so they control whether or not a gene is expressed (used to make a protein).

mRNA Carries The Code to The Ribosomes

- 1) Proteins are made in the cell <u>cutoplasm</u> on tiny structures called <u>ribosomes</u>.
- 2) To make proteins, ribosomes use the code in the DNA. DNA is found in the cell nucleus and can't move out of it because it's really big. So the cell needs to get the code from the DNA to the ribosome.
- 3) This is done using a molecule called mRNA which is made by copying the code from DNA. The mRNA acts as a messenger between the DNA and the ribosome it carries the code between the two.
- 4) The correct amino acids are brought to the ribosomes in the correct order by carrier molecules.

Proteins Have Many Different Functions

When a chain of amino acids has been assembled, it <u>folds</u> into a <u>unique shape</u> which allows the protein to <u>perform</u> the task it's meant to do. Here are a few examples of types of protein:

- 1) ENZYMES act as biological catalysts to speed up chemical reactions in the body (see page 28).
- HORMONES used to <u>carry messages</u> around the body. E.g. <u>insulin</u> is a hormone released into the blood by the pancreas to regulate the blood sugar level.
- STRUCTURAL PROTEINS are physically <u>strong</u>. E.g. <u>collagen</u> is a structural protein that strengthens <u>connective tissues</u> (like ligaments and cartilage).

Q: What do DNA and a game of rounders have in common?

A: four bases. Remember — the order of these bases in the DNA is copied into mRNA, which then moves to the ribosomes. Here, the chain of amino acids is assembled according to the order of the bases.



How many amino acids does it code for?

b) Give the sequence of bases for the complementary strand.

[1 mark] [1 mark]

strand. [1 mark]

Mutations

Sometimes the sequence of DNA bases can be changed. These changes are called mutations. Read on...

Mutations are Changes to the Genetic Code

- Occasionally a gene may <u>mutate</u>. A mutation is a random <u>change</u> in an organism's <u>DNA</u>. They can sometimes be <u>inherited</u>.
- Mutations occur <u>continuously</u>. They can occur <u>spontaneously</u>, e.g. when a chromosome isn't quite replicated properly. However, the <u>chance</u> of mutation is <u>increased</u> by exposure to certain <u>substances</u> or <u>some types of radiation</u>.
- 3) Mutations change the sequence of the <u>DNA bases</u> in a gene, which produces a <u>genetic variant</u> (a different form of the gene). As the <u>sequence</u> of DNA bases <u>codes</u> for the sequence of <u>amino acids</u> that make up a <u>protein</u> (see page 84), mutations to a gene <u>sometimes</u> lead to <u>changes</u> in the protein that it codes for.
- Most mutations have very little or no effect on the protein. Some will change
 it to such a small extent that its function or appearance is unaffected.
- 5) However, some mutations can <u>seriously affect</u> a protein. Sometimes, the mutation will code for an altered protein with a change in its <u>shape</u>. This could affect its ability to perform its <u>function</u>. E.g.
 - 1) If the shape of an enzyme's active site is changed, its substrate may no longer be able to bind to it.
 - Enzymes are proteins see page 28.
 - Structural proteins like collagen could lose their strength if their shape is changed, making them pretty useless at providing structure and support.
- 6) If there's a mutation in the <u>non-coding DNA</u>, it can alter how genes are <u>expressed</u>.

There's more on a non-coding DNA on the previous page.

There Are Different Types of Mutation, For Example...

Insertions

- Insertions are where a <u>new base</u> is <u>inserted</u> into the DNA base sequence where it shouldn't be.
- You should remember from page 85 that every three bases in a DNA base sequence codes for a particular amino acid.
- An insertion changes the way the groups of three bases are 'read', which can change the amino acids that they code for.
- Insertions can change more than one amino acid as they have a knock-on effect on the bases further on in the sequence.



Deletions

- Deletions are when a random base is deleted from the DNA base sequence.
- Like insertions, they change the way that the base sequence is 'read' and have knock-on effects further down the sequence.

Substitutions

<u>Substitution</u> mutations are when a random base in the DNA base sequence is <u>changed</u> to a different base.



Mutations — sometimes known to give ninja superpowers...

Remember, changes in bases can affect the amino acid sequence, and therefore the protein coded for.

- Q1 What is a mutation? [1 mark]
- O2 Explain why the function of a protein may be affected by a gene mutation.

[3 marks]

Reproduction


Ooo err, reproduction... Surely you knew it'd come up at some point. It can happen in two different ways...

Sexual Reproduction Produces Genetically Different Cells

- Sexual reproduction is where genetic information from two organisms (a father and a mother) is combined to produce offspring which are genetically different to either parent.
- In sexual reproduction, the mother and father produce gametes by meiosis (see next page) — e.g. egg and sperm cells in animals.
- 3) In humans, each gamete contains <u>23 chromosomes</u> half the number of chromosomes in a normal cell. (Instead of having two of each chromosome, a gamete has just one of each.)
- 4) The egg (from the mother) and the sperm cell (from the father) then <u>fuse together</u> (<u>fertilisation</u>) to form a cell with the full number of chromosomes (half from the father, half from the mother).

<u>SEXUAL REPRODUCTION</u> involves the fusion of male and female gametes.

Because there are <u>TWO</u> parents, the offspring contain a mixture of their parents' genes.



- 5) This is why the offspring inherits features from both parents it's received a mixture of chromosomes from its mum and its dad (and it's the chromosomes that decide how you turn out).
- 6) This mixture of genetic information produces variation in the offspring. Pretty cool, eh.
- 7) Flowering plants can reproduce in this way too. They also have egg cells, but their version of sperm is known as pollen. Hmm... I'm having second thoughts about frolicking in that meadow now.

Asexual Reproduction Produces Genetically Identical Cells

- 1) In asexual reproduction there's only one parent so the offspring are genetically identical to that parent.
- Asexual reproduction happens by <u>mitosis</u> an <u>ordinary cell</u> makes a new cell by <u>dividing in two</u> (see page 15).
- The new cell has exactly the same genetic information (i.e. genes) as the parent cell
 — it's called a clone.

In <u>ASEXUAL REPRODUCTION</u> there's only <u>ONE</u> parent. There's <u>no fusion</u> of gametes, <u>no mixing</u> of chromosomes and <u>no genetic variation</u> between parent and offspring. The offspring are <u>genetically identical</u> to the parent — they're <u>clones</u>.

Bacteria, some plants and some animals reproduce asexually.



You need to reproduce these facts in the exam...

The main messages on this page are that: 1) sexual reproduction needs two parents and forms cells that are genetically different to the parents, so there's lots of genetic variation. And 2) asexual reproduction needs just one parent to make genetically identical cells, so there's no genetic variation in the offspring.

Q1 What type of cell division is involved in asexual reproduction?

[1 mark]

O2 Suggest why there is variation in the offspring of sexual reproduction.

[2 marks]


Meiosis

Now I bet you're wondering how gametes end up with half the number of chromosomes of a normal cell...

Gametes Are Produced by Meiosis

- As you know from the previous page, gametes only have one copy of each <u>chromosome</u>, so that when gamete <u>fusion</u> takes place, you get the <u>right amount</u> of <u>chromosomes</u> again (two copies of each).
- 2) To make gametes which only have <u>half</u> the original number of chromosomes, cells divide by <u>meiosis</u>. This process involves <u>two cell divisions</u>. In humans, it <u>only</u> happens in the <u>reproductive organs</u> (the ovaries in females and testes in males).

Meiosis Produces Cells With Half the Normal Number of Chromosomes



Before the cell starts to divide, it <u>duplicates</u> its <u>genetic</u> <u>information</u>, forming two armed chromosomes — one arm of each chromosome is an <u>exact copy</u> of the other arm. After replication, the chromosomes arrange themselves into <u>pairs</u>.

The genetic information is stored in IDNA — Is see p.84.

In the <u>first division</u> in meiosis the chromosome pairs <u>line up</u> in the centre of the cell.

The pairs are then <u>pulled apart</u> so each new cell only has one copy of each chromosome. <u>Some</u> of the father's chromosomes (shown in blue) and <u>some</u> of the mother's chromosomes (shown in red) go into each new cell.

In the <u>second division</u>, the chromosomes <u>line up</u> again in the centre of the cell.

The arms of the chromosomes are <u>pulled apart</u>.

You get four gametes, each with only a <u>single set</u> of chromosomes in it. Each of the gametes is <u>genetically different</u> from the others because the chromosomes all get <u>shuffled up</u> during meiosis and each gamete only gets half of them, at random.

The Cell Produced by Gamete Fusion Replicates Itself

- After two gametes have fused during fertilisation, the resulting new cell divides by mitosis to make a copu of itself.
- 2) Mitosis repeats many times to produce lots of new cells in an embryo.
- As the embryo develops, these cells then start to differentiate (see page 14) into the different tupes of specialised cell that make up a whole organism.

There's loads on 3 = mitosis on page 15.

Now that I have your undivided attention...

In humans, meiosis only occurs in reproductive organs, for making gametes.

Q1 Human body cells contain 46 chromosomes each. The graph on the right shows how the mass of DNA per cell changed as some cells divided by meiosis in a human ovary. How many chromosomes were present in each cell when they reached stage 6? [I mark]



More on Reproduction

As the line in Oliver Twist goes, "Please sir, I want some more... information about the types of reproduction". Something like that anyway. I'm feeling like a generous workhouse master today, so here's a full page.

Sexual Reproduction Has Advantages Over Asexual Reproduction

- Offspring from sexual reproduction have a <u>mixture</u> of two sets of chromosomes. The organism inherits genes (and therefore features) from <u>both</u> parents, which produces <u>variation</u> in the offspring (see p.87).
- 2) Variation increases the chance of a species <u>surviving</u> a <u>change in the environment</u>. While a change in the environment could <u>kill</u> some <u>individuals</u>, it's likely that <u>variation</u> will have led to <u>some</u> of the offspring being able to <u>survive</u> in the new environment. They have a <u>survival advantage</u>.
- 3) Because individuals with characteristics that make them <u>better adapted</u> to the environment have a <u>better</u> chance of <u>survival</u>, they are <u>more likely</u> to breed <u>successfully</u> and pass the genes for the characteristics on. This is known as natural selection (see p.96).
- 4) We can use selective breeding to speed up natural selection. This allows us to produce animals with desirable characteristics. Selective breeding is where individuals with a desirable characteristic are bred to produce offspring that have the desirable characteristic too (see p.98). This means that we can increase food production, e.g. by breeding animals that produce a lot of meat.

Asexual Reproduction Has Advantages Over Sexual Reproduction

- 1) There only needs to be one parent.
- This means that asexual reproduction uses less energy than sexual reproduction, because organisms don't have to find a mate.
- 3) This also means that asexual reproduction is faster than sexual reproduction.
- 4) Many identical offspring can be produced in favourable conditions.

Some Organisms Can Reproduce by Both Methods

Some organisms can reproduce sexually or asexually depending on their <u>circumstances</u>. You need to know these examples:

- Malaria is caused by a parasite that's spread by mosquitoes see page 47. When a mosquito
 carrying the parasite bites a human, the parasite can be transferred to the human. The parasite
 reproduces sexually when it's in the mosquito and asexually when it's in the human host.
- 2) Many species of fungus can reproduce both sexually and asexually. These species release <u>spores</u>, which can become <u>new</u> fungi when they land in a suitable place. Spores can be produced sexually and asexually. <u>Asexually-produced</u> spores form fungi that are <u>genetically identical</u> to the parent fungus. <u>Sexually-produced</u> spores introduce <u>variation</u> and are often produced in response to an unfavourable <u>change</u> in the <u>environment</u>, increasing the chance that the population will <u>survive</u> the change.
- 3) Loads of species of plant produce seeds sexually, but can also reproduce asexually.

 Asexual reproduction can take place in different ways. For example, strawberry plants produce runners. These are stems that grow horizontally on the surface of the soil away from a plant. At various points along the runner, a new strawberry plant forms that is identical to the original plant. Another example is in plants that grow from bulbs (e.g. daffodils).

 New bulbs can form from the main bulb and divide off. Each new bulb can grow into a new identical plant.

Well that's reproduction sorted... Well that's reproduction sorted...

It's key here to remember that both types of reproduction have their own different advantages. Make sure you also know about organisms that can reproduce by both methods.

O1 Give three advantages of asexual reproduction over sexual reproduction.

[3 marks]

X and Y Chromosomes


Now for a couple of very important little chromosomes...

Your Chromosomes Control Whether You're Male or Female

There are <u>23 pairs</u> of chromosomes in every human body cell (page 15). Of these, <u>22</u> are <u>matched pairs</u> of chromosomes that just control <u>characteristics</u>. The <u>23rd pair</u> are labelled <u>XY</u> or <u>XX</u>. They're the two chromosomes that <u>decide</u> your <u>sex</u> — whether you turn out <u>male</u> or <u>female</u>.

Males have an X and a Y chromosome: XY
The Y chromosome causes male characteristics.
Females have two X chromosomes: XX

The XX combination allows female characteristics to develop.




When making sperm, the X and Y chromosomes are drawn apart in the first division in meiosis (see page 88). There's a 50% chance each sperm cell gets an X-chromosome and a 50% chance it gets a Y-chromosome.


A similar thing happens when making eggs. But the original cell has two X-chromosomes, so all the eggs have one X-chromosome.

Genetic Diagrams Show the Possible Gamete Combinations

- 1) To find the probability of getting a boy or a girl, you can draw a genetic diagram.
- Genetic diagrams are just models that are used to show all the possible genetic <u>outcomes</u> when you cross together different genes or chromosomes.
- Put the possible gametes (eggs or sperm) from one parent down the side, and those from the other parent along the top.
- 4) Then in each middle square you fill in the letters from the top and side that line up with that square. The pairs of letters in the middle show the possible combinations of the gametes.



- 5) There are two XX results and two XY results, so there's the same probability of getting a boy or a girl.
- 6) Don't forget that this 50:50 ratio is only a probability at each pregnancy.



The other type of genetic diagram looks a bit more complicated, but it shows exactly the same thing.

- 1) At the top are the parents.
- The middle circles show the <u>possible gametes</u> that are formed. One gamete from the female combines with one gamete from the male (during fertilisation).
- 3) The criss-cross lines show all the possible ways the X and Y chromosomes could combine. The possible combinations of the offspring are shown in the bottom circles.
- Remember, only one of these possibilities would actually happen for any one offspring.

Have you got the Y-factor...

Most genetic diagrams you'll see in exams concentrate on a gene instead of a chromosome. But the principle's the same. Don't worry — there are loads of other examples on the following pages.

What combination of sex chromosomes do human females have?

[1 mark]

Genetic Diagrams

Genetic diagrams, eh. They're not as scary as they look — you just need to practise them...

Some Characteristics are Controlled by Single Genes


- 1) What genes you inherit control what characteristics you develop.
- <u>Different</u> genes control <u>different</u> characteristics. <u>Some</u> characteristics are controlled by a <u>single</u> gene, e.g. <u>mouse fur colour</u> and <u>red-green colour blindness</u> in humans.
- 3) However, most characteristics are controlled by several genes interacting.
- 4) All genes exist in different versions called alleles (which are represented by letters in genetic diagrams).
- 5) You have two versions (alleles) of every gene in your body one on each chromosome in a pair.
- 6) If an organism has two alleles for a particular gene that are the same, then it's homozygous for that trait. If its two alleles for a particular gene are different, then it's heterozygous.
- 7) If the two alleles are <u>different</u>, only one can determine what <u>characteristic</u> is present. The allele for the <u>characteristic that's shown</u> is called the <u>dominant</u> allele (use a capital letter for dominant alleles e.g. 'C'). The other one is called recessive (and you show these with small letters e.g. 'c').
- 8) For an organism to display a <u>recessive</u> characteristic, <u>both</u> its alleles must be <u>recessive</u> (e.g. cc). But to display a <u>dominant</u> characteristic the organism can be <u>either</u> CC or Cc, because the dominant allele overrules the recessive one if the plant/animal/other organism is heterozygous.
- Your <u>genotype</u> is the combination of <u>alleles</u> you have. Your alleles work at a molecular level to determine what characteristics you have — your <u>phenotype</u>.

Genetic Diagrams Show the Possible Alleles of Offspring


Suppose you start breeding <u>hamsters</u> with <u>superpowers</u>. The allele which causes hamsters to have superpowers is recessive ("b"), whilst normal (boring) behaviour is due to a dominant allele ("B").

1) A superpowered hamster must have the genotype bb. But a normal hamster could be BB or Bb.

2) Here's what happens if you breed from two homozugous hamsters:



3) If two of these offspring now breed, you'll get the next generation:



4) That's a 3:1 ratio of normal to superpowered offspring in this generation (a 1 in 4 or 25% probability of superpowers).

Your meanotype determines how nice you are to your sibling...

You need to be able to produce and interpret both of these types of genetic diagram for the exam.

O1 Define genotype and phenotype.

[2 marks]

More Genetic Diagrams

You've got to be able to predict and explain the outcomes of crosses between individuals for each possible combination of dominant and recessive alleles of a gene. You should be able to draw a genetic diagram and work it out — but it'll be easier if you've seen them all before. So here are a couple more examples for you. You also need to know how to interpret another tupe of genetic diagram called a family tree...

All the Offspring are Normal


Let's take another look at the superpowered hamster example from page 91:


In this cross, a homozugous dominant hamster (BB) is crossed with a homozugous recessive hamster (bb). All the offspring are normal (boring).

But, if you crossed a homozygous dominant hamster (BB) with a heterozugous hamster (Bb). you would also get all normal (boring) offspring.

For a reminder on the terms homozygous

and heterozygous, head to page 91.






To find out which it was you'd have to breed the offspring together and see what kind of ratio you got that time — then you'd have a good idea. If it was 3:1, it's likely that you originally had BB and bb.

There's a 1:1 Ratio in the Offspring

A cat with long hair was bred with another cat with short hair. The long hair is caused by a dominant allele 'H', and the short hair by a recessive allele 'h'.

They had 8 kittens — 4 with long hair and 4 with short hair. This is a 1:1 ratio — it's what you'd expect when a parent with only one dominant allele (heterozygous - Hh) is crossed with a parent with two recessive alleles (homozygous recessive — hh).



You Need to be Able to Interpret Family Trees

Knowing how inheritance works can help you to interpret a family tree — this is one for cystic fibrosis (p.93).

- 1) From the family tree, you can tell that the allele for custic fibrosis isn't dominant because plenty of the family carry the allele but don't have the disorder.
- 2) There is a 25% chance that the new babu will have the disorder and a 50% chance that it will be a carrier, as both of its parents are carriers but are unaffected. The case of the new baby is just the same as in the genetic diagram on page 93 - so

John Susan Key Cystic fibrosis Fve unaffected ar new babu

the baby could be unaffected (FF), a carrier (Ff) or have cystic fibrosis (ff).

It's enough to make you go cross-eyed...

In the exam, you might get a family tree showing the inheritance of a dominant allele — in this case, there won't be any carriers shown. Now, here's a fascinating practice question about peas...




01 Round peas are caused by the dominant allele, R. The allele for wrinkly peas, r, is recessive. Using a Punnett square, predict the ratio of plants with round peas to plants with wrinkly peas for a cross between a heterozygous pea plant and a pea plant that is homozygous recessive. [3 marks]

Inherited Disorders

Some disorders can be inherited from your parents. Many of these can be screened for in embryos.

Cystic Fibrosis is Caused by a Recessive Allele

<u>Custic fibrosis</u> is a <u>genetic disorder</u> of the <u>cell membranes</u>. It <u>results</u> in the body producing a lot of thick sticky <u>mucus</u> in the <u>air passages</u> and in the <u>pancreas</u>.




- The allele which causes cystic fibrosis is a <u>recessive allele</u>. 'f', carried by about 1 <u>berson in 25</u>.
- Because it's recessive, people with only one copy of the allele won't have the disorder — they're known as carriers.
- 3) For a child to have the disorder, both parents must be either carriers or have the disorder themselves.
- 4) As the diagram shows, there's a <u>lin 4 chance</u> of a child having the disorder if <u>both</u> parents are <u>carriers</u>.

Polydactyly is Caused by a Dominant Allele

Polydactyly is a genetic disorder where a baby's born with extra fingers or toes.

It doesn't usually cause any other problems so isn't life-threatening.

- The disorder is caused by a dominant allele, 'D', and so can be inherited if just one parent carries the defective allele.
- The parent that has the defective allele will have the condition too since the allele is dominant.
- As the genetic diagram shows, there's a 50% chance of a child having the disorder if one parent has one D allele.



Embryos Can Be Screened for Genetic Disorders

- During in vitro fertilisation (IVF), embryos are fertilised in a laboratory, and then implanted into the mother's womb.
- 2) Before being implanted, it's possible to remove a cell from each embryo and analuse its genes.
- 3) Many genetic disorders can be detected in this way, such as cystic fibrosis.
- 4) It's also possible to get DNA from an embryo in the womb and test that for disorders.
- 5) There are lots of ethical, social and economic concerns surrounding embryo screening.
- 6) Embryonic screening is quite controversial because of the decisions it can lead to.
- 7) For embryos produced by IVF after screening, embryos with 'bad' alleles would be destroyed.
- 8) For embryos in the womb screening could lead to the decision to terminate the pregnancy.
- 9) Here are some more arguments for and against screening:

Against Embryonic Screening

- It implies that <u>people</u> with <u>genetic problems</u> are 'undesirable' — this could increase prejudice.
- There may come a point where everyone wants to screen their embryos so they can pick the most 'desirable' one, e.g. they want a blue-eyed, blond-haired, intelligent boy.
- 3) Screening is expensive.

For Embryonic Screening

- It will help to stop people suffering.
- Treating disorders costs the Government (and the taxpayers) a lot of money.
- There are laws to stop it going too far.
 At the moment parents cannot even select the sex of their baby (unless it's for health reasons).

Embryo screening — it's a tricky one...

Try writing a balanced argument for and against embryo screening — it's good practice.

O1 Why won't someone heterozygous for the cystic fibrosis allele have the disorder?

[3 marks]


The Work of Mendel

Some people forget about Mendel but I reckon he's the Granddaddy of Genetics. Here's a whole page on him.

Mendel Did Genetic Experiments with Pea Plants

Gregor Mendel was an Austrian monk who trained in mathematics and natural history at the University of Vienna. On his garden plot at the monastery in the mid 19th century, Mendel noted how characteristics in plants were passed on from one generation to the next. The results of his research were published in 1866 and eventually became the foundation of modern genetics.

The diagrams show two crosses for height in pea plants that Mendel carried out...



Mendel had shown that the height characteristic in pea plants was determined by separately inherited "hered lary units" passed on from each parent. The ratios of tall and dwarf plants in the offspring showed that the unit for tall plants, <u>I</u>, was <u>dominant</u> over the unit for dwarf plants, <u>t</u>.

Mendel Reached Three Important Conclusions

Mendel reached these three important conclusions about heredity in plants:

- 1) Characteristics in plants are determined by "hereditary units".
- 2) Hereditary units are passed on to offspring unchanged from both parents, one unit from each parent.
- 3) Hereditary units can be dominant or recessive if an individual has both the dominant and the recessive unit for a characteristic, the dominant characteristic will be expressed.

It Took a While For People to Understand His Work

- Mendel's work was <u>cutting edge</u> and <u>new</u> to the scientists of the day. They didn't have the background knowledge to <u>properly understand</u> his findings — they had <u>no idea</u> about <u>genes</u>, <u>DNA</u> and <u>chromosomes</u>.
- 2) It wasn't until after his death that people realised how significant his work was.
- 3) Using Mendel's work as a starting point, the observations of many different scientists have contributed to the understanding of genes that we have today. For example:
 - In the <u>late 1800s</u>, scientists became familiar with <u>chromosomes</u>.
 They were able to observe how they <u>behaved</u> during <u>cell division</u>.
 - 2) Then in the <u>early 20th century</u>, scientists realised that there were striking <u>similarities</u> in the way that <u>chromosomes</u> and <u>Mendel's "units"</u> acted. Based on this, it was proposed that the "units" were <u>found</u> on the <u>chromosomes</u>. We now know these "units" as <u>genes</u>.
 - In 1953, the structure of DNA was determined (see page 85).
 This allowed scientists to go on and find out exactly how genes work.

Clearly, being a monk in the 1800s was a right laugh...

There was no TV in those days, you see, so Mendel filled his time by growing lots and lots of pea plants.

1 Suggest why the importance of Mendel's work wasn't realised straight away. [1 mark]

Variation

You'll probably have noticed that not all people are identical. There are reasons for this.

Organisms of the Same Species Have Differences

- 1) Different species look... well... different my dog definitely doesn't look like a daisy.
- But even organisms of the same species will usually look at least slightly different —
 e.g. in a room full of people you'll see different colour hair, individually shaped noses, etc.
- These differences are called the <u>variation</u> within a species, and there are <u>two</u> types of variation — <u>genetic variation</u> and <u>environmental variation</u>.



Different Genes Cause Genetic Variation

- 1) All plants and animals have characteristics that are in some ways similar to their parents'.
- 2) This is because an organism's <u>characteristics</u> are determined by the <u>genes inherited</u> from their <u>parents</u>. (Genes are the <u>codes</u> inside your cells that <u>control</u> how you're made — more about these on page 84).
- 3) These genes are passed on in sex cells (gametes), from which the offspring develop (see page 87).
- 4) Most animals (and quite a lot of plants) get some genes from the mother and some from the father.
- This combining of genes from two parents causes <u>genetic variation</u> no two of the species are <u>genetically identical</u> (other than identical twins).
- 6) Some characteristics are determined only by genes (e.g. violet flower colour). In animals these include: eye colour, blood group and inherited disorders (e.g. haemophilia or cystic fibrosis).

Characteristics are also Influenced by the Environment

- The environment, including the conditions that organisms live and grow in, also causes differences
 between members of the same species this is called environmental variation.
- 2) For example, a plant grown in plenty of <u>sunlight</u> would be <u>luscious</u> and <u>green</u>, but the <u>same</u> plant grown in darkness would grow tall and spindly and have yellow leaves. These are environmental variations.
- Environmental variation covers a <u>wide range</u> of differences from <u>losing your toes</u> in a piranha attack, to getting a <u>suntan</u>, to having <u>yellow leaves</u> (never happened to me yet though), and so on.

Most Characteristics are Due to Genes AND the Environment

- Most characteristics (e.g. body weight, height, skin colour, condition of teeth, academic or athletic prowess, etc.) are determined by a mixture of genetic and environmental factors.
- For example, the <u>maximum height</u> that an animal or plant could grow to is determined by its <u>genes</u>.
 But whether it actually grows that tall depends on its <u>environment</u> (e.g. how much food it gets).

Mutations Introduce Variation



- 1) Mutations are changes to the sequence of bases in <u>DNA</u>. As you saw on page 86, mutations can lead to changes in the <u>protein</u> that a gene codes for. <u>Most mutations</u> have <u>no effect</u> on the <u>protein</u> the gene codes for, so <u>most mutations</u> have <u>no effect</u> on the organism's <u>phenotype</u>. Some have a <u>small influence</u> on phenotype and so only alter characteristics <u>slightly</u>. However, although it's very rare, mutations can result in a <u>new phenotype</u> being seen in a species.
- If the <u>environment changes</u>, and the new phenotype makes an <u>individual more suited</u> to the new environment, it can become common <u>throughout</u> the species <u>relatively quickly</u> by natural selection — see the next page.

My mum's got no trousers — cos I've got her jeans...

So you can't blame all of your faults on your parents — the environment usually plays a role too.

O1 Explain what is meant by environmental variation.

[2 marks]

Evolution

THEORY OF EVOLUTION: All of today's species have evolved from simple life forms that first started to develop over three billion years ago.

1

Charles Darwin

Only the Fittest Survive

<u>Charles Darwin</u> came up with a really important theory about <u>evolution</u>. He used the <u>observations</u> he made on a huge <u>round-the-world trip</u>, along with <u>experiments</u>, <u>discussions</u> and new <u>knowledge</u> of <u>fossils</u> and <u>geology</u>, to suggest the theory of <u>evolution by natural selection</u>.

- Darwin knew that organisms in a species show wide variation in their characteristics (phenotypic variation). He also knew that organisms have to compete for limited resources in an ecosystem.
- 2) Darwin concluded that the organisms with the most <u>suitable characteristics</u> for the <u>environment</u> would be <u>more successful competitors</u> and would be <u>more likely to survive</u>.

 This idea is called the 'survival of the fittest'.
- The successful organisms that <u>survive</u> are more likely to <u>reproduce</u> and <u>pass on</u> the genes for the characteristics that made them successful to their offspring.
- 4) The organisms that are <u>less well adapted</u> would be <u>less likely to survive</u> and <u>reproduce</u>, so they are less likely to pass on their genes to the next generation.
- Over time, <u>beneficial characteristics</u> become <u>more common</u> in the population and the species changes — it evolves.

New Discoveries Have Helped to Develop the Theory

- Darwin's theory wasn't perfect. Because the relevant scientific knowledge <u>wasn't available</u> at the time, he couldn't give a good <u>explanation</u> for <u>why</u> new characteristics <u>appeared</u> or exactly <u>how</u> individual organisms <u>passed on</u> beneficial adaptations to their offspring.
- 2) We now know that <u>phenotype</u> is controlled by <u>genes</u>. New <u>phenotypic variations</u> arise because of <u>genetic variants</u> produced by <u>mutations</u> (changes in DNA see page 86). <u>Beneficial variations</u> are passed on to <u>future generations</u> in the <u>genes</u> that parents contribute to their offspring.

The Development of a New Species is Called Speciation

Over a long period of time, the phenotype of organisms can change so much because
of natural selection that a completely new species is formed. This is called speciation.

There's more = on speciation = on p.102.

 Speciation happens when populations of the same species change enough to become reproductively isolated — this means that they can't interbreed to produce fertile offspring.

Extinction is When No Individuals of a Species Remain

The fossil record contains many species that don't exist any more — these species are said to be extinct.

Species become extinct for these reasons:

- 1) The environment changes too quickly (e.g. destruction of habitat).
- 2) A new predator kills them all (e.g. humans hunting them).
- 3) A new disease kills them all.
- 4) They can't compete with another (new) species for food.
- A <u>catastrophic event</u> happens that kills them all (e.g. a volcanic eruption or a collision with an asteroid).

Dodos are now extinct. Humans not only hunted them, but introduced other animals which ate all their eggs, and we destroyed the forest where they lived — they really didn't stand a chance...

"Natural selection" — sounds like vegan chocolates...

Natural selection's all about the organisms with the best characteristics surviving to pass on their genes.

Q1 The sugary nectar in some orchid flowers is found at the end of a long tube behind the flower.

There are moth species with long tongues that can reach the nectar.

Explain how natural selection could have led to the moths developing long tongues. [4 marks]



More About Evolution

There's a lot of <u>evidence</u> for the theory of evolution by natural selection.

But back in the day, poor Charlie Darwin didn't have half as much evidence to convince people.

Not Everyone Agreed with Darwin...

When Darwin proposed his theory in his book "On the Origin of Species" in 1859, his idea was very controversial for various reasons...

It went against common religious beliefs about how life on Earth developed — it was the first plausible explanation for the existence of life on earth without the need for a "Creator" (God).

Darwin couldn't explain why these new, useful characteristics <u>appeared</u> or <u>how</u> they were <u>passed on</u> from individual organisms to their offspring. But then he didn't know anything about <u>genes</u> or <u>mutations</u> — they weren't discovered 'til 50 years after his theory was published.

There wasn't enough evidence to convince many <u>scientists</u>, because not many <u>other studies</u> had been done into how organisms change over time.

...and Lamarck had Different Ideas

There were different scientific hypotheses about evolution around at the same time, such as Lamarck's:

- Jean-Baptiste Lamarck (1744-1829) argued that changes that an organism acquires during
 its lifetime will be passed on to its offspring e.g. he thought that if a characteristic
 was used a lot by an organism, then it would become more developed during its
 lifetime, and the organism's offspring would inherit the acquired characteristic.
- For example, using this theory, if a rabbit <u>used</u> its legs to run a lot (to escape predators), then
 its legs would get <u>longer</u>. The offspring of that rabbit would then be <u>born</u> with <u>longer legs</u>.

Scientists Develop Different Hypotheses from Observations

- 1) Often scientists come up with different hypotheses to explain similar observations.
- Scientists might develop different hypotheses because they have different beliefs (e.g. religious) or they have been influenced by different people (e.g. other scientists and their way of thinking)... or they just darn well think differently.
- There's more about making hypotheses on page 1.
- 3) The only way to <u>find out</u> whose hypothesis is right is to find evidence to <u>support</u> or <u>disprove</u> each one.
- 4) For example, Lamarck and Darwin both had different hypotheses to explain how evolution happens. In the end...
 - Lamarck's hypothesis was eventually <u>rejected</u> because experiments <u>didn't support</u>
 <u>his hypothesis</u>. You can see it for yourself, e.g. if you dye a hamster's fur <u>bright pink</u>
 (not recommended), its offspring will still be born with the <u>normal</u> fur colour because
 the new characteristic won't have been passed on.
 - The discovery of genetics <u>supported</u> Darwin's idea because it provided an <u>explanation</u> of how organisms born with beneficial characteristics can <u>pass them on</u> (i.e. via their genes). Other evidence was also found by looking at <u>fossils</u> of <u>different ages</u> (the <u>fossil record</u>) this allows you to see how <u>changes</u> in organisms <u>developed slowly over time</u>. The relatively recent discovery of how <u>bacteria</u> are able to evolve to become <u>resistant to antibiotics</u> also further supports evolution by <u>natural selection</u>.
- 5) There's so much evidence for Darwin's idea that it's now an accepted hypothesis (a theory).

Here's to crazy new ideas...

Science is fuelled by new ideas. Without them, we'd still think the world is flat and that mullets are cool.

Q1 Explain why Darwin's theory was considered controversial when it was first proposed. [3 marks]

Selective Breeding

'<u>Selective breeding</u>' sounds like it has the potential to be a tricky topic, but it's actually dead simple.

You take the <u>best</u> plants or animals and breed them together to get the best possible <u>offspring</u>. That's it.

Selective Breeding is Very Simple

Selective breeding is when humans <u>artificially select</u> the plants or animals that are going to <u>breed</u> so that the genes for particular characteristics <u>remain</u> in the population.

Organisms are <u>selectively bred</u> to develop features that are <u>useful</u> or <u>attractive</u>, for example:

- · Animals that produce more meat or milk.
- · Crops with disease resistance.
- Dogs with a good, gentle temperament.
- Decorative plants with big or unusual flowers.

This is the basic process involved in selective breeding:

- 1) From your existing stock, select the ones which have the characteristics you're after.
- 2) Breed them with each other.
- 3) Select the best of the offspring, and breed them together.
- Continue this process over <u>several generations</u>, and the desirable trait gets <u>stronger</u> and <u>stronger</u>. Eventually, <u>all</u> the offspring will have the characteristic.


Selective breeding
is also known as
artificial selection'.

In <u>agriculture</u> (farming), selective breeding can be used to <u>improve yields</u>. E.g. to improve <u>meat yields</u>, a farmer could breed together the <u>cows</u> and <u>bulls</u> with the <u>best characteristics</u> for producing <u>meat</u>, e.g. large size. After doing this for <u>several generations</u> the farmer would get cows with a <u>very high meat yield</u>.

5) Selective breeding is <u>nothing new</u> — people have been doing it for <u>thousands</u> of years. It's how we ended up with <u>edible crops</u> from <u>wild plants</u> and how we got <u>domesticated</u> animals like cows and dogs.

The Main Drawback is a Reduction in the Gene Pool

- The main problem with selective breeding is that it reduces the gene pool the <u>number of different alleles</u> (forms of a gene) in a population. This is because the farmer keeps breeding from the "<u>best</u>" animals or plants which are all <u>closely related</u>. This is known as <u>inbreeding</u>.
- 2) Inbreeding can cause <u>health problems</u> because there's more chance of the organisms inheriting <u>harmful genetic defects</u> when the <u>gene pool</u> is <u>limited</u>. Some <u>dog breeds</u> are particularly susceptible to <u>certain defects</u> because of inbreeding e.g. pugs often have breathing problems.
- 3) There can also be serious problems if a new disease appears, because there's not much variation in the population. All the stock are closely related to each other, so if one of them is going to be killed by a new disease, the others are also likely to succumb to it.



I use the same genes all the time too — they flatter my hips...

Different breeds of dog came from selective breeding. For example, somebody thought 'I really like this small, yappy wolf — I'll breed it with this other one'. After thousands of generations, we got poodles.



Explain how you could selectively breed for floppy ears in rabbits.

O2 What potential issues can selective breeding cause? [3 marks]

01


Genetic Engineering

Genetic engineering is an interesting area of science with exciting possibilities, but there might be dangers too...

Genetic Engineering Transfers Genes Between Organisms

The basic idea of genetic engineering is to <u>transfer</u> a <u>gene</u> responsible for a <u>desirable characteristic</u> from one organism's genome into <u>another</u> organism, so that it also has the <u>desired characteristic</u>.

- 1) A useful gene is isolated (cut) from one organism's genome using enzymes and is inserted into a vector.
- The vector is usually a <u>virus</u> or a <u>bacterial plasmid</u> (a fancy piece of circular DNA found in bacterial cells), depending on the type of organism that the gene is being transferred to.
- 3) When the vector is introduced to the target organism, the useful gene is inserted into its cell(s).
- 4) Scientists use this method to do all sorts of things. For example:
 - Bacteria have been genetically modified to produce human insulin that can be used to treat diabetes.
 - Genetically modified (GM) crops have had their genes modified, e.g. to improve the size and quality of their fruit, or make them resistant to disease, insects and herbicides (chemicals used to kill weeds).
 - Sheep have been genetically engineered to produce substances, like drugs, in their milk that can be used to treat human diseases.



5) In some cases, the transfer of the gene is carried out when the organism receiving the gene is at an <u>early stage</u> of <u>development</u> (e.g. egg or embryo). This means that the organism <u>develops</u> with the <u>characteristic</u> coded for by the gene.

Genetic Engineering is a Controversial Topic

- Genetic engineering is an exciting area of science, which has the potential for solving many
 of our problems (e.g. treating diseases, more efficient food production etc.), but not everyone
 thinks it's a great idea.
- There are <u>worries</u> about the long-term effects of genetic engineering that changing an organism's
 genes might <u>accidentallu</u> create unplanned <u>problems</u>, which could get passed on to <u>future generations</u>.

There Are Pros and Cons of GM Crops

- Some people say that growing GM crops will affect the number of wild flowers (and so the
 population of insects) that live in and around the crops reducing farmland biodiversity.
- 2) Not everyone is convinced that GM crops are <u>safe</u> and some people are concerned that we might not <u>fully understand</u> the effects of eating them on <u>human health</u>. E.g. people are worried they may develop <u>allergies</u> to the food although there's probably no more risk for this than for eating usual foods.
- 3) A big concern is that <u>transplanted genes</u> may get out into the <u>natural environment</u>. For example, the <u>herbicide resistance</u> gene may be picked up by weeds, creating a new <u>'superweed'</u> variety.
- 4) On the plus side, the characteristics chosen for GM crops can increase the yield, making more food.
- 5) People living in developing nations often lack <u>nutrients</u> in their diets. GM crops could be <u>engineered</u> to contain the nutrient that's <u>missing</u>. For example, 'golden rice' is a GM rice crop that contains beta-carotene lack of this substance causes <u>blindness</u>.
- 6) GM crops are already being grown in some places, often without any problems.

If only there was a gene to make revision easier...

Make sure you've got everything on this page firmly in your noggin. You need to understand the lot.

Outline one benefit and one concern about GM crops.

[2 marks]

Cloning

We can clone plants and animals in several different ways. Cool. But some of them have potential problems...

Plants Can Be Cloned by Tissue Culture and from Cuttings

This is where a few plant cells are put in a growth medium with hormones, and they grow into new plants — clones of the parent plant. These plants can be made very quickly, in very little space, and be grown all year. Tissue culture is used by scientists to preserve rare plants that are hard to reproduce naturally and by plant nurseries to produce lots of stock quickly.

Gardeners can take cuttings from good parent plants, and then plant them to produce genetically identical copies (clones) of the parent plant.





2) These plants can be produced quickly

and cheaply. This is an older, simpler method than tissue culture.

You Can Make Animal Clones Using Embryo Transplants

Farmers can produce cloned offspring from their best bull and cow — using embruo transplants.

- Sperm cells are taken from a prize bull and egg cells are taken from a prize cow. The sperm are then used to artificially fertilise an egg cell. The embryo that develops is then split many times (to form clones) before any cells become specialised.
- 2) These cloned embruos can then be implanted into lots of other cows where they grow into baby calves (which will all be genetically identical to each other).
- 3) Hundreds of "ideal" offspring can be produced every year from the best bull and cow.

Adult Cell Cloning is Another Way to Make a Clone

- Adult cell cloning involves taking an unfertilised egg cell and removing its nucleus. The nucleus is then removed from an adult body cell (e.g. skin cell) and is inserted into the 'empty' egg cell.
- 2) The egg cell is then stimulated by an electric shock this makes it divide, just like a normal embryo.
- 3) When the embryo is a ball of cells, it's implanted into the womb of an adult female. It grows into a genetically identical copy (clone) of the original adult body cell as it has the same genetic information.
- 4) This technique was used to create Dolly the famous cloned sheep.

There are Many Issues Surrounding Cloning

- Cloning quickly gets you lots of "ideal" offspring. But you also get a "reduced gene pool" this means there are fewer different alleles in a population. If a population are all closely related and a new disease appears, they could all be wiped out — there may be no allele in the population giving resistance to it.
- 2) But the study of animal clones could lead to greater understanding of the development of the embruo, and of ageing and age-related disorders.
- 3) Cloning could also be used to help preserve endangered species.
- 4) However, it's possible that cloned animals might not be as healthy as normal ones, e.g. Dolly the sheep had arthritis, which tends to occur in older sheep (but the jury's still out on if this was due to cloning).
- 5) Some people worry that humans might be cloned in the future. If it was allowed, any success may follow many unsuccessful attempts, e.g. children born severely disabled.

Thank goodness they didn't do that with my little brother...

Cloning can be controversial — especially when it's to do with cloning animals (and especially humans).

Describe how animals can be cloned by adult cell cloning.

[5 marks]

Fossils

Fossils are great. If they're <u>well-preserved</u>, you can see what oldy-worldy creatures <u>looked</u> like.

They also show how living things have <u>evolved</u>. Although we're not sure how life started in the first place...

Fossils are the Remains of Plants and Animals

Fossils are the <u>remains</u> of organisms from <u>many thousands of years ago</u>, which are found in <u>rocks</u>. They provide the <u>evidence</u> that organisms lived ages ago. Fossils can tell us a lot about <u>how much</u> or <u>how little</u> organisms have <u>changed</u> (<u>evolved</u>) over time. Fossils form in rocks in one of <u>three</u> ways:

1) FROM GRADUAL REPLACEMENT BY MINERALS (Most fossils happen this way.)

- 1) Things like teeth, shells, bones etc., which don't decay easily, can last a long time when buried.
- They're eventually <u>replaced by minerals</u> as they decay, forming a <u>rock-like substance</u> shaped like the original hard part.
- The surrounding sediments also turn to rock, but the fossil stays distinct inside the rock and eventually someone digs it up.



2) FROM CASTS AND IMPRESSIONS

- Sometimes, fossils are formed when an organism is <u>buried</u> in a <u>soft</u> material like clay.
 The clay later <u>hardens</u> around it and the organism decays, leaving a <u>cast</u> of itself.
 An animal's <u>burrow</u> or a <u>plant's roots</u> (rootlet traces) can be preserved as casts.
- Things like footprints can also be <u>pressed</u> into these materials when soft, leaving an <u>impression</u> when it hardens.

3) FROM PRESERVATION IN PLACES WHERE NO DECAY HAPPENS

- In amber (a clear yellow 'stone' made from fossilised resin) and tar pits there's no oxygen or moisture so decay microbes can't survive.
- 2) In glaciers it's too cold for the decay microbes to work.
- Peat bogs are too acidic for decay microbes.
 (A fully preserved man they named 'Pete Marsh' was found in a bog.)



But No One Knows How Life Began

Fossils show how much or how little different organisms have changed (evolved) as life has developed on Earth over millions of years. But where did the first living thing come from...

- 1) There are various hupotheses suggesting how life first came into being, but no one really knows.
- 2) Maybe the first life forms came into existence in a primordial <u>swamp</u> (or under the <u>sea</u>) here on <u>Earth</u>. Maybe simple organic molecules were brought to Earth on <u>comets</u> these could have then become more <u>complex</u> organic molecules, and eventually very simple <u>life forms</u>.
- 3) These hypotheses can't be supported or disproved because there's a lack of good, valid evidence:
 - Many early forms of life were <u>soft-bodied</u>, and soft tissue tends to decay away <u>completely</u> — so the fossil record is <u>incomplete</u>.
 - Fossils that did form millions of years ago may have been destroyed by geological activity, e.g. the movement of tectonic plates may have crushed fossils already formed in the rock.

Don't get bogged down by all this information...

It's a bit mind-boggling really how fossils can still exist even millions of years after the organism died. They really are fascinating things, and scientists have learned a whole lot from studying them in detail.

O1 Suggest what makes low-oxygen environments suitable for the formation of fossils.

[2 marks]

Speciation

Evolution (see page 96) leads to the development of lots of different species. This process is called speciation.

Speciation is the Development of a New Species

- 1) A species is a group of similar organisms that can reproduce to give fertile offspring.
- 2) Speciation is the development of a new species.
- Speciation occurs when <u>populations</u> of the <u>same species</u> become so <u>different</u> that they can <u>no longer successfully interbreed</u> to produce <u>fertile offspring</u>.




Isolation and Natural Selection Lead to Speciation

Isolation is where populations of a species are separated. This can happen due to a physical barrier.

E.g. floods and earthquakes can cause barriers that geographically isolate some individuals from the main population. Conditions on either side of the barrier will be slightly different, e.g. they may have different climates. Because the environment is different on each side, different characteristics will become more common in each population due to natural selection operating differently on the populations.

- 1) Each population shows genetic variation because they have a wide range of alleles.
- In each population, individuals with characteristics that make them better adapted to their environment have a <u>better chance of survival</u> and so are more likely to <u>breed</u> successfully.
- So the <u>alleles</u> that control the <u>beneficial characteristics</u> are more likely to be <u>passed on</u> to the <u>next generation</u>.

Eventually, individuals from the different populations will have changed so much that they won" to able to breed with one another to produce fertile offspring. The two groups will have become separate species.



Wallace Was a Pioneer of the Theory of Speciation

Alfred Russel Wallace was a scientist working at the same time as Charles Darwin.
He was one of the early scientists working on the idea of speciation. His observations greatly contributed to how we understand speciation today. Our current understanding developed as more evidence became available over time.

There's more
on Darwin and
natural selection
on page 96.

- 2) During his career, Wallace independently came up with the idea of natural selection and published work on the subject together with Darwin in 1858. This then prompted Darwin to publish 'On the Origin of Species' in 1859.
- 3) Observations made by Wallace as he travelled the world provided lots of evidence to support the theory of evolution by natural selection. For example, he realised that warning colours are used by some species (e.g. butterflies) to deter predators from eating them and that this was an example of a beneficial characteristic that had evolved by natural selection.
- 4) It's this work on warning colours and his work on speciation that he's most famous for.

If only Darwin's name had been Gromit...

Well isn't speciation riveting... Put some time into learning this page, then maybe grab a biscuit.

O1 Why might natural selection work differently on two isolated populations of a species?

[3 marks]

more common in the

population because of

page 96 for more.

= natural selection — see

Antibiotics don't kill

viruses — see p.51.

Antibiotic-Resistant Bacteria

The discovery of <u>antibiotics</u>, like <u>penicillin</u>, was a huge benefit to medicine — suddenly bacterial infections that had often been fatal could be cured. But unfortunately they might not be a permanent solution.

Bacteria can Evolve and Become Antibiotic-Resistant

- 1) Like all organisms, bacteria sometimes develop random mutations (see p.86) in their DNA.

 These can lead to changes in the bacteria's characteristics, e.g. being less affected by a particular antibiotic. This can lead to antibiotic-resistant strains forming as the gene for antibiotic resistance becomes more common in the population.
- To make matters worse, because bacteria are so rapid at reproducing, they can evolve quite quickly.
- 3) For the bacterium, the ability to resist antibiotics is a big advantage. It's better able to survive, even in a host who's being treated to get rid of the infection, and so it lives for longer and reproduces many more times. This increases the population size of the antibiotic-resistant strain.
- Antibiotic-resistant strains are a problem for people who become <u>infected</u> with these bacteria because
 they aren't immune to the new strain and there is no effective treatment. This means that the infection
 <u>easily spreads</u> between people. Sometimes drug companies can come up with a <u>new</u> antibiotic that's
 effective, but 'superbugg' that are resistant to most known antibiotics are becoming more common.
- MRSA is a relatively common 'superbug' that's really hard to get rid of.
 It often affects people in hospitals and can be fatal if it enters their bloodstream.

Antibiotic Resistance is Becoming More Common

- For the last few decades, we've been able to deal with <u>bacterial infections</u> pretty easily using <u>antibiotics</u>. The <u>death rate</u> from infectious bacterial diseases (e.g. pneumonia) has <u>fallen</u> dramatically.
- But the problem of <u>antibiotic resistance</u> is getting <u>worse</u> partly because
 of the <u>overuse</u> and <u>inappropriate use</u> of antibiotics, e.g. doctors prescribing
 them for <u>non-serious conditions</u> or infections caused by <u>viruses</u>.

3) The more often antibiotics are used, the bigger the problem of antibiotic resistance becomes, so it's important that doctors only prescribe antibiotics when they really need to:

It's not that antibiotics actually <u>cause</u> resistance — they create a situation where naturally resistant bacteria have an advantage and so increase in numbers.

- 4) It's also important that you take all the antibiotics a doctor prescribes for you:
 - Taking the <u>full course</u> makes sure that <u>all</u> the bacteria are <u>destroyed</u>, which means that there are <u>none</u> left to mutate and develop into <u>antibiotic-resistant strains</u>.
- 5) In <u>farming</u>, antibiotics can be given to animals to <u>prevent</u> them becoming <u>ill</u> and to make them <u>grow faster</u>. This can lead to the development of <u>antibiotic-resistant bacteria</u> in the animals which can then <u>spread to humans</u>, e.g. during meat preparation and consumption. Increasing concern about the <u>overuse</u> of antibiotics in agriculture has led to some countries <u>restricting their use</u>.
- 6) The increase in antibiotic resistance has encouraged drug companies to work on developing <u>new</u> antibiotics that are <u>effective</u> against the resistant strains. Unfortunately, the <u>rate of development</u> is <u>slow</u>, which means we're <u>unlikely</u> to be able to keep up with the <u>demand</u> for new drugs as <u>more</u> antibiotic-resistant strains develop and spread. It's also a very <u>costly</u> process.

Aaargh, a giant earwig! Run from the attack of the superbug...

The reality of 'superbugs' is even scarier than giant earwigs. Microorganisms that are resistant to all our drugs are a worrying thought. It'll be like going back in time to before antibiotics were invented.

- a worrying thought. It if he like going back in time to before antibiotics were invented.

 Q1 Suggest a situation where antibiotics could be prescribed inappropriately. [1 mark]
- Q2 Explain why it's important that people take the full course of antibiotics they are prescribed. [2 marks]

Classification

It seems to be a basic human urge to want to classify things — that's the case in biology anyway...

Classification is Organising Living Organisms into Groups

- Traditionally, organisms have been <u>classified</u> according to a system first proposed in the 1700's by <u>Carl Linnaeus</u>, which <u>groups</u> living things according to their <u>characteristics</u> and the <u>structures</u> that make them up.
- In this system (known as the <u>Linnaean system</u>), living things are first divided into <u>kingdoms</u> (e.g. the plant kingdom).
- The kingdoms are then <u>subdivided</u> into smaller and smaller groups — <u>phylum</u>, <u>class</u>, <u>order</u>, <u>family</u>, <u>genus</u>, <u>species</u>.

Kingdom Phylum Class Order Family Genus Species

Classification Systems Change Over Time

- As knowledge of the <u>biochemical processes</u> taking place inside organisms developed and <u>microscopes</u> <u>improved</u> (which allowed us to find out more about the <u>internal structures</u> of organisms), scientists put forward <u>new</u> models of classification.
- 2) In 1990, Carl Woese proposed the <a href="https://doi.org/10.1009/jhttps://doi.org/10.1009
- 3) In the three-domain system, organisms are first of all split into three large groups called domains:
 - ARCHAEA Organisms in this domain were once thought to be <u>primitive bacteria</u>, but they're actually a different type of <u>prokaryotic cell</u>. They were first found in <u>extreme places</u> such as hot springs and salt lakes.
 - BACTERIA This domain contains true bacteria like E. coli and Staphylococcus. Although they
 often look similar to Archaea, there are lots of biochemical differences between them.
 - EUKARYOTA This domain includes a <u>broad range</u> of organisms including <u>fungi</u> (page 46), <u>plants</u>, <u>animals</u> and <u>protists</u> (page 46).
- 4) These are then subdivided into smaller groups kingdom, phylum, class, order, family, genus, species.

Organisms Are Named According to the Binomial System

- 1) In the binomial system, every organism is given its own two-part Latin name.
- 2) The <u>first</u> part refers to the <u>genus</u> that the organism belongs to. This gives you information on the organism's <u>ancestru</u>. The <u>second</u> part refers to the <u>species</u>.


E.g. humans are known as Homo sapiens. 'Homo' is the genus and 'sapiens' is the species.

> distant ancestor

3) The binomial system is used <u>worldwide</u> and means that scientists in <u>different countries</u> or who speak different languages all refer to a particular species by the same name — avoiding potential confusion.

Evolutionary Trees Show Evolutionary Relationships

- 1) Evolutionary trees show how scientists think different species are related to each other.
- 2) They show <u>common ancestors</u> and relationships between species. The more <u>recent</u> the common ancestor, the more <u>closely related</u> the two species — and the more <u>characteristics</u> they're likely to share.
- 3) Scientists analyse lots of different types of <u>data</u> to work out evolutionary relationships. For <u>living</u> organisms, they use the <u>current classification data</u> (e.g. DNA analysis and structural similarities). For <u>extinct</u> species, they use information from the <u>fossil record</u> (see page 96).



Whales and dolphins have a recent common ancestor so are closely related. They're both more distantly related to sharks.

Binomial system — uh oh, sounds like maths...

Sometimes, the genus in a binomial name is abbreviated to a capital letter with a full stop after it.

Q1 The evolutionary tree on the right shows the relationship between four species, A-D.

Which two species shown in the tree are the most closely related? [1 mark] A B C D



Revision Questions for Topic 6

You've finished Topic 6 — Hoorah. Now have a go at these questions...

- Try these questions and tick off each one when you get it right.
- When you're completely happy with a sub-topic, tick it off.





DNA, Genes and Mutations (p.84-86) 1) What do genes code for? 2) List the parts of a DNA nucleotide. 3) How many bases in a DNA sequence code for one amino acid? 4) Give a function of the non-coding parts of DNA. 5) Explain how mutations can lead to a change in the protein coded for by a gene. 6) True or False: "Most mutations have little or no effect on the protein coded for bu a gene". Reproduction and Meiosis (p.87-89) 7) Name the male and female gametes of animals. 8) Which tupe of reproduction produces genetically identical cells? 9) State the tupe of cell division used to make gametes in humans. 10) How does variation in a population increase its chance of surviving environmental change? 11) Describe the methods of sexual and asexual reproduction used by strawberry plants. Sex Chromosomes, Genetic Diagrams and Inherited Disorders (p.90-93) 12) What is the probability that offspring will have the XX combination of sex chromosomes? 13) How many genes are responsible for controlling fur colour in mice? 14) What are alleles? 15) What does it mean if someone is heterozugous for a gene? 16) What is the chance of a child being born with poludactulu if one parent has a single dominant allele for the gene that controls it? 17) Give two arguments for and two arguments against screening embryos for genetic disorders. Mendel, Variation and Evolution (p.94-97) 18) What do we now know Mendel's "units" as? 19) What is variation? 20) Explain how beneficial characteristics can become more common in a population over time. 21) What was Jean-Baptiste Lamarck's theory about evolution? Selective Breeding, Genetic Engineering and Cloning (p.98-100) 22) What is selective breeding? 23) How might farmers use selective breeding? 24) What is genetic engineering? 25) How can embryo transplants be used to create animal clones? Fossils, Speciation and Antibiotic-Resistant Bacteria (p.101-103) 26) Give two ways that fossils can be formed. 27) What is speciation and when can it occur? 28) What leads to the formation of antibiotic-resistant strains of bacteria? Classification (p.104)

29) Name the groups that organisms are classified into in the Linnaean system.
30) Who proposed the 'three-domain system' of classification in 1990?

Competition

Ecology is all about organisms and the environment they live in, and how the two interact. Simples.

First Learn Some Words to Help You Understand Ecology...

This topic will make a lot more sense if you become familiar with these terms first:

- 1) Habitat the place where an organism lives.
- 2) Population all the organisms of one species living in a habitat.
- 3) Community the populations of different species living in a habitat.
- 4) Abiotic factors non-living factors of the environment, e.g. temperature.
- 5) Biotic factors living factors of the environment, e.g. food.
- Ecosystem the interaction of a community of living organisms (biotic) with the non-living (abiotic) parts of their environment.



Organisms Compete for Resources to Survive

Organisms need things from their environment and from other organisms in order to survive and reproduce:

- 1) Plants need light and space, as well as water and mineral ions (nutrients) from the soil.
- 2) Animals need space (territory), food, water and mates.

Organisms compete with other species (and members of their own species) for the same resources.


Any Change in Any Environment can Have Knock-on Effects

In a community, each species <u>depends</u> on other species for things such as <u>food</u>, <u>shelter</u>, <u>pollination</u> and <u>seed dispersal</u> — this is called interdependence.

The <u>interdependence</u> of all the living things in an ecosystem means that any major <u>change</u> in the ecosystem (such as one species being removed) can have <u>far-reaching effects</u>.

The diagram on the right shows part of a <u>food web</u> (a diagram of what eats what) from a stream.

Stonefly larvae are particularly sensitive to pollution. Suppose pollution killed them in this stream. The table below shows some of the effects this might have on some of the other organisms in the food web.



	-		MINIMUM AND A
Organism	Effect of loss of stonefly larvae	Fffect on population	Remember that 1000 =
Blackfly	Less competition for algae	Increase	webs are very complex =
larvae	More likely to be eaten by predators	Decrease	- are difficult to =
Water spider	Less food	Decrease	predict accurately.
Stickleback	Less food (if water spider or mayfly larvae numbers decrease)	Decrease	1 ////////

In some communities, all the species and environmental factors are in <u>balance</u> so that the <u>population sizes</u> are <u>roughly constant</u> (they may go up and down in cycles — see p.109). These are called <u>stable communities</u>. Stable communities include <u>tropical rainforests</u> and <u>ancient oak woodlands</u>.

I'm dependent on the cocoa tree...

If my source of chocolate was removed, it would have far-reaching effects on my revision and grades. Seriously though, make sure you know what organisms compete for in an ecosystem. Then try these...



[3 marks]

Q2 Using the food web above, suggest what might happen to the frog population if the stickleback population decreased.

[2 marks]



Abiotic and Biotic Factors

The environment in which organisms live changes all the time. The things that change are either abiotic (non-living) or biotic (living) factors. These can have a big effect on a community...

Abiotic Factors Can Vary in an Ecosystem...

Abiotic factors are non-living factors. For example:

- 1) Moisture level
- 2) Light intensity
- 3) Temperature
- 4) Carbon dioxide level (for plants)

- 5) Wind intensity and direction
- 6) Oxugen level (for aquatic animals)
- 7) Soil pH and mineral content

A change in the environment could be an increase or decrease in an abiotic factor, e.g. an increase in temperature. These changes can affect the size of populations in a community. This means they can also affect the population sizes of other organisms that depend on them (see previous page).

For example, a decrease in light intensity, temperature or level of carbon dioxide could decrease the rate of photosynthesis in a plant species (see p.57). This could affect plant growth and cause a decrease in the population size.

For example, a decrease in the mineral content of the soil (e.g. a lack of nitrates) could cause nutrient deficiencies (see p.55). This could also affect plant growth and cause a decrease in the population size.

2000 minimining Animals depend on plants for food, so a decrease in a plant population could affect = the animal species in a = community.

...and So Can Biotic Factors

Biotic factors are living factors. Here are some examples:


- 1) New predators arriving
- 2) Competition one species may outcompete another so that numbers are too low to breed
- New pathogens
- 4) Availability of food

A change in the environment could be the introduction of a new biotic factor, e.g. a new predator or pathogen. These changes can also affect the size of populations in a community, which can have knock-on effects because of interdependence (see previous page).

For example, a new predator could cause a decrease in the preu population. There's more about predator-prey populations on p.109.

For example, red and grey squirrels live in the same habitat and eat the same food. Grey squirrels outcompete the red squirrels - so the population of red squirrels is decreasing

The following graph shows the effect of a new pathogen on Species A. The population size of species A was increasing up until 1985, when it decreased rapidly until 1990 - suggesting that 1985 was the year that the new pathogen arrived. The population started to rise again after 1990.



Exams — a type of abiotic factor affecting my environment...

So, two lists of factors that would be a good idea to learn. I reckon this is a prime time for shutting the book, scribbling them all down and then checking how you did. Also, have a gander at that squirrel example.

- Give four examples of abiotic factors that could affect a plant species. 01 [4 marks]
- Q2 Cutthroat trout are present in lakes in Yellowstone National Park. In the last few decades, lake trout have been introduced to the lakes. However, lake trout have emerged as predators of the cutthroat trout. Give two other biotic factors that could affect the size of the cutthroat trout population. [2 marks]

Adaptations

Life exists in so many different environments because the organisms that live in them have adapted to them.

Adaptations Allow Organisms to Survive

Organisms, including microorganisms, are adapted to live in different environmental conditions. The features or characteristics that allow them to do this are called adaptations. Adaptations can be:

1) Structural

These are features of an organism's body structure — such as shape or colour. For example:

Arctic animals like the Arctic fox have white fur so they're camouflaged against the snow. This helps them avoid predators and sneak up on prey.

Animals that live in cold places (like whales) have a thick layer of blubber (fat) and a low surface area to volume ratio to help them retain heat.

Animals that live in hot places (like camels) have a thin layer of fat and a large surface area to volume ratio to help

them lose heat.

2) Behavioural

These are ways that organisms behave. Many species (e.g. swallows) migrate to warmer climates during the winter to avoid the problems of living in cold conditions.

3) Functional

These are things that go on inside an organism's body that can be related to processes like reproduction and metabolism (all the chemical reactions happening in the body). For example:

Desert animals conserve water by producing very little sweat and small amounts of concentrated urine.

Brown bears hibernate over winter. They lower their metabolism which conserves energy, so they don't have to hunt when there's not much food about.



Microorganisms Have a Huge Variety of Adaptations...

...so that they can live in a wide range of environments:

Some microorganisms (e.g. bacteria) are known as extremophiles — they're adapted to live in very extreme conditions. For example, some can live at high temperatures (e.g. in super hot volcanic vents), and others can live in places with a high salt concentration (e.g. very salty lakes) or at high pressure (e.g. deep sea vents).




In a nutshell, it's horses for courses...

In the exam, you might have to say how an organism is adapted to its environment. Look at its characteristics (e.g. colour/shape) as well as the conditions it has to cope with (e.g. predation/temperature) and you'll be sorted.

- 01 The diagram on the right shows a penguin. Penguins live in the cold, icy environment of the Antarctic. They swim in the sea to hunt for fish to eat. Some penguins also huddle together in large groups to keep warm.
 - a) What type of adaptation is being described when penguins 'huddle together'?
 - b) Explain one structural adaptation

[2 marks]







Food Chains

If you like food, and you like chains, then food chains might just blow your mind. Strap yourself in and prepare for some 'edge of your seat' learning, because the show is about to begin...

Food Chains Show What's Eaten by What in an Ecosystem

- 1) Food chains always start with a producer.
 - Producers make (produce) their own food using energy from the Sun.
- 2) Producers are usually green plants or algae they make glucose by photosynthesis (see page 57).
- 3) When a green plant produces glucose, some of it is used to make other biological molecules in the plant.
- 4) These biological molecules are the plant's biomass the mass of living material.
- 5) Biomass can be thought of as energy stored in a plant.
- 6) Energy is transferred through living organisms in an ecosystem when organisms eat other organisms.
- 7) Producers are eaten by primary consumers. Primary consumers are then eaten by secondary consumers and secondary consumers are eaten by tertiary consumers. Here's an example of a food chain:

Producers Primary consumers Secondary consumer 5000 dandelions... feed... 100 rabbits... which feed... 1 fox.

that eat other organisms. 'Primary' means 'first', so primary consumers are the first consumers in a food chain. Secondary consumers are second and tertiary consumers are third.


Populations of Prey and Predators Go in Cycles

Consumers that hunt and kill other animals are called predators, and their preu are what they eat. In a <u>stable community</u> containing <u>prey</u> and <u>predators</u> (as most of them do of course):

1) The population of any species is usually limited by the amount of food available.

For more about a stable community see page 106.

2) If the population of the <u>preu</u> increases, then so will the population of the <u>predators</u>. 3) However as the population of predators increases, the number of prey will decrease.



E.g. More grass means more rabbits More rabbits means more foxes. But more foxes means fewer rabbits.

Eventually fewer rabbits will mean fewer foxes again. This up and down pattern continues...

4) Predator-prey cycles are always out of phase with each other. This is because it takes a while for one population to respond to changes in the other population. E.g. when the number of rabbits goes up, the number of foxes doesn't increase immediately because it takes time for them to reproduce.

When the TV volume goes up... my revision goes down...

You might think that the start of a food chain always has to be a plant. In most cases it is, but sometimes organisms like algae can be too because they photosynthesise. No wonder algae always looks smug...



- Look at the following food chain for a particular area: grass \rightarrow grasshopper \rightarrow rat \rightarrow snake
 - a) Name the producer in the food chain. [1 mark]
 - b) How many consumers are there in the food chain? [1 mark]
 - Name the primary consumer in the food chain. [1 mark]
 - d) All the rats in the area are killed. Explain two effects that this could have on the food chain.

[4 marks]



Using Quadrats

This is where the <u>fun</u> starts. Studying <u>ecology</u> gives you the chance to <u>rummage around</u> in bushes, get your hands <u>dirty</u> and look at some <u>real organisms</u>, living in the <u>wild</u>. Hold on to your hats folks...

Organisms Live in Different Places Because The Environment Varies

- 1) As you know from page 106, a habitat is the place where an organism lives, e.g. a playing field.
- 2) The distribution of an organism is where an organism is found, e.g. in a part of the playing field.
- 3) Where an organism is found is affected by environmental factors (see page 107). An organism might be more common in one area than another due to differences in environmental factors between the two areas. For example, in the playing field, you might find that daisies are more common in the open than under trees, because there's more light available in the open.
- 4) There are a couple of ways to study the distribution of an organism. You can:
 - · measure how common an organism is in two sample areas (e.g. using quadrats) and compare them.
 - study how the distribution changes across an area, e.g. by placing quadrats along a <u>transect</u> (p.III).
 Both of these methods give quantitative data (numbers) about the distribution.

Use Quadrats to Study The Distribution of Small Organisms

A <u>quadrat</u> is a <u>square</u> frame enclosing a <u>known area</u>, e.g. 1 m². To compare <u>how common</u> an organism is in <u>two sample areas</u> (e.g. shady and sunny spots in that playing field) just follow these simple steps:

1) Place a Im² quadrat on the ground at a random point within the first sample area. A quadrat E.g. divide the area into a grid and use a random number generator to pick coordinates.



3) Repeat steps 1 and 2 as many times as you can.

4) Work out the mean number of organisms per quadrat within the first sample area.





Anna counted the number of daisies in 7 quadrats within her first sample area and recorded the following results: 18, 20, 22, 23, 23, 25

Here the MEAN is:
$$\frac{\text{TOTAL number of organisms}}{\text{NUMBER of quadrats}} = \frac{154}{7} = 22 \text{ daisies per quadrat}$$

- 5) Repeat steps 1 to 4 in the second sample area.
- Finally compare the two means. E.g. you might find 2 daisies per m² in the shade, and 22 daisies per m² (lots more) in the open field.

You Can Also Work Out the Population Size of an Organism in One Area

EXAMPLE

Students used quadrats, each with an area of 0.5 m², to randomly sample daisies on an open field. The students found a mean of 10.5 daisies per quadrat. The field had an area of 800 m². Estimate the population of daisies on the field.

1) Work out the <u>mean number of organisms per m²</u>. $1 \div 0.5 = 2$ $2 \times 10.5 = 21$ daisies per m²

Then multiply the mean by the total area (in m²)
of the habitat.

800 × 21 = 16 800 daisies on the open field The population size of an organism is sometimes called its abundance.

If your quadrat has an area of 1 m², the mean number of organisms per m² is just the same as the mean number per quadrat.

Drat, drat, and double drat - my favourite use of quadrats...

It's key that you make sure you put your quadrat down in a random place before you start counting.

Q1 A 1200 m² field was randomly sampled for buttercups using a quadrat with an area of 0.25 m². A mean of 0.75 buttercups were found per quadrat.

Estimate the total population of buttercups.

[2 marks]

Using Transects




So, now you think you've learnt all about distribution. Well hold on — there's more ecology fun to be had.

Use Transects to Study The Distribution of Organisms Along a Line

You can use lines called <u>transects</u> to help find out how organisms (like plants) are <u>distributed</u> across an area — e.g. if an organism becomes <u>more or less common</u> as you move from a hedge towards the middle of a field. Here's what to do:

- Mark out a line in the area you want to study using a tape measure.
 Then collect data along
- the line.

 3) You can do this by
- just <u>counting</u> all the organisms you're interested in that <u>touch</u> the line.




 Or, you can collect data by using <u>quadrats</u> (see previous page). These can be placed <u>next to</u> each other along the line or <u>at intervals</u>, for example, every 2 m.

You Can Estimate the Percentage Cover of a Quadrat

If it's difficult to count all the individual organisms in the quadrat (e.g. if they're grass) you can calculate the <u>percentage cover</u>. This means estimating the percentage <u>area</u> of the quadrat covered by a particular type of organism, e.g. by <u>counting</u> the number of little squares covered by the organisms.

EXAMPLE Sor sch

Some students were measuring the distribution of organisms from one corner of a school playing field to another, using quadrats placed at regular intervals along a transect. Below is a picture of one of the quadrats. Calculate the percentage cover of each organism, A and B.



You count a square if it's more than half covered.

- Count the <u>number of squares</u> covered by organism A.
- Make this into a <u>percentage</u> divide the number of squares covered by the organism by the total number of squares in the quadrat (100), then multiply the result by 100.
- 3) Do the same for organism B.

Type A = 42 squares

(42/100) × 100 = 0.42 × 100 = 42%

Type B = 47 squares $(47/100) \times 100$ = 0.47 × 100 = 47%

A slug that's been run over — definitely a widely-spread organism

So if you want to measure the distribution of a organism across an area, you could use a transect. You can either use them alone or along with quadrats. Now who's for a game of tennis... I've got my transect up.

- Q1 What is a transect? [1 mark]
- Q2 Some students want to measure how the distribution of dandelions changes across a field, from one corner to another. Describe a method they could use to do this.
- O3 How could you estimate the number of organisms in a quadrat, if they are difficult to count?

[1 mark]

[2 marks]

Environmental Change & The Water Cycle

Now you know how to measure the distribution of organisms (see pages 110-111), it's time to learn why their distribution may change over time. Then it's on to something completely different — the water cycle.

Environmental Changes Affect The Distribution of Organisms

<u>Environmental changes</u> can cause the <u>distribution of organisms</u> to change. A change in distribution means a change in <u>where</u> an organism <u>lives</u>. Environmental changes that can affect organisms in this way include:

1) A change in the AVAILABILITY of WATER. For example:

The <u>distribution</u> of some <u>animal</u> and <u>plant species</u> in the <u>tropics</u> changes between the <u>wet</u> and the <u>dry</u> seasons — i.e. the times of year where there is more or less <u>rainfall</u>, and so more or less <u>water available</u>. E.g. each year in Africa, large numbers of <u>glant</u> wildebeest migrate, moving north and then back south as the rainfall patterns change.

2) A change in the TEMPERATURE. For example:

The distribution of <u>bird species</u> in Germany is changing because of a rise in average <u>temperature</u>. E.g. the <u>European bee-eater</u> bird is a <u>Mediterranean</u> species but it's now present in parts of <u>Germany</u>.



3) A change in the composition of ATMOSPHERIC GASES. For example:


The distribution of some species <u>changes</u> in areas where there is more <u>air pollution</u>. E.g. some species of <u>lichen</u> can't grow in areas where <u>sulfur dioxide</u> is given out by certain <u>industrial processes</u>.

These environmental changes can be caused by <u>seasonal</u> factors, <u>geographic</u> factors or <u>human interaction</u>. For example, the rise in average temperature (see point 2 above) is due to <u>global warming</u>, which has been caused by <u>human activity</u> (see pages 116-117).

The Water Cycle Means Water is Endlessly Recycled

The water here on planet Earth is constantly <u>recycled</u>. Strange but true...

- Energy from the <u>Sun</u> makes water <u>evaporate</u> from the land and sea, turning it into <u>water</u> <u>vapour</u>. Water also evaporates from plants
 this is known as <u>transpiration</u> (see p.43).
- The warm water vapour is <u>carried upwards</u>
 (as warm air rises). When it gets higher up it cools and condenses to form clouds.
- Water falls from the clouds as <u>precipitation</u> (usually rain, but sometimes snow or hail) onto <u>land</u>, where it provides <u>fresh water</u> for <u>plants</u> and <u>animals</u>.
- If then <u>drains</u> into the <u>sea</u>, before the whole process starts again.



Come on out, it's only a little water cycle, it won't hurt you...

Because organisms rely on the biotic and abiotic factors affecting the environment (see page 107) it's no wonder that when the factors change, it affects where organisms live. Some of the non-living (abiotic) factors that affect us are in a constant cycle — read about the water cycle here and then the carbon cycle on the next page.

Q1 Give two environmental changes that can affect the distribution of living organisms. [2 marks]

Q2 a) In the water cycle, how does water move from the land into the air? [1 mark]

[1 mark]

b) How does the water cycle benefit plants and animals?


The Carbon Cycle

Recycling may be a buzz word for us but it's old school for nature. All the nutrients in our environment are constantly being recycled — there's a nice balance between what goes in and what goes out again.

Elements are Cycled Back to the Start of the Food Chain by Decay

- 1) Living things are made of materials they take from the world around them. E.g. plants turn elements like carbon, oxugen, hudrogen and nitrogen from the soil and the air into the complex compounds (carbohydrates, proteins and fats) that make up living organisms. These get passed up the food chain.
- 2) These materials are returned to the environment in waste products, or when the organisms die and decay.
- 3) Materials decay because they're broken down (digested) by microorganisms. This happens faster in warm, moist, aerobic (oxugen rich) conditions because microorganisms are more active in these conditions.
- 4) Decay puts the stuff that plants need to grow (e.g. mineral ions see point 1) back into the soil.
- 5) In a stable community, the materials that are taken out of the soil and used by plants etc. are balanced by those that are put back in. There's a constant cucle happening.

The Constant Cycling of Carbon is called the Carbon Cycle



That can look a bit complicated at first, but it's actually pretty simple:

- 1) CO is removed from the atmosphere by green plants and algae during photosunthesis. The carbon is used to make glucose, which can be turned into carbohydrates, fats and proteins that make up the bodies of the plants and algae.
- 2) When the plants and algae respire, some carbon is returned to the atmosphere as CO_a.
- 3) When the plants and algae are eaten by animals, some carbon becomes part of the fats and proteins in their bodies. The carbon then moves through the food chain.
- 4) When the animals respire, some carbon is returned to the atmosphere as CO_a.
- 5) When plants, algae and animals die, other animals (called detritus feeders) and microorganisms feed on their remains. When these organisms respire, CO2 is returned to the atmosphere.
- 6) Animals also produce waste that is broken down by detritus feeders and microorganisms.
- 7) The combustion (burning) of wood and fossil fuels also releases CO, back into the air.
- 8) So the carbon (and energy) is constantly being cycled from the air, through food chains (via plants, algae and animals, and detritus feeders and microorganisms) and eventually back out into the air again.

What goes around comes around...

Carbon is very important for living things — it's the basis for all the organic molecules in our bodies.

Q1 What causes materials to decay?

O2 Describe how carbon is removed from the atmosphere in the carbon cycle. [1 mark] [1 mark]

algae get from

photosynthesis is

transferred up the food chain.

Decay

Microorganisms break down plant and animal material and waste to get energy. Not my idea of a good meal, but to them it's positively gourmet. This process of decay can be really useful to us — read on...

Decomposition Can Produce Compost

Compost is <u>decomposed organic matter</u> (e.g. food waste) that is used as a <u>natural fertiliser</u> for crops and garden plants. Farmers and gardeners try to provide the <u>ideal conditions</u> for quick decay to make <u>compost</u>.

The Rate of Decay is Affected by Several Factors

<u>Microorganisms</u> such as <u>bacteria</u> and <u>fungi</u>, as well as <u>detritus feeders</u> (p.113) are the critters responsible for <u>decomposition</u> (decay). You'd think that organisms that feed on dead stuff wouldn't be that <u>picky</u>, but they're a bit like Goldilocks — everything has to be <u>just right</u> for them to work at their best.

- TEMPERATURE Warmer temperatures make things decompose <u>quicker</u> because they <u>increase</u>
 the <u>rate</u> that the <u>enzymes</u> (see p.28) involved in decomposition work at. If it's <u>too hot</u> though,
 decomposition <u>slows down</u> or <u>stops</u> because the enzymes are <u>destroyed</u> and the <u>organisms die.</u>
 Really <u>cold</u> temperatures slow the rate of decomposition too.
- OXYGEN AVAILABILITY Many organisms need oxygen to respire, which they need to do to survive. The microorganisms involved in anaerobic decay (see below) don't need oxygen though.
- WATER AVAILABILITY Decay takes place faster in moist environments because
 the organisms involved in decay need water to carry out biological processes.
- NUMBER OF DECAY ORGANISMS The more microorganisms and detritus feeders there are, the faster decomposition happens.

Biogas is Made by Anaerobic Decay of Waste Material

- 1) Biogas is mainly made up of methane, which can be burned as a fuel.
- 2) Lots of different microorganisms are used to produce biogas. They decay plant and animal waste anaerobically (without oxygen). This type of decay produces methane gas. Sludge waste from, for example, sewage works or sugar factories, is used to make biogas on a large scale.
- 3) Biogas is made in a simple fermenter called a digester or generator.
- 4) Biogas generators need to be kept at a constant temperature to keep the microorganisms respiring away.
- 5) Biogas can't be stored as a liquid (it needs too high a pressure), so it has to be used straight away for heating, cooking, lighting, or to power a turbine to generate electricity.

Not All Biogas Generators Are the Same

There are two main types of biogas generator — batch generators and continuous generators.

Batch generators make biogas in small batches.
They're manually loaded up with waste, which is left to digest, and the by-products are cleared away at the end of each session.

Continuous generators make biogas all the time.
Waste is continuously fed in, and biogas is
produced at a steady rate. Continuous generators
are more suited to large-scale biogas projects.

Inlet for

The diagram on the right shows a <u>simple biogas generator</u>. Whether it's a continuous or batch generator, it needs to have the following:

- 1) an inlet for waste material to be put in,
- 2) an outlet for the digested material to be removed through,
- 3) an outlet so that the biogas can be piped to where it is needed.

waste material Biogas outlet Gas Outlet for digested material needed. Waste material (to be used as fertiliser)

What a rotten subject...

Wow... So we can use poo (and other waste) to make a fuel. It's pretty gross, but I reckon it could come in handy.

Other than temperature, give three factors that affect the rate of decay.

[3 marks]

Investigating Decay



It's time for a practical. I hope you're not looking forward to a milky brew, because this might just put you off.


You Can Investigate the Effect of Temperature on the Rate of Decay

You can investigate <u>decay</u> by observing the action of the enzyme <u>lipase</u> on a sample of <u>milk</u> that has been made <u>alkaline</u>. When the lipase breaks the milk down, the <u>pH</u> of the milk <u>decreases</u>.

measuring cylinder

- Measure out <u>5 cm</u>³ of <u>lipase</u>
 <u>solution</u> and add it to a <u>test tube</u>.

 <u>Label</u> this tube with an 'L' for lipase.
- 2) Measure out <u>5 cm</u>³ of <u>milk</u> and add it to a different test tube.
- 3) Add <u>5 drops</u> of <u>phenolphthalein</u> indicator to the tube containing milk.
- 4) Then measure out 7 cm² of sodium carbonate solution and add it to the tube containing milk and phenolphthalein. This makes the solution in the tube alkaline, so it should turn pink.
- 5) Put both tubes into a water bath set to 30 °C and leave them to reach the temperature of the water bath. You could stick a thermometer into the milk tube to check this.
- 6) Once the tubes have <u>reached 30 °C</u>, use a <u>calibrated dropping pipette</u> (a dropping pipette with a scale) to put <u>1 cm</u>³ of the <u>lipase solution</u> into the milk tube and start a <u>stopwatch</u> straight away.
- 7) Stir the contents of the tube with a glass rod.
 The enzume will start to decompose the milk.
- As soon as the solution loses its pink colour, <u>stop</u> the <u>stopwatch</u> and record <u>how long</u> the <u>colour change</u> took in a table.
- 9) Repeat the experiment at a range of different temperatures (e.g. 10 °C, 20 °C, 40 °C, 50 °C). Make sure you carry out the experiment three times at each temperature, then calculate the mean time taken for the colour change to occur at each temperature. (You can make a water bath capable of temperatures below room temperature by adding ice cubes to a beaker of water and measuring the temperature with a thermometer.)
- 10) You can use your results to <u>calculate</u> the <u>rate of decay</u> using this formula. The <u>units</u> will be s⁻¹ since rate is given per unit of time.



1000

time

Rate =

You could also investigate decomposition in your gym bag...

I found that leaving my wet PE kit in my gym bag sure made for some interesting moulds... and smells.

- Q1 Ted was investigating the effect of temperature on the rate of decay of milk by measuring the time taken for a change in the pH. His results are shown in the table.
 - a) Calculate the rate of decay at 20 °C. Give your answer in s⁻¹. [1 mark]
 - b) Describe the trend in rate of decay shown by Ted's results. [1 mai

Temperature (°C)	10	20	30	40
		480		

You've met this formula

before, on page 29.

pink to white

Biodiversity and Waste Management

Unfortunately, human activity can negatively affect the planet and its variety of life. Read on for bad news...


Earth's Biodiversity is Important

Biodiversity is the variety of different species of organisms on Earth, or within an ecosystem.

- High biodiversity is important. It makes sure that ecosystems (see p.106) are <u>stable</u> because
 different species depend on each other for things like <u>shelter</u> and <u>food</u>. Different species can
 also help to maintain the right <u>physical environment</u> for each other (e.g. the acidity of the soil).
- 2) For the human species to survive, it's important that a good level of biodiversity is maintained.
- 3) Lots of human actions, including <u>waste production</u> (see below) and <u>deforestation</u> (see p.118), as well as <u>global warming</u> (see next page) are reducing biodiversity. However, it's only <u>recently</u> that we've started <u>taking measures</u> to <u>stop</u> this from continuing.

There are Over Seven Billion People in the World...

- The <u>population</u> of the world is currently <u>rising</u> very quickly, and it's not slowing down — look at the graph...
- This is mostly due to modern <u>medicine</u> and <u>farming</u> methods, which have reduced the number of people duing from disease and hunger.
- This is great for all of us <u>humans</u>, but it means we're having a bigger effect on the environment we live in.



...With Increasing Demands on the Environment

When the <u>Earth's population</u> was much smaller, the effects of <u>human activity</u> were usually <u>small</u> and <u>local</u>. Nowadays though, our actions can have a far more <u>widespread</u> effect.

- Our increasing <u>population</u> puts pressure on the <u>environment</u>, as we take the resources we need to survive.
- 2) But people around the world are also demanding a higher <u>standard of living</u> (and so demand luxuries to make life more comfortable cars, computers, etc.). So we use more <u>raw materials</u> (e.g. oil to make plastics), but we also use more <u>energy</u> for the manufacturing processes. This all means we're taking more and more resources from the environment more and more <u>quickly</u>.
- 3) Unfortunately, many raw materials are being used up quicker than they're being replaced. So if we carry on like we are, one day we're going to run out.

We're Also Producing More Waste

As we make more and more things we produce more and more <u>waste</u>, including <u>waste chemicals</u>. And unless this waste is properly handled, more <u>harmful pollution</u> will be caused. Pollution affects <u>water</u>, <u>land</u> and <u>air</u> and <u>kills</u> plants and animals, <u>reducing biodiversity</u>.



<u>Sewage</u> and <u>toxic chemicals</u> from industry can pollute lakes, rivers and oceans, affecting the plants and animals that rely on them for survival (including humans). And the <u>chemicals</u> used on land (e.g. fertilisers, pesticides and herbicides) can be washed into water.



We use <u>toxic chemicals</u> for farming (e.g. pesticides and herbicides). We also bury <u>nuclear waste</u> underground, and we dump a lot of <u>household waste</u> in landfill sites.



Smoke and acidic gases released into the atmosphere can pollute the air, e.g. sulfur dioxide can cause acid rain.

More people, more mess, less space, fewer resources...

Biodiversity's a useful thing, but the increasing rate of species extinction means that it's being reduced every day.

What is meant by the term 'biodiversity'?

[1 mark]

Global Warming

The Earth is getting <u>warmer</u>. Climate scientists are now trying to work out what the <u>effects</u> of global warming might be — sadly, it's not as simple as everyone having nicer summers.


Carbon Dioxide and Methane Trap Energy from the Sun

- 1) The temperature of the Earth is a balance between the energy it
- gets from the Sun and the energy it radiates back out into space.

 2) Gases in the atmosphere naturally act like an insulating layer.

 They absorb most of the energy that would normally be radiated out into space, and re-radiate it in all directions (including back towards the Earth).

 This increases the temperature of the planet.
- 3) If this didn't happen, then at night there'd be nothing to keep any energy in, and we'd quickly get very cold indeed. But recently we've started to worry that this effect is getting a bit out of hand...
- 4) There are several different gases in the atmosphere which help keep the energy in. They're called "greenhouse gases", and the main ones whose levels we worry about are carbon dioxide (CO₂) and methane because the levels of these two gases are rising quite sharply.
- 5) The Earth is gradually heating up because of the increasing levels of greenhouse gases this is global warming. Global warming is a type of climate change and causes other types of climate change, e.g. changing rainfall patterns.



The Consequences of Global Warming Could be Pretty Serious

There are several reasons to be worried about global warming. Here are a few:

- 1) Higher temperatures cause <u>seawater</u> to <u>expand</u> and <u>ice</u> to <u>melt</u>, causing the sea level to <u>rise</u>. It has <u>risen</u> a little bit over the last 100 years. This is beginning to <u>increase</u> the <u>frequency</u> of <u>flooding</u> in some areas. If sea level keeps <u>rising</u>, it'll be bad news for people and animals living in <u>low-luing places</u>, and could result in the <u>loss of habitats</u> (where organisms live).
- 2) The <u>distribution</u> of many <u>wild animal</u> and <u>plant species</u> is changing as <u>temperatures increase</u> and the amount of <u>rainfall changes</u> in different areas. Some species are becoming <u>more</u> widely distributed, e.g. species that need <u>warner temperatures</u> are spreading <u>further</u> as the conditions they <u>thrive</u> in exist over a <u>wider</u> area. Other species are becoming <u>less</u> widely distributed, e.g. species that need <u>cooler temperatures</u> have <u>smaller</u> ranges as the conditions they <u>thrive</u> in exist over a <u>smaller</u> area.
- There have been <u>changes in migration patterns</u>, e.g. some birds may be migrating <u>further north</u>, as more northern areas are getting warmer.
- Biodiversity (see p.116) could be <u>reduced</u> if some species are unable to survive a change in the climate, so become <u>extinct</u>.



The greenhouse effect — when you start growing into a tomato...

Global warming is rarely out of the news. Most scientists accept that it's happening and that human activity has caused most of the recent warming. However, they don't know exactly what the effects will be.

O1 Explain how global warming could lead to the loss of low-lying habitats.

[3 marks]

Deforestation and Land Use

Trees and peat bogs trap carbon dioxide and lock it up. The problems start when it escapes...

Humans Use Lots of Land for Lots of Purposes

- 1) We use land for things like building, quarruing, farming and dumping waste.
- 2) This means that there's less land available for other organisms.
- 3) Sometimes, the way we use land has a <u>bad effect</u> on the environment for example, if it requires <u>deforestation</u> or the <u>destruction</u> of habitats like <u>peat bogs</u> and other <u>areas of peat.</u>

Deforestation Means Chopping Down Trees

<u>Deforestation</u> is the <u>cutting down</u> of <u>forests</u>. This causes big problems when it's done on a <u>large-scale</u>, such as cutting down rainforests in <u>tropical areas</u>. It's done for various reasons, including:

- To <u>clear land</u> for farming (e.g. cattle or rice crops) to provide <u>more food</u>.
- To grow crops from which biofuels based on ethanol can be produced.

Deforestation Can Cause Many Problems

LESS CARBON DIOXIDE TAKEN IN

 Cutting down <u>loads of trees</u> means that the amount of carbon dioxide removed from the atmosphere during <u>photosunthesis</u> is <u>reduced</u>. More CO₂ in the atmosphere causes global warming (see previous page), which leads to climate change.

 Trees 'lock up' some of the <u>carbon</u> that they absorb during photosynthesis in their wood, which can remove it from the <u>atmosphere</u> for hundreds of years. Removing trees means that less is locked up.

MORE CARBON DIOXIDE IN THE ATMOSPHERE

- Carbon dioxide is <u>released</u> when trees are <u>burnt</u> to clear land. (Carbon in wood doesn't contribute to atmospheric pollution until it's released by burning.)
- 2) Microorganisms feeding on bits of dead wood release carbon dioxide as a waste product of respiration.

LESS BIODIVERSITY

- 1) Biodiversity (p.116) is the variety of different species the more species, the greater the biodiversity.
- Habitats like forests can contain a <u>huge number</u> of different species of <u>plants</u> and <u>animals</u>, so when
 they are destroyed there is a danger of <u>many species becoming extinct</u> biodiversity is <u>reduced</u>.

Destroying Peat Bogs Adds More CO₂ to the Atmosphere

- Bogs are areas of land that are acidic and waterlogged. Plants that live in bogs don't fully decay when
 they die, because there's not enough oxygen. The partly-rotted plants gradually build up to form peat.
- 2) So the carbon in the plants is stored in the peat instead of being released into the atmosphere.
- 3) However, peat bogs are often drained so that the area can be used as farmland, or the peat is cut up and dried to use as fuel. It's also sold to gardeners as compost. Peat is being used faster than it forms.
- 4) When peat is drained, it comes into more contact with air and some <u>microorganisms</u> start to <u>decompose</u> it. When these microorganisms <u>respire</u>, they <u>use oxygen</u> and <u>release carbon dioxide</u>, contributing to global warming (see the previous page).
- 5) Carbon dioxide is also released when peat is burned as a fuel.
- 6) Destroying the bogs also destroys (or reduces the area of) the <u>habitats</u> of some of the animals, plants and microorganisms that live there, so reduces biodiversity.

Pete Boggs Demolition Ltd — the name in habitat destruction...

So removing trees and peat results in more atmospheric CO,, which contributes to global warming. Bad times.

21 Suggest why deforestation can result in a higher CO, concentration in the atmosphere. [3 marks]

Maintaining Ecosystems and Biodiversity

It's really important that biodiversity is maintained, but other factors also have to be taken into account.

Programmes Can be Set Up to Protect Ecosystems and Biodiversity

It's important that biodiversity is maintained at a high enough level to make sure that ecosystems are stable (see page 106). In some areas, programmes have been set up by concerned citizens and scientists to minimise damage by human activities (see p.116) to ecosystems and biodiversity. Here are a few examples:

- 1) Breeding programmes have been set up to help prevent endangered species from becoming extinct. These are where animals are bred in captivity to make sure the species survives if it dies out in the wild. Individuals can sometimes be released into the wild to boost or re-establish a population.
- Programmes to protect and regenerate rare habitats like mangroves, heathland and coral reefs
 have been started. Protecting these habitats helps to protect the species that live there —
 preserving the ecosystem and biodiversity in the area.
- 3) There are programmes to reintroduce hedgerows and field margins around fields on farms where only a single tupe of crop is grown. Field margins are areas of land around the edges of fields where wild flowers and grasses are left to grow. Hedgerows and field margins provide a habitat for a wider variety of organisms than could survive in a single crop habitat.
- 4) Some governments have introduced regulations and programmes to <u>reduce</u> the level of <u>deforestation</u> taking place and the amount of <u>carbon dioxide</u> being released into the atmosphere by businesses. This could reduce the increase of <u>global warming</u> (see page 117).
- 5) People are encouraged to recycle to <u>reduce</u> the amount of <u>waste</u> that gets dumped in <u>landfill</u> sites. This could <u>reduce</u> the amount of <u>land</u> taken over for landfill, leaving <u>ecosystems</u> in place.

Conflicting Pressures Can Affect How Biodiversity is Maintained

Sadly for noble biodiversity warriors, maintaining biodiversity <u>isn't</u> as simple as you would hope. There are lots of <u>conflicting pressures</u> that have to be taken into account. For example:

- Protecting biodiversity costs money. For example, governments sometimes pay farmers a <u>subsidy</u> to reintroduce hedgerows and field margins to their land. It can also cost money to keep a <u>watch</u> on whether the programmes and regulations designed to maintain biodiversity are being <u>followed</u>. There can be <u>conflict</u> between <u>protecting biodiversity</u> and <u>saving money</u> — money may be <u>prioritised</u> for other things.
- 2) Protecting biodiversity may come at a <u>cost</u> to local people's <u>livelihood</u>. For example, reducing the amount of deforestation is <u>great</u> for <u>biodiversity</u>, but the people who were <u>previously employed</u> in the tree-felling industry could be <u>left unemployed</u>. This could affect the <u>local economy</u> if people move away with their family to find work.
- 3) There can be conflict between protecting biodiversity and protecting our <u>food security</u>. Sometimes certain organisms are seen as <u>pests</u> by farmers (e.g. locusts and foxes) and are killed to <u>protect crops</u> and <u>livestock</u> so that <u>more food</u> can be produced. As a result, however, the food chain and biodiversity can be affected.
- 4) <u>Development is important</u>, but it can affect the environment. Many people want to <u>protect</u> biodiversity in the face of development, but sometimes <u>land</u> is in such <u>high demand</u> that <u>previously untouched</u> land with <u>high biodiversity</u> has to be used for development, e.g. for housing developments on the edge of towns, or for new agricultural land in developing countries.

Revision or sleep — now that's a conflicting pressure...

Like many situations in ecology, maintaining biodiversity isn't black and white. There are lots of factors to take into account before decisions on the best way to go forward can be made.

Q1 Give an example of how biodiversity can be increased in areas that farm single crops.

[2 marks]

O2 How could wild populations of endangered species be preserved by breeding programmes?

[2 marks]

Trophic Levels

And now for something slightly different... The word 'trophic' comes from the Greek word trophe meaning 'nourishment'. This page must contain something about food then. Probably worth reading on I reckon...

Food Chains Can be Divided into Trophic Levels

- You might recognise some of this stuff about food chains from page 109. Unfortunately, now you need to learn a bit more about it. Enter the exciting world of trophic levels.
- Trophic levels are the different stages of a food chain. They consist of one or more organisms that perform a specific role in the food chain.
- 3) Trophic levels are named after their location in the food chain using numbers. The first level is called trophic level 1. Each level after that is numbered in order based on how far along the food chain the organisms in the trophic level are. For example, the organisms second in line in the food chain belong to trophic level 2, organisms third in line are in trophic level 3, and so on.

Trophic Level 1 Contains Producers

- Producers are the organisms at the <u>starting point</u> of a food chain, e.g. plants and algae.
- 2) They're called <u>producers</u> because they <u>make their own food</u> by <u>photosynthesis</u> using energy from the Sun.

Trophic Level 2 Contains Primary Consumers

- 1) Herbivores that eat the plants and algae are primary consumers.
- 2) Herbivores eat only plants and algae.


Trophic Level 3 Contains Secondary Consumers

- 1) Carnivores that eat the primary consumers are secondary consumers.
- 2) Carnivores are meat eaters.

Trophic Level 4 Contains Tertiary Consumers

- Carnivores that eat other carnivores (the secondary consumers) are tertiary consumers.
- Carnivores that have no predators are at the top of the food chain, so they're always in the <u>highest</u> trophic level.
 They're known as <u>apex predators</u>.

There can be more than four trophic levels in a food chain, but there are only usually 4 or 5 because so much energy is lost from the food chain at each trophic level (see p.122).



(producer)

Decomposers Break Down Uneaten Remains and Waste

- 1) Decomposers such as bacteria and fungi play an important role in ecosystems.
- 2) They decompose any dead plant or animal material left in an environment.
- They can do this by <u>secreting</u> (releasing) <u>enzymes</u> that <u>break</u> the dead stuff <u>down</u> into <u>small soluble food molecules</u>.

 These then <u>diffuse into</u> the microorganisms.

This process also releases nutrients into the environment, which the producers need in order to grow.

G, G, G, E, F, F, F, D — sorry, just decomposing Beethoven's 5th...

So, trophic level is just the fancy term for a stage in a food chain. Each trophic level always contains organisms that perform the same role. For example, the organisms in trophic level 3 are always the secondary consumers. And don't forget about those environmental champions, the decomposers. They clean up after everyone else.

O1 Explain how decomposers break down dead material in an environment.

[1 mark]

Pyramids of Biomass

OK, I'll be straight with you. This isn't the most interesting page in the world, but hey — life's like that.

Pyramids of Biomass Show the Relative Mass of Each Trophic Level

There's less energy and less biomass every time you move up a stage (trophic level) in a food chain. There are usually fewer organisms every time you move up a level too. Let's revisit our woodland friends from p.109

triends from p.IO: as an example:




Biomass just
means the mass
of living material.

100 dandelions... feed... 10 rabbits... which feed... one fox.

This <u>isn't</u> always true though — for example, if <u>500 fleas</u> are feeding on the fox, the number of organisms has <u>increased</u> as you move up to that stage in the food chain. So a better way to look at the food chain is often to think about <u>biomass</u> instead of number of organisms. You can use information about biomass to construct a <u>pyramid</u> of biomass to represent the food chain:

- Each bar on a <u>pyramid of biomass</u> shows the <u>relative mass of living material</u> at a <u>trophic level</u>
 — basically how much all the organisms at each level would "<u>weigh</u>" if you put them <u>all together</u>.
- So the one fox above would have a <u>big biomass</u> and the <u>hundreds of fleas</u> would have a <u>very small biomass</u>. Biomass pyramids are practically <u>always pyramid-shaped</u>:




The big bar along the bottom of the pyramid shows trophic level 1. It always represents the <u>producer</u> (e.g. plants or algae). The next bar will be the <u>primary consumer</u> (the animal that eats the producer), then the <u>secondary consumer</u> (the animal that eats the primary consumer) and so on up the food chain. Easy.

Pyramids of Biomass Give You Information About Food Chains

 It's easy to look at pyramids of biomass and explain what they show about the <u>food chain</u> just remember, the <u>biomass</u> at each stage should be <u>drawn to</u> scale. For example:

 If you're given actual numbers, you can use them to draw bars of the correct scale. Don't forget that the order of organisms in the pyramid must follow the order of the food chain, and each bar must be labelled.



Even if you know nothing about the natural world, you're probably aware that a tree is quite a bit binger than an aphid. So what's going on here is that lots (probably thousands) of aphids are feeding on a great big tree. Quite a lot of ladybirds, are then eating the aphids, and a few partidges are eating the ladybirds. Biomass and energy are still decreasing as you go up the levels — it's just that one tree can have a very bing biomass, and can fix a lot of the Sun's energy using all those leaves.

Constructing pyramids is a breeze — just ask the Egyptians...

There are a few exceptions where pyramids of biomass aren't quite pyramid-shaped. It happens when the producer has a very short life but reproduces loads, like with plankton at certain times of year. But it's very rare, so forget I mentioned it. Sorry. When you're done here, do something fun like sing to a squirrel.

Clover — Snails — Thrushes — Buzzard
Biomass: 96 kg 42 kg 8 kg 1 kg

Q1 Draw an accurate pyramid of biomass for this food chain.
Include the scale you have used. [4 marks]

Biomass Transfer

Great, steak for dinner — some biomass from a cow to help you grow. But you'll need some of that biomass for respiration, so you can move about and revise. And some will be indigestible. This is how biomass is lost...

Biomass is Lost Between Each Trophic Level

- 1) Energy from the Sun is the source of energy for nearly all life on Earth.
- 2) Producers, such as green plants and algae, use energy transferred by light from the Sun to make food (glucose) during photosynthesis. Of the energy that hits these producers, only about 1% is transferred for photosynthesis.
- Some of the glucose is used by the plants and algae to make <u>biological molecules</u>. These biological molecules make up the plant's biomass — the mass of living material. Biomass stores energy.
- 4) <u>Biomass</u> is <u>transferred</u> through a <u>food chain</u> in an ecosystem when organisms <u>eat</u> other organisms. However, <u>not much biomass</u> gets transferred from one <u>trophic level</u> to the <u>next</u>. In fact, only about <u>10%</u> of the biomass is passed on to the next level.
- 5) Biomass is lost between the levels for a few different reasons:
 - Organisms don't always eat every single part of the organism they're consuming.
 For example, some material that makes up plants and animals is inedible (e.g. bone).
 This means that not all the biomass can be passed to the next stage of the food chain.
 - Organisms don't absorb all of the stuff in the food they ingest (take in). The stuff that they
 don't absorb is egested (released) as faeces (ahem... pooped out, to you and me).
 - Some of the biomass taken in is converted into <u>other substances</u> that are lost as <u>waste</u>. For example, organisms use a lot of <u>glucose</u> (obtained from the biomass) in respiration to provide energy for movement and keeping warm, etc. rather than to make more biomass. This process produces lots of waste <u>carbon dioxide</u> and <u>water</u> as by-products. <u>Urea</u> is another waste substance, which is released in urine with water when the proteins in the biomass are broken down.

You Can Calculate the Efficiency of Biomass Transfer



- The numbers show the <u>amount of biomass</u> available to the <u>next level</u>. So <u>43 kg</u> is the amount of biomass available to the <u>greenflies</u>, and <u>4.2 kg</u> is the amount available to the <u>ladybirds</u>.
- 2) You can work out how much biomass has been <u>lost</u> at each level by taking away the biomass that is available <u>at that</u> level from the biomass that was available at the <u>previous</u> level. For example:
 Biomass lost at 2nd trophic level = 43 kg 4.2 kg = 38.8 kg.
- 3) You can also calculate the efficiency of biomass transfer between trophic levels:

efficiency = $\frac{\text{biomass transferred to the next level}}{\text{biomass available at the previous level}} \times 100$

So, the efficiency of biomass transfer from the 1st trophic level = 4.2 kg \div 43 kg \times 100 = 9.8%.

Spaghetti necklaces are my favourite type of food chain...

Remember, hardly any of the biomass from one trophic level is passed on to the next level — most is lost.

Q1 Give two ways that biomass is lost between trophic levels.

[2 marks] Q2 Vid

Q2 A shark with 110 kg of available biomass eats some large fish with 995 kg of available biomass.

[1 mark] [1 mark]

a) Calculate the amount of biomass lost between the fish and the shark.

b) Calculate the percentage efficiency of biomass transfer between the fish and the shark.

Sustainable methods of food

production are needed so that enough food can be made to feed everyone now and in the

future. Sustainable production means making enough food

without using resources faster

than they renew.

Food Security and Farming

There are more than 7 billion people on our planet and they all need feeding, so maintaining our food supply is important. We need to make sure that we use sustainable methods of food production wherever we can.

Lots of Factors Affect Food Security

<u>Food security</u> is having <u>enough food</u> to <u>feed</u> a <u>population</u>.

There's a <u>wide range</u> of things that can <u>threaten</u> food security. For example:

- 1) The world population keeps increasing, with the birth rate of many developing countries rising quickly.
- As diets in <u>developed countries change</u>, the <u>demand</u> for certain foods to be <u>imported</u> from <u>developing</u> countries can increase. This means that already scarce food resources can become more scarce.
- 3) Farming can be affected by new pests and pathogens (e.g. bacteria and viruses) or changes in the environmental conditions (e.g. a lack of rain). This can result in the loss of crops and livestock, and can lead to widespread famine.
- 4) The high input costs of farming (e.g. the price of seeds, machinery and livestock) can make it too expensive for people in some countries to start or maintain food production, meaning that there sometimes aren't enough people producing food in these areas to feed the people.
- In some parts of the world, there are <u>conflicts</u> that affect the <u>availability</u> of <u>food</u> and <u>water</u>.

Overfishing is Decreasing Fish Stocks

- 1) Fish stocks are declining in the oceans because we're fishing so much.
- 2) This means there's less fish for us to eat, the ocean's food chains are affected and some species of fish may disappear altogether in some areas for example, cod are at risk of disappearing from the north west Atlantic.
- 3) To tackle this problem, we need to maintain fish stocks at a level where the fish continue to breed.

 This is <u>sustainable food production</u>. Fish stocks can be <u>maintained</u> (<u>conserved</u>) in these ways:

FISHING QUOTAS — there are <u>limits</u> on the number and size of fish that can be caught in certain areas. This <u>prevents</u> certain species from being <u>overfished</u>.

NET SIZE — there are different limits of the mesh size of the fish net, depending on what's being fished. This is to reduce the number of 'unwanted' and discarded fish — the ones that are accidently caught, e.g. shrimp caught along with cod. Using a bigger mesh size will let the 'unwanted' species escape. It also means that younger fish will slip through the net, allowing them to reach breeding age.

Food Production Can be Made More Efficient

<u>Limiting</u> the <u>movement</u> of livestock and keeping them in a <u>temperature-controlled environment</u> reduces the <u>transfer</u> of <u>energy</u> from livestock to the environment. This makes farming <u>more efficient</u> as the animals use less energy <u>moving</u> around and <u>controlling</u> their own <u>body temperature</u>. This means that <u>more energy</u> is available for <u>growth</u>, so <u>more food</u> can be produced from the <u>same input</u> of resources.

- 1) Livestock like calves and chickens can be factory farmed. This involves raising them in small pens.
- 2) Fish can also be factory farmed in cages where their movement is restricted.
- 3) Some animals are also fed high-protein food to further increase their growth.

 Some factory farming methods are controversial. Because the animals are kept so controversial. Because the animals are kept so ethical objections, as some people think that making animals live in unnatural and uncomfortable conditions is gruel.

Fishermen are just too effishent...

Food security is seriously important, but there are lots of issues that threaten it. There are ways to reduce the threats though. Make sure you're familiar with the stuff on this page and your exam mark security may be vastly improved.

O1 Explain how food production from livestock farming can be made more efficient. [4 marks]

Biotechnology

Biotechnology is where living things and biological processes are manipulated to produce a useful product.

Mycoprotein — Food from Fungi

- Using modern biotechnology techniques, large amounts of microorganisms can be cultured (grown) industrially under controlled conditions in large vats for use as a food source.
- 2) Mycoprotein is used to make high-protein meat substitutes for vegetarian meals, e.g. Quorn™.
- 3) It's made from the fungus <u>Fusarium</u>, which is grown in aerobic conditions on glucose surup, which it uses as food.
- 4) The fungal biomass is harvested and purified to produce the mucoprotein.



sticky ends

plasmid

The bacteria are

grown like mad

in a vat.

Restriction enzumes

...ligase joins the two bits of DNA together...

... and the

recombinant DNA

is inserted into

the bacteria.

cut the gene out

and cut open

the plasmid...

Bacteria can be Engineered to Produce Human Insulin

Genetic engineering is transferring a <u>useful gene</u> from one organism to another (see page 99).

Bacteria can be genetically engineered to <u>make human insulin</u>: human insulin gene

- 1) A <u>plasmid</u> (a loop of DNA) is <u>removed</u> from a bacterium.
- 2) The insulin gene is cut out of a human chromosome using a restriction enzyme. Restriction enzymes recognise specific sequences of DNA and cut the DNA at these points. The cut leaves one of the DNA strands with unpaired bases this is called a 'sticky end'.
- The <u>plasmid</u> is cut open using the <u>same</u> restriction enzyme — leaving the <u>same sticky ends</u>.
- 4) The plasmid and the human insulin gene are mixed together.
- Ligase (an enzyme) is added. This joins the sticky ends together to produce recombinant DNA (two different bits of DNA stuck together).
- 6) The recombinant DNA is inserted into a bacterium.
- 7) The modified bacterium is grown in a vat under controlled conditions. You end up with millions of bacteria that produce insulin. The insulin can be harvested and purified to treat people with diabetes.

Crops Can be Genetically Modified

Many people in the world today don't have <u>enough</u> food to eat (or the diet they have isn't <u>varied</u>). This mostly happens in <u>developing countries</u> — like those in <u>Africa</u> and parts of <u>Asia</u>.

Biotechnology could help...

- 1) Genetically modified (GM) crops can be produced that are resistant to pests improving crop yields.
- 2) They can be genetically modified to grow better in drought conditions again improving crop yields.
- 3) And some crops can be modified to provide more nutritional value, e.g. 'Golden Rice' has been genetically engineered to produce a chemical that's converted in the body to vitamin A.

...But not everyone agrees

- Many people argue that people go hungry because they <u>can't afford</u> to buy food, not because there <u>isn't</u> any food about. So they argue that you need to <u>tackle poverty first</u>.
- 2) There are fears that countries may become dependent on companies who sell GM seeds.
- 3) Sometimes poor soil is the main reason why crops fail, and even GM crops won't survive.

I'll have a fungus sandwich with a side of GM rice please...

Genetic modification is a powerful biotechnological tool. It could hold the solution to a wide range of problems.

Q1 Rennin is an enzyme used to make cheese. It is naturally produced by stomach cells in cows, which contain the rennin gene. Suggest how a bacterial cell could be genetically engineered to produce rennin.

ns. Ol Video Solution

[3 marks]



Revision Questions for Topic 7

That's Topic 7 done with. It must be time for some revision questions...

- Try these questions and tick off each one when you get it right.

27) How can bacteria be used to produce human insulin?

For even more practice, try the

Retrieval Quiz for Topic 7 — just

scan this QR code!

Topic 7 Ouiz

• When you're completely happy with a sub-topic, tick it off.	Topic I Quiz
Competition, Abiotic and Biotic Factors, and Adaptations (p.106-108)	Quiz
1) Define 'habitat'.	\checkmark
2) What things do animals compete for in an ecosystem?	\checkmark
3) What are biotic and abiotic factors?	\checkmark
4) What are functional adaptations?	\checkmark
Food Chains (p.109)	
5) What do food chains always start with?	\checkmark
6) Explain what happens to the population size of a predator	_
if its prey becomes more common in an ecosystem.	\checkmark
Quadrats and Transects (p.110-111)	
7) Explain how a quadrat can be used to investigate the distribution of clover plants in two areas.	\checkmark
8) Suggest why you might use a transect when investigating the distribution of organisms.	\checkmark
Environmental Change, and The Water and Carbon Cycles (p.112-113)	
9) How might a change in the availability of water affect the distribution of species?	\checkmark
10) When water vapour cools and condenses in the atmosphere, what does it change into?	\checkmark
11) Explain how microorganisms return carbon to the atmosphere.	\checkmark
Decay (p.114-115)	
12) Explain why temperature affects the rate of decay.	
13) Is biogas produced by aerobic or anaerobic decay?	
Human Impacts on the Planet (p.116-119)	
14) Suggest why it's important to have high biodiversity in an ecosystem.	V
15) Name two gases linked to global warming.	<u></u>
16) Give an example of how global warming could reduce biodiversity.	
17) Why might humans carry out deforestation?	
18) Explain why the destruction of peat bogs adds more carbon dioxide to the atmosphere.	<u></u>
19) How can recycling programmes help to protect ecosystems?	\checkmark
Trophic Levels and Biomass (p.120-122)	
20) What is a trophic level?	
21) In which trophic level are the tertiary consumers found?	
22) Does the bottom bar on a pyramid of biomass represent the tertiary consumer or the producer?	
23) Approximately what percentage of biomass is lost between trophic levels?	
Food Security and Biotechnology (p.123-124)	
24) Give an example of a factor that can affect food security.	
25) How can fish stocks be maintained at a sustainable level?	
26) What is mycoprotein?	

Measuring Substances

Get your lab coats on, it's time to find out about the skills you'll need in experiments...

Use the Right Apparatus to Take Readings

1) Mass

To weigh a solid, start by putting the <u>container</u> you are weighing your substance into on a <u>balance</u>. Set the balance to exactly <u>zero</u> and then weigh out the correct amount of your substance. Easy peasy.

2) Temperature

You can use a <u>thermometer</u> to measure the temperature of a solution. Make sure that the <u>bulb</u> of the thermometer is <u>completely submerged</u> in the solution and that you wait for the temperature to <u>stabilise</u> before you take your initial reading. Read off the scale on the thermometer at <u>eue level</u> to make sure it's correct.

When you're reading off a scale, write down the value of the graduation that the amount is closest to.

Read volume

from here —

3) Volume of a Liquid


There's more than one way to measure the volume of a <u>liquid</u>. Whichever method you use, always read the volume from the <u>bottom of the meniscus</u> (the curved upper surface of the liquid) when it's at <u>eye level</u>.

Using a pipette — Pipettes are used to suck up and transfer volumes of liquid between containers. Dropping pipettes are used to transfer drops of liquid. Graduated pipettes are used to transfer accurate volumes. A pipette filler is attached to the end of a graduated pipette, to control the amount of liquid being drawn up.

<u>Using a measuring cylinder</u> — <u>Measuring cylinders</u> come in all different <u>sizes</u>. Make sure you choose one that's the right size for the measurement you want to make. It's no good using a huge I dm³ cylinder to measure out 2 cm² of a liquid — the graduations will be too big, and you'll end up with <u>massive errors</u>. It'd be much better to use one that measures up to 10 cm³.

4) Volume of a Gas

- 1) To accurately measure the volume of gas, you should use a gas syringe.
- 2) Alternatively, you can use an <u>upturned measuring cylinder filled with water.</u>
 The gas will <u>displace</u> the water so you can <u>read the volume</u> off the <u>scale</u>.



4) When you're measuring a gas, you need to make sure that the equipment is set up so that none of the gas can escape, otherwise your results won't be accurate.

5) pH

The method you should use to measure pH depends on what your experiment is.

- Indicators are dues that change colour depending on whether they're in an acid or an alkali.
 You use them by adding a couple of drops of the indicator to the solution you're interested in.
 Universal indicator is a mixture of indicators that changes colour gradually as pH changes.
 It's useful for estimating the pH of a solution based on its colour.
- Indicator paper is useful if you don't want to colour the entire solution that
 you're testing. It changes colour depending on the pH of the solution it touches.
 You can also hold a piece of damp indicator paper in a gas sample to test its pH.

3) pH meters have a digital display that gives an accurate value for the pH of a solution.

Blue litmus paper turns red in acidic conditions and red litmus paper turns blue in alkaline conditions.

Experimentus apparatus...

Wizardry won't help you here, unfortunately. It's best you just get your head down and learn this stuff.



Heating, Safety and Ethics

More lab stuff coming up next — a bit about heating things up and then some important things to think about...

Bunsen Burners Have a Naked Flame

Bunsen burners are good for heating things quicklu. But you need to make sure you're using them safelu:

- You should always use a Bunsen burner on a <u>heat-proof mat</u>.
- If your Bunsen burner is alight but not heating anything, make sure
 you close the hole so that the flame becomes yellow and clearly visible.
- Use the <u>blue</u> flame to heat things. If you're heating a vessel in the flame, hold
 it at the <u>top</u> (e.g. with <u>tongs</u>) and point the opening <u>away from</u> yourself (and others).
- If you're heating something over the flame (e.g. a beaker of water), you should
 put a tripod and gauze over the Bunsen burner before you light it, and place the vessel on this.



The Temperature of Electric Water Baths & Electric Heaters Can Be Set

- A <u>water bath</u> is a container filled with water that can be heated to a <u>specific temperature</u>.
 A <u>simple</u> water bath can be made by heating a <u>beaker of water</u> over a <u>Bunsen burner</u> and monitoring the temperature with a <u>thermometer</u>. However, it is difficult to keep the temperature of the water <u>constant</u>.
- 2) An electric water bath will monitor and adjust the temperature for you. Here's how you use one:
 - · Set the temperature on the water bath, and allow the water to heat up.
 - To make sure it's reached the right temperature, use a thermometer.
 - Place the vessel containing your substance in the water bath using test tube holders or tongs.
 The level of water outside the vessel should be <u>just above</u> the level of the substance in the vessel.
 - The substance will then be warmed to the <u>same temperature</u> as the water.
 As the substance in the vessel is surrounded by water, the heating is very <u>even</u>.
- 3) Electric heaters are often made up of a metal plate that can be heated to a specified temperature. The vessel containing the substance you want to heat is placed on top of the hot plate. The vessel is only heated from below, so you'll usually have to stir the substance inside to make sure it's heated evenly.

Make Sure You're Working Safely in the Lab

- 1) <u>Before</u> you start any experiment, make sure you know about any <u>safety precautions</u> to do with your <u>method</u> or the <u>chemicals</u> you're using. You need to <u>follow</u> any instructions that your teacher gives you <u>carefully</u>. The chemicals you're using may be <u>hazardous</u> for example, they might <u>be flammable</u> (<u>catch fire easilu</u>), or they might <u>irritate</u> or <u>burn</u> your <u>skin</u> if it comes into contact with them.
- 2) Make sure that you're wearing <u>sensible clothing</u> when you're in the lab (e.g. open shoes won't protect your feet from spillages). When you're doing an experiment, you should wear a <u>lab coat</u> to protect your skin and clothing. Depending on the experiment, you may need to also wear <u>safety goggles</u> and <u>gloves</u>.
- You also need to be aware of general safety in the lab, e.g. keep anything flammable away from lit Bunsen burners, don't directly touch any hot equipment, handle glassware carefully so it doesn't break, etc.

You Need to Think About Ethical Issues In Your Experiments

- 1) Any organisms involved in your investigations need to be treated safely and ethically.
- Animals need to be treated <u>humanely</u> they should be <u>handled carefully</u> and any wild animals
 captured for studying (e.g. during an investigation of the distribution of an organism) should be
 returned to their original habitat.
- Any animals kept in the lab should also be <u>cared for</u> in a humane way, e.g. they should not be kept in <u>overcrowded conditions</u>.
- 4) If you are carrying out an experiment involving other <u>students</u> (e.g. investigating the effect of caffeine on reaction time), they should not be forced to participate <u>against their will</u> or feel <u>pressured</u> to take part.

Potometers and Microscopes


We're on to potometers and microscopes now, oh my...

Potometers Should Be Set Up Underwater

A potometer is a special piece of apparatus used to measure the water uptake by a plant. Here's how to set one up:

1) Cut a shoot underwater to prevent air from entering the xylem. Cut it at a slant to increase the surface area available for water uptake.

- 2) Assemble the potometer in water and insert the shoot under water, so no air can enter.
- 3) Remove the apparatus from the water but keep the end of the capillary tube submerged in a beaker of water.
- 4) Check that the apparatus is watertight and airtight.
- 5) Dry the leaves, allow time for the shoot to acclimatise and then shut the tap.
- 6) Remove the end of the capillary tube from the beaker of water until one air bubble has formed, then put the end of the tube back into the water.
- 7) A potometer can be used to estimate the transpiration rate of a plant. There's more about this on page 44.




You Can Measure the Size of a Single Cell

When viewing cells under a microscope. you might need to work out their size.

To work out the size of a single cell:

- 1) Place a clear, plastic ruler on top of your microscope slide. Clip the ruler and slide onto the stage.
- 2) Select the objective lens that gives an overall magnification of $\times 100$.
- 3) Adjust the focus to get a clear image of the cells.
- 4) Move the ruler so that the cells are lined up along 1 mm. Then count the number of cells along this 1 mm sample.
- 5) 1 mm = 1000 μ m. So to calculate the length of a single cell in μ m, you just need to divide 1000 μm by the number of cells in the sample. E.g. if you counted 4 cells in 1 mm, the length of a single cell would be: 1000 ÷ 4 = 250 µm

Use the Cell Size to Work out the Length of a Scale Bar






3) To draw a 500 µm scale bar, drawn length of cell (µm) x 500 just use this formula: scale bar length (μ m) = actual length of cell (μm)

Single cell, looking for love...

These techniques might seem a bit tricky at first, but if you practice them they'll become second nature. Honest.



drawing of cell

500 μm

Sampling

I love <u>samples</u>... especially when I'm a bit <u>peckish</u> in the supermarket and they're handing out <u>free cheese</u>. Unfortunately, this page isn't about those samples. It's a lot more useful than that...

Sampling Should be Random

- When you're investigating a population, it's generally not possible to study every single organism
 in the population. This means that you need to take <u>samples</u> of the population you're interested in.
- The sample data will be used to <u>draw conclusions</u> about the <u>whole</u> population, so it's important that it <u>accurately</u> represents the <u>whole population</u>.
- 3) To make sure a sample represents the population, it should be random.


If a sample doesn't represent the population as a whole, it's said to be biased.


Organisms Should Be Sampled At Random Sites in an Area

- If you're interested in the <u>distribution</u> of an organism in an area, or its <u>population size</u>, you can take population samples in the area you're interested in using <u>quadrats</u> or <u>transects</u> (see pages 110-111).
- If you only take samples from one part of the area, your results will be biased they may not give
 an accurate representation of the whole area.
- 3) To make sure that your sampling isn't biased, you need to use a method of choosing sampling sites in which every site has an equal chance of being chosen. For example:

If you're looking at plant species in a field...

- 1) Divide the field into a grid.
- Label the grid along the bottom and up the side with numbers.
- Use a <u>random number</u> <u>generator</u> (on a computer or calculator) to select coordinates, e.g. (2,6).
- Take your samples at these coordinates.





Health Data Should be Taken from Randomly Selected People

- 1) As mentioned above, it's not practical (or even possible) to study an entire human population.
- 2) You need to use <u>random sampling</u> to choose members of the population you're interested in. For example:

A <u>health professional</u> is investigating <u>how many</u> people diagnosed with <u>Type 2 diabetes</u> in a particular country <u>also</u> have <u>heart disease</u>:

- All the people who have been diagnosed with Type 2 diabetes in the country of interest are identified by hospital records. In total, there are 270 196 people.
- 2) These people are assigned a number between 1 and 270 196.
- A random number generator is used to choose the sample group (e.g. it selects the individuals #72 063, #11 822, #193 123, etc.)
- 4) The proportion of people in the sample that have heart disease can be used to estimate the total number of people with Type 2 diabetes that also have heart disease.

'Eeny, meeny, miny, moe' just doesn't cut it any more...

Sampling is an important part of an investigation. It needs to be done randomly, or the data won't be worth much.

Comparing Results

Being able to compare your results is really important. Here are some ways you might do it. I spoil you.

Percentage Change Allows you to Compare Results

- 1) When investigating the change in a variable, you may want to compare results that didn't have the same initial value. For example, you may want to compare the change in mass of potato cylinders left in different concentrations of sugar solution that had different initial masses (see page 21).
- 2) One way to do this is to calculate the percentage change. You work it out like this:

percentage (%) change =
$$\frac{\text{final value - original value}}{\text{original value}} \times 100$$

EXAMPLE

A student is investigating the effect of the concentration of sugar solution on potato cells.

She records the mass of potato cylinders before and after placing them in sugar solutions of different concentrations.

The table on the right shows some of her results. Which potato cylinder had the largest percentage change?

cylinder	(mol/dm ³)	start (g)	end (g)
1	0.0	7.5	8.7
2	1.0	8.O	6.8

1) Stick each set of results into the equation:

$$\% change = \frac{final \ value - original \ value}{original \ value} \times 100$$

2) Sompare the results. 16% is greater than 15%, so the potato cylinder in the O.O mol/dm3 sugar solution had the largest percentage change.


Percentiles Tell you Where in Your Data Set a Data Point Lies

- Percentiles are useful if you want to compare the value of one data point to the rest of your data.
- 2) To find a percentile, you rank your data from smallest to largest, then divide it into one hundred equal chunks. Each chunk is one percentile.
- 3) This means that each percentile represents one percent of the data, and so the value of a percentile tells you what percentage of the data has a value lower than the data points in that percentile.

E.g. Mike the Meerkat is in the 90th percentile for height in his gang. This means that 90% of the gang are shorter than Mike.



4) Percentiles can be used to give a more realistic idea of the spread of data than the range (see p.6) by finding the range between the 10th and 90th percentiles in a data set (the middle 80% of the data), you can look at the spread of the data while ignoring any outlying results.



Percentage change in how much I love maths after this page — 0%

Aaaand that's the end of Practical Skills, folks. Go forth, and science like you've never scienced before...

p.11 — Cells

Q1 Any two from: e.g. prokaryotic cells are smaller than eukaryotic cells [1 mark]. / Prokaryotic cells do [1 mark]. / Prokaryotic cells do [1 mark]. / Prokaryotic cells don't have a true nucleus but eukaryotic cells do [1 mark]. / Prokaryotic cells have circular DNA but eukaryotic cells have circular DNA but eukaryotic cells don't [1 mark].

p.12 - Microscopy

Q1 real size = image size + magnification = 2.4 mm ÷ 40 = 0.06 mm [1 mark] 0.06 × 1000 = 60 µm [1 mark]

p.13 — More on Microscopy

Q1 To highlight objects within the sample by adding colour to them [1 mark].

p.14 — Cell Differentiation and

Specialisation

Q1 E.g. they have few subcellular structures.

/ They're joined end to end. / They're long.

// mark//

Q2 The cell has a hair-like shape, which gives it a large surface area [1 mark] to absorb water and minerals from the soil [1 mark].

p.15 — Chromosomes and Mitosis

Q1a) 11 ÷ (62 + 11) = 0.150... 0.150... × 100 = 15% [1 mark]

b) E.g. she could see the X-shaped chromosomes in the middle of the cells. / She could see the arms of the chromosomes being pulled apart [1 mark].

p. 16 — Binary Fission

Q1 Lots of nutrients [1 mark] and a warm environment [1 mark].

Q2 2 hours = 60 × 2 = 120 minutes 120 minutes + 40 minutes = 160 minutes 160 ÷ 20 = 8 divisions [1 mark] 2⁶ = 2 × 2 × 2 × 2 × 2 × 2 × 2 × 2 = 256 cells [1 mark]

p.18 - Culturing Microorganisms

Q1a) A [1 mark]

b) diameter = 13 mm radius = 13 ÷ 2 = 6.5 mm [1 mark] $\pi r^2 = \pi \times 6.5^2 = 132.7...$ = 133 mm² [1 mark]

 c) E.g. a paper disc soaked in sterile water [1 mark].

p.19 - Stem Cells

Q1 Copies of the plant can be made by taking stem cells from the meristem of the plant [1 mark] and growing them into new, genetically identical plants (clones) [1 mark].

p.20 - Diffusion

Q1a) The ink will diffuse / spread out through the water [1 mark]. This is because the ink particles will move from where is a higher concentration of them (the drop of ink) to where there is a lower concentration of them (the surrounding water) [1 mark].

b) The ink particles will diffuse / spread out faster [1 mark].

p.21 — Osmosis

Q1 Water will move out of the piece of potato by osmosis [1 mark], so its mass will decrease [1 mark].

p.22 - Active Transport

Q1 Active transport allows nutrients such as glucose to move from a lower concentration in the gut to a higher concentration in the blood (against the concentration gradient) 11 markl.

p.23 - Exchange Surfaces

Q1 Surface area: (2 × 2) × 2 = 8

(2 × 1) × 4 = 8 8 + 8 = 16 µm² [1 mark]

Volume: $2 \times 2 \times 1 = 4 \mu m^3$ [1 mark] So the surface area to volume ratio is 16 : 4, or 4 : 1 [1 mark].

p.24 — Exchanging Substances

Q1 E.g. they have a large surface area. / They have a moist lining for dissolving gases. / They have very thin walls. / They have a good blood supply. [1 mark]

Q2 The damage to the villi is likely to reduce the surface area for absorption [1 mark]. Therefore, less iron can be absorbed from the digested food in the small intestine into the blood [1 mark].

p.25 — More on Exchanging Substances

Substances

11 Any two from: e.g. it is made up of gill filaments which give a large surface area. / Each gill filament is covered in lamellae, which further increases the surface area. / The lamellae have a thin surface layer of cells. / The lamellae have olds of capillaries. / A large concentration gradient is maintained between the water and the blood. / 2 marks)

p.27-Cell Organisation

Q1 That it is made up of different tissues [1 mark] that work together to perform a particular function [1 mark].

p.28 — Enzymes

Q1 If the pH is too high or too low, it can interfere with the bonds holding the enzyme together. This changes the shape of the active site [1 mark] and denatures the enzyme [1 mark].

p.29 — Investigating Enzymatic Reactions

Q1 2 minutes = 2 × 60 = 120 seconds 36 ÷ 120 = 0.3 cm³/s [1 mark]

p.30 - Enzymes and Digestion

Ot Bile is alkaline, so it neutralises the stomach acid and makes conditions in the small intestine alkaline [1 mark]. The enzymes of the small intestine far mark]. The stail in these alkaline conditions [1 mark]. It also emulsifies faits/breaks down fats into tiny droplets [1 mark]. This gives a bigger surface area of fat for the enzyme lipase to work on, making digestion faster [1 mark].

p.31 — More on Enzymes and Digestion

Q1 Stomach [1 mark], pancreas [1 mark] and small intestine [1 mark].

p.32 - Food Tests

Q1 iodine solution [1 mark]

p.33 — The Lungs

Q1 495 ÷ 12

= 41 breaths per minute. [1 mark]

p.34 — Circulatory System —

The Heart
The right ventricle [1 mark].

Q2 They supply oxygenated blood to the heart itself [1 mark].

p.35 — Circulatory System — Blood Vessels

Q1 2.175 | × 1000 = 2175 ml [1 mark] 2175 + 8.7 = 250 ml/min [1 mark]

p.36 - Circulatory System - Blood

Q1 They help the blood to clot at a wound [1] mark].

Q2 To carry oxygen from the lungs to all the cells in the body [1 mark].

p.37 — Cardiovascular Disease

Q1a) Stents can be inserted into the coronary arteries to keep them open [1 mark]. This ensures that the supply of oxygenated blood to the heart isn't interrupted [1 mark].

b) Any two from: e.g. there is a risk of complications such as a heart attack during the operation. / There's a risk of infection from surgery. / There's a risk of the patient developing a blood clot/thrombosis near the stent [2 marks].

p.38 — More on Cardiovascular Disease

Q1a) The heart valves might not be able to open fully, meaning that less blood can flow through then [1 mark]. They can also become leaky, meaning that blood is able to flow in both directions [1 mark].

b) By replacing the valve with a biological or mechanical valve [1 mark].

Q2 Any one from: e.g. surgery to fit an artificial heart can lead to bleeding and infection.
/ Parts of the artificial heart could wear out. / The electric motor could fail. / Blood doesn't flow through the heart as smoothly, which could cause clots and lead to strokes. / The patient has to take blood thinning drugs to prevent blood clots, which could cause problems with bleeding if they're hurt in an accident / mark!.

p.39 — Health and Disease

Q1 The state of physical and mental wellbeing [1 mark].

Q2 It can be spread from person to person [1 mark].

p.40 - Risk Factors for

Non-Communicable Diseases

The presence of certain substances in the body. / The presence of certain substances in the environment [1 mark].

p.41 - Cancer

- Q1 Body cells dividing out of control [1 mark]. Q2 Any three from: e.g. smoking / obesity / UV
- exposure / viral infection [1 mark for each correct answer, up to 3 marks].

p.42 - Plant Cell Organisation

Q1 Meristem tissue is found at the growing tips of roots and shoots [1 mark] and is able to differentiate into lots of different types of plant cell [1 mark].

p.43 — Transpiration and Translocation

Q1 Xylem is made up of dead cells joined together end to end [1 mark] with no end walls between them and a hole down the middle [1 mark]. It is strengthened by lignin [1 mark]

p.44 - Transpiration and Stomata

Q1 Aloe vera, because the transpiration rate will be higher in the hot, dry area [1 mark]. so the aloe vera will have fewer stomata to help conserve water [1 mark].

p.46 - Communicable Disease

Q1 Viruses replicate themselves by using the machinery of the cell they live in to produce many copies of themselves [1 mark]. The cell will usually then burst, releasing all the new viruses [1 mark].

p.47 - Viral, Fungal and Protist Diseases

- a red skin rash [1 mark]
- Using a fungicide [1 mark] and by stripping the plants of the affected leaves and destroying them [1 mark].

p.48 - Bacterial Diseases and Preventing Disease

- Strains of the bacteria becoming resistant to antibiotics/penicillin [1 mark]
- To prevent the contamination of food by disease-causing pathogens [1 mark].

p.49 — Fighting Disease

- Q1 It is when white blood cells engulf foreign cells and digest them [1 mark].
- They secrete mucus to trap pathogens [1 mark]. They have cilia [1 mark], which waft the mucus up to the back of the throat where it can be swallowed [1 mark]

p.50 — Fighting Disease — Vaccination

Q1 Basia's white blood cells recognise the antigens on the flu virus and rapidly produce antibodies, which kill the pathogen [1 mark]. Cassian's white blood cells don't recognise the antigens, so it takes a while for them to produce antibodies and he becomes ill in the meantime [1 mark].

p.51 — Fighting Disease — Drugs

Q1 bacteria [1 mark]

p.52 — Developing Drugs

- Whether the drug works and produces the effect you're looking for [1 mark].
- To make sure that the drug doesn't have any harmful side effects when the body is working normally [1 mark]
- To help to prevent false claims about the results [1 mark].

p.53 - Monoclonal Antibodies

Q1 A B-lymphocyte [1 mark] and a tumour cell [1 mark].

p.54 - More on Monoclonal Antibodies

The fluorescent antibodies will bind to the antigens on the bacteria [1 mark], so any bacteria that are present in the sample will be detected / will glow [1 mark].

p.55 - Plant Diseases and Defences Q1 It will have stunted growth [1 mark].

p.57 — Photosynthesis and Limiting Factors

Glucose [1 mark] and oxygen [1 mark]. Light intensity [1 mark], volume of CO. [1 mark] and amount of chlorophyll [1 mark].

p.60 - The Rate of Photosynthesis

E.g. light intensity [1 mark], CO2 [1 mark]. light intensity $\propto \frac{1}{\text{distance}^2}$

1 ÷ 152 = 0.00444... a.u. [1 mark] 1 + 52 = 0.04 a.u. [1 mark] 0.04 ÷ 0.00444... = 9 [1 mark]

p.61 - Respiration and Metabolism

- Q1 Any two from: e.g. to build up larger molecules from smaller ones. / To contract muscles. / To keep body temperature steady. [2 marks]
- The sum of all of the reactions that happen in a cell or the body [1 mark].

p.62 - Aerobic and Anaerobic Respiration

- Glucose [1 mark] and oxygen [1 mark].
- Q2 fermentation [1 mark]

p.63 - Exercise

Running [1 mark]. It raises the pulse rate the most, so it is the most vigorous type of exercise [1 mark]. The more vigorous the exercise, the more anaerobic respiration will be taking place in the muscles [1 mark]. Anaerobic respiration produces lactic acid, so running will lead to the greatest build up of lactic acid in the blood [1 mark].

p.65 — Homeostasis

- Q1 To maintain the right conditions for cells to function properly and for enzyme action [1 mark]
- Q2 receptor (cell) [1 mark]

p.66 - The Nervous System

Q1 Muscles [1 mark], glands [1 mark].

p.67 — Synapses and Reflexes

Q1 A rapid, automatic response to a stimulus that doesn't involve the conscious part of the brain [1 mark].

Q2a) muscle [1 mark]

b) The heat stimulus is detected by receptors in the hand [1 mark], which send impulses along a sensory neurone to the CNS [1 mark]. The impulses are transferred to a relay neurone [1 mark]. They are then transferred to a motor neurone and travel along it to the effector/muscle [1 mark].

p.68 — Investigating Reaction Time

Q1a) 242 + 256 + 253 + 249 + 235 = 1235 [1 mark] 1235 ÷ 5 = 247 ms [1 mark]

b) Any two from: e.g. the hand each person used to click the mouse / the computer equipment/programme used / the amount of energy drink they consumed / the type of energy drink used / the time between consuming the energy drink and taking the test [2 marks].

p.69 - The Brain

Q1 cerebellum [1 mark]

p.70 — The Eye

Q1 cornea [1 mark]

Q2 The ciliary muscles contract [1 mark], which slackens the suspensory ligaments

[1 mark]. The lens becomes fat/more curved [1 mark]. This increases the amount by which it refracts light, so the image is focused on the retina [1 mark].

p.71 - Correcting Vision Defects

- Q1 The lens is the wrong shape / refracts the light too much, or the eveball is too long [1 mark]. So images of distant objects are brought into focus in front of the retina [1 mark].
- Q2 The lenses refract the light rays so they focus on the retina [1 mark].

p.72 — Controlling Body Temperature Q1 thermoregulatory centre/hypothalamus [1 mark]

p.73 — The Endocrine System

Q1 Because it produces many hormones that act on other glands to regulate body conditions [1 mark].

p.74 — Controlling Blood Glucose

Q1 Curve 1, because the secretion rate is high when the blood glucose level is low / the secretion rate decreases as the blood glucose level rises [1 mark]. Glucagon increases the blood ducose level, so it is secreted when the blood glucose level becomes too low [1 mark].

p.75 — The Kidneys

Q1 The kidneys will reabsorb less water [1 mark], so too much water may be lost from the body in urine [1 mark].

p.76 — Kidney Failure

Q1 It allows ions and waste substances to diffuse out of the blood into the dialysis fluid [1 mark], but not large molecules like proteins (so they won't be lost from the blood) [1 mark].

p.77 — Puberty and the Menstrual Cycle

- Q1 FSH/follicle-stimulating hormone [1 mark]
- Q2 testes [1 mark]

p.78 — Controlling Fertility

Q1 The pill/oral contraceptives [1 mark].
The contraceptive patch [1 mark].

p.79 — More on Controlling Fertility

- Q1 They stimulate several eggs to mature [1 mark].
- Q2 E.g. it doesn't always work. / It can be expensive. / Too many eggs can be stimulated, resulting in unexpected multiple pregnancies. [1 mark]

p.80 - Adrenaline and Thyroxine

- Q1 thyroid gland [1 mark]
- Q2 When the high level of thyroxine is detected, the secretion of TSH from the pitultary gland is inhibited [I mark]. This reduces the amount of thyroxine released from the thyroid gland [I mark], so the level in the blood falls back towards normal If mark].

p.81 - Plant Hormones

Q1 More auxin accumulates on the side of the shoot that's in the shade [1 mark]. This causes the shaded side of the shoot to elongate faster [1 mark] so the shoot bends towards the light.

p.82 — Commercial Uses of Plant Hormones

Q1 E.g. to stimulate seed germination / to control dormancy / to stimulate flowering / to grow larger fruit [1 mark].

p.84 — DNA

- Q1 A small section of DNA found on a chromosome [1 mark] that codes for a particular sequence of amino acids [1 mark] that are put together to make a specific protein [1 mark].
- Q2 The entire set of genetic material in an organism [1 mark].

p.85 — The Structure of DNA and Protein Synthesis

Protein Synthesi Q1a) Four [1 mark].

b) GAGGATGCACTA [1 mark]

p.86 - Mutations

- Q1 A mutation is a random change in an organism's DNA [1 mark].
- Q2 Mutations change the DNA base sequence [1 mark]. This can alter the amino acids that the bases code for [1 mark]. This can change the shape of the protein the amino acids make up, which can affect its function [1 mark].

p.87 — Reproduction

- Q1 mitosis [1 mark]
- Q2 Because there are two parents, the offspring contain a mixture of their parents' genes [1 mark]. This mixture of genetic information produces variation [1 mark].

p.88 — Meiosis

Q1 23 [1 mark]

p.89 — More on Reproduction

Q1 Any three from: e.g. there only needs to be one parent unlike sexual reproduction, which requires two. If takes less energy than sexual reproduction because organisms don't need to find a mate. It is faster than sexual reproduction. / Many identical offspring can be produced in favourable conditions. [1 mark for each correct answer, up to 3 marks]

p.90 - X and Y Chromosomes

Q1 XX [1 mark]

p.91 — Genetic Diagrams

Q1 Your genotype is the combination of alleles you have [1 mark]. Your phenotype is the characteristics you have [1 mark].

p.92 - More Genetic Diagrams

Q1



round peas : wrinkly peas

[1 mark for correct gametes, 1 mark for correct offspring genotypes and 1 mark for correct ratio.]

p.93 — Inherited Disorders

Q1 Because the allele which causes cystic fibrosis is recessive [1 mark], so you have to have two recessive alleles to have the disorder [1 mark]. Heterozygous people have one dominant and one recessive allele [1 mark].

p.94 — The Work of Mendel

Q1 Scientists of the time didn't have the background knowledge to properly understand Mendel's findings because they didn't know about genes, DNA and chromosomes [1 mark].

p.95 - Variation

Q1 Differences between members of the same species [1 mark] that have been caused by the environment/conditions something lives in [1 mark].

p.96 - Evolution

Q1 There was a variety of tongue lengths in the moth population [1 mark]. Moths with longer tongues got more food/nectar and were more likely to survive [1 mark]. These moths were more likely to reproduce and pass on the genes responsible for their long tongues [1 mark]. So, over time, longer tongues became more common in the moth population [1 mark].

p.97 - More About Evolution

Q1 It contradicted the common religious beliefs of the time that all life on earth was created by God [f mark]. Darwin lacked the evidence that he needed to convince many other scientists [f mark]. He also didn't have any explanation for why new beneficial characteristics appeared, or how they were passed on to offspring [f mark].

p.98 - Selective Breeding

- 61 Select rabbits with floppy ears [1 mark] and breed them together to produce offspring [1 mark]. Select offspring with floppy ears and breed them together [1 mark]. Repeat this over many generations until all of the offspring have floppy ears [1 mark].
- Q2 Selective breeding reduces the gene pool [1 mark]. This causes an increased chance of organisms inheriting harmful genetic defects [1 mark]. There is also an increased chance that a population could be wiped out by a new disease [1 mark].

p.99 — Genetic Engineering

Q1 Benefit - E.g. the characteristics chosen for GM crops can mean that they have an increased yield. / GM crops can be engineered to contain certain nutrients, which some people in developing nations may lack from their diets [1 mark]. Concern - E.g. some people are not convinced that GM crops are safe and are concerned that we might not fully understand the effects of eating them on human health. / Some people say that growing GM crops will affect the number of wild flowers and insects that live in and around the crops. / There is concern that the transplanted genes could get out into the natural environment, which could lead to the creation of 'superweeds' [1 mark].

p.100 - Cloning

Q1 The nucleus is removed from an unfertilised egg [I mark] and a complete set of chromosomes from an adult body cell is inserted into the egg [I mark]. The egg cell is then stimulated to divide to from an embryo by being given an electric shock [I mark]. The embryo is the implanted into an adult female [I mark], where it develops into a clone of the original adult body cell [I mark].

p.101 — Fossils

Q1 The microbes that cause decay can't survive in low oxygen conditions [1 mark], so the dead organisms are preserved rather than decayed [1 mark].

p.102 — Speciation

O1 The two populations may be in different environments [1 mark] and so different characteristics will be beneficial [1 mark]. Natural selection will work differently in each population so that individuals with the beneficial characteristics are more likely to survive and reproduce [1 mark].

p.103 - Antibiotic-Resistant Bacteria Q1 Any one from: e.g. when the illness is only

- minor. / When the infection is being caused by a virus [1 mark]. Taking the complete course makes sure
- that all the bacteria are destroyed [1 mark] This means that there are none left to mutate and develop into antibiotic-resistant strains [1 mark].

p.104 — Classification

Q1 B and C [1 mark].

p.106-Competition

- Q1 Any three from: light / space / water / mineral ions [1 mark for each correct answer, up to 3 marks].
- Q2 E.g. the frog population might increase as there might be more water spiders available for them to eat [1 mark] because there will be fewer sticklebacks to eat the water spiders / less competition for food [1 mark]. / The frog population might decrease as they are more likely to be eaten by pike [1 mark] because there will be fewer sticklebacks for the pikes to eat [1 mark]

p. 107 - Abiotic and Biotic Factors

- Q1 Any four from: moisture level / light intensity / temperature / carbon dioxide level / wind intensity / wind direction / soil pH / mineral content of soil [1 mark for each correct answer, up to 4 marks).
- Q2 Any two from: availability of food / competition for resources / new pathogens [1 mark for each correct answer, up to 2 marks].

p.108 - Adaptations

Q1a) A behavioural adaptation [1 mark].

b) E.g. it has flippers [1 mark] so it can swim for food [1 mark]. / A thick layer of fat [1 mark] so it retains heat [1 mark]. / A low surface area to volume ratio [1 mark] so it retains heat [1 mark].

p.109 - Food Chains

Q1a) grass [1 mark]

- b) three [1 mark]
- c) grasshopper [1 mark]
- d) The population of grasshoppers could increase [1 mark] as there's nothing to eat them [1 mark]. The population of snakes could decrease [1 mark] as there's nothing for them to eat [1 mark].

p.110 - Using Quadrats

Q1 $0.75 \times 4 = 3$ buttercups per m² [1 mark]. 3 × 1200 = 3600 buttercups in total [1 mark]

p.111 - Using Transects

- Q1 A line used to help find out how organisms are distributed across an area [1 mark].
- They could mark out a line across the field, from one corner to the other [1 mark]. Then they could count all of the dandelions that touch the line [1 mark]
- You could estimate the percentage of the quadrat covered by the organisms [1 mark]

p.112 - Environmental Change & The Water Cycle

- Q1 Any two from: e.g. a change in the availability of water / a change in temperature / a change in atmospheric gases [1 mark for each correct answer. up to 2 marks].
- Q2a) By evaporation / transpiration [1 mark]. b) By providing them with fresh water [1 mark].

p.113 — The Carbon Cycle

- Microorganisms/detritus feeders break them down/digest them [1 mark].
- By green plants and algae in photosynthesis [1 mark]

p.114 - Decay

Q1 Oxygen availability [1 mark], water availability [1 mark] and the number of microorganisms present [1 mark].

p.115 — Investigating Decay

Q1a) Rate = 1000

Rate = $\frac{1000}{480}$ = 2.08 s⁻¹ [1 mark]

b) As the temperature increases, the rate of decay increases [1 mark].

p.116 - Biodiversity and Waste

Management

The variety of different species of organisms on Earth, or within an ecosystem [1 mark].

p.117 — Global Warming

Global warming causes higher temperatures, which cause ice to melt and seawater to expand [1 mark]. This causes the sea level to rise [1 mark], which could lead to flooding of low-lying land and therefore the loss of habitats [1 mark].

p.118 - Deforestation and Land Use

Q1 Trees 'lock up' some of the carbon that they absorb during photosynthesis, so if lots of trees are removed, less carbon will be locked up from the atmosphere [1 mark]. If the land is cleared by burning the trees, this means that lots of carbon dioxide is released [1 mark]. Microorganisms feeding on leftover dead wood release carbon dioxide as a waste product of respiration [1 mark].

p.119 - Maintaining Ecosystems and Biodiversity


- Hedgerows and field margins can be reintroduced around single-crop fields [1 mark]. These provide a habitat for organisms that would otherwise be unable to live in the area [1 mark].
- Breeding programmes breed endangered animals in captivity to make sure the species survives if they die out in the wild [1 mark]. Individuals can sometimes be released into the wild to boost or re-establish a population [1 mark].

p. 120 - Trophic Levels

Q1 They secrete enzymes that break down the dead material into small soluble food molecules [1 mark].

p.121 - Pyramids of Biomass

Q1 E.a.



[2 marks for drawing all bars correctly to scale, otherwise 1 mark for any three bars drawn correctly to scale. 1 mark for all bars correctly labelled, 1 mark for a correct scale.]

p. 122 — Biomass Transfer

- Q1 Any two from: e.g. organisms don't always eat every single part of the organism they're consuming. / Organisms don't absorb all of the material in the food they ingest. / Some of the biomass is converted into other substances and released as waste. [1 mark for each correct answer, up to 2 marks]
- Q2a) 995 110 = 885 kg [1 mark] b) (110 ÷ 995) × 100 = 11.1% [1 mark]

p.123 - Food Security and Farming

Q1 The movement of livestock can be restricted [1 mark] and they can be kept in a temperature-controlled environment [1 mark]. This means that they use less energy moving around and controlling their own body temperature [1 mark], so more energy is available for growth [1 mark].

p. 124 - Biotechnology

Q1 A plasmid could be removed from the bacterial cell and cut open with an enzyme [1 mark]. The rennin gene could be cut out of a cow chromosome using an enzyme [1 mark]. The plasmid and rennin gene could then be joined together and the plasmid re-inserted into the bacterial cell [1 mark]

Glossary

Abiotic factor	A non-living factor of the environment.
Accommodation	The ability to focus on near or distant objects by changing the shape of the lens in the eye.
Accurate result	A result that is very close to the true answer.
Active transport	The movement of particles against a concentration gradient (i.e. from an area of lower concentration to an area of higher concentration) using energy transferred during respiration.
Adaptation	A feature or characteristic that helps an organism to survive in its natural environment.
Aerobic respiration	Respiration taking place in the presence of oxygen.
Allele	An alternative version of a gene.
Alveolus	A tiny air sac in the lungs, where gas exchange occurs.
Amino acid	A small molecule that is a building block of proteins.
Anaerobic respiration	Respiration taking place in the absence of oxygen.
Anomalous result	A result that doesn't seem to fit with the rest of the data.
Antibiotic	A drug used to kill or prevent the growth of bacteria.
Antibiotic resistance	When bacteria aren't killed by an antibiotic.
Antibody	A protein produced by white blood cells in response to the presence of an antigen.
Antigen	A molecule on the surface of a cell or a pathogen. Foreign antigens trigger white blood cells to produce antibodies.
Antitoxin	A protein produced by white blood cells that counteracts toxins made by invading bacteria.
Artery	A blood vessel that carries blood away from the heart.
Asexual reproduction	Where organisms reproduce by mitosis to produce genetically identical offspring.
Auxin	A plant hormone that controls the growth of a plant in response to different stimuli.
Behavioural adaptation	A way in which an organism behaves that helps it to survive in its environment.
Bias	Prejudice towards or against something.
Binary fission	A type of simple cell division carried out by prokaryotic cells where the cell makes copies of its genetic material, before splitting into two daughter cells.
Binomial system	The system used in classification for naming organisms using a two-part Latin name.
Biodiversity	The variety of different species of organisms on Earth, or within an ecosystem.
Biomass	The mass of living material in an organism or a group of organisms.
Biotic factor	A living factor of the environment.
Capillary	A type of blood vessel involved in the exchange of materials at tissues.
Carbohydrase	A type of digestive enzyme that catalyses the breakdown of a carbohydrate into sugars.
Cardiovascular disease	Disease of the heart or blood vessels.
Catalyst	A substance that increases the speed of a reaction, without being changed or used up.
Categoric data	Data that comes in distinct categories (e.g. flower colour or blood group).
Cell membrane	A membrane surrounding a cell, which holds it together and controls what goes in and out.

Cell wall	A structure surrounding some cell types, which gives strength and support.
Cellulose	A molecule which strengthens cell walls in plants and algae.
Central Nervous System (CNS)	The brain and spinal cord. It's where reflexes and actions are coordinated.
Chlorophyll	A green substance found in chloroplasts which absorbs light for photosynthesis.
Chloroplast	A structure found in plant cells and algae. It is the site of photosynthesis.
Chromosome	A long molecule of DNA found in the nucleus. Each chromosome carries many genes.
Clinical trial	A set of drug tests on human volunteers.
Clone	An organism that is genetically identical to another organism.
Communicable disease	A disease that can spread between individuals.
Community	The populations of different species living in a habitat.
Continuous data	Numerical data that can have any value within a range (e.g. length, volume or temperature).
Contraceptive	A method of preventing pregnancy, which can be hormonal or non-hormonal.
Control experiment	An experiment that's kept under the same conditions as the rest of an investigation, but doesn't have anything done to it.
Control variable	A variable in an experiment that is kept the same.
Coordination centre	An organ (e.g. the brain, spinal cord or pancreas) that processes information from receptors and organises a response from the effectors.
Coronary artery	A blood vessel which supplies blood to the heart muscle.
Coronary heart disease	A disease in which the coronary arteries are narrowed by the build up of fatty deposits.
Correlation	A relationship between two variables.
Cutting (plants)	A small piece of a plant (usually with a new bud on it) that can be grown into a new plant.
Cystic fibrosis	An inherited disorder of the cell membranes caused by a recessive allele.
Cytoplasm	A gel-like substance in a cell where most of the chemical reactions take place.
Deforestation	The cutting down of forests (large areas of trees).
Dependent variable	The variable in an experiment that is measured.
Diabetes	A condition that affects the body's ability to control its blood glucose level.
Dialysis	A way of artificially filtering the blood to remove waste products and keep the concentration of dissolved ions in the blood at normal levels. It is used to treat patients with kidney failure.
Differentiation	The process by which a cell becomes specialised for its job.
Diffusion	The spreading out of particles from an area of higher concentration to an area of lower concentration.
Discrete data	Numerical data that can only take certain values (e.g. number of people).
Distribution	Where organisms are found in a particular area.
DNA	Deoxyribonucleic acid. The molecule in cells that stores genetic information.
Dominant allele	The allele for the characteristic that's shown by an organism if two different alleles are present for that characteristic.
Double-blind trial	A clinical trial where neither the doctors nor the patients know who has received the drug and who has received the placebo until all the results have been gathered.

Glossary

	101
Ecosystem	The interaction of a community of living organisms with the abiotic parts of their environment.
Effector	Either a muscle or gland which responds to nervous impulses.
Efficacy	Whether something, e.g. a drug, works or not.
Endothermic reaction	A reaction where energy is transferred from the environment.
Enzyme	A protein that acts as a biological catalyst.
Eukaryotic cell	A complex cell, such as a plant or animal cell.
Evolution	The changing of the inherited characteristics of a population over time.
Excretion	The removal of waste products from the body.
Exothermic reaction	A reaction that transfers energy to the environment.
Extinction	When no living individuals of a species remain.
Extremophile	An organism that's adapted to live in extreme conditions.
Family tree	A diagram that shows how a characteristic is inherited in a group of related people.
Fermentation	The process of anaerobic respiration in yeast cells.
Fertilisation	The fusion of male and female gametes during sexual reproduction.
Fertility	The ability to conceive a child.
Fishing quota	A limit on the number and size of fish that can be caught in a certain area.
Food security	Having enough food to feed the population.
Fossil	The remains of an organism from many years ago, which is found in rock.
Fossil record	The history of life on Earth preserved as fossils.
Functional adaptation	Something that goes on inside an organism's body that helps it to survive in its environment.
Gamete	A sex cell, e.g. an egg cell or a sperm cell in animals.
Gene	A short section of DNA, found on a chromosome, which contains the instructions needed to make a protein (and so controls the development of a characteristic).
Genetic engineering	The process of cutting out a useful gene from one organism's genome and inserting it into another organism's cell(s).
Genetically modified (GM) crop	A crop which has had its genes modified through genetic engineering.
Genome	All of the genetic material in an organism.
Genotype	What alleles you have, e.g. Tt.
Geotropism	See gravitropism.
Gibberellin	A type of plant hormone that stimulates seed germination, stem growth and flowering.
Gland	An organ that hormones are produced and secreted from.
Global warming	The rise in the average global temperature.
Glucagon	A hormone produced and secreted by the pancreas when blood glucose level is too low.
Glycogen	A molecule that acts as a store of glucose in liver and muscle cells.
Gravitropism	The growth of a plant in response to gravity. Also known as geotropism.
Guard cell	A type of cell found on either side of a stoma. A pair of these cells control the stoma's size.

Habitat	The place where an organism lives.
Haemoglobin	A red pigment found in red blood cells that carries oxygen.
Hazard	Something that has the potential to cause harm (e.g. fire, electricity, etc.).
Heterozygous	Where an organism has two alleles for a particular gene that are different.
Homeostasis	The regulation of conditions inside your body (and cells) to maintain a stable internal environment, in response to changes in both internal and external conditions.
Homozygous	Where an organism has two alleles for a particular gene that are the same.
Hormone	A chemical messenger which travels in the blood to activate target cells.
Hybridoma	A cell made by fusing a mouse lymphocyte with a tumour cell, which is used to produce lots of monoclonal antibodies.
Hyperopia	Long-sightedness.
Hypothesis	A possible explanation for a scientific observation.
Inbreeding	When closely related animals or plants are bred together.
Independent variable	The variable in an experiment that is changed.
Inherited disorder	A disorder caused by a faulty allele, which can be passed on to an individual's offspring.
Insulin	A hormone produced and secreted by the pancreas when blood glucose level is too high.
Interdependence	Where, in a community, each species depends on other species for things such as food, shelter, pollination and seed dispersal.
IVF	In vitro fertilisation. The artificial fertilisation of eggs in the lab.
Limiting factor	A factor which prevents a reaction from going any faster.
Lipase	A type of digestive enzyme that catalyses the breakdown of lipids into fatty acids and glycerol.
Mean (average)	A type of average found by adding up all the data and dividing by the number of values.
Median (average)	The middle value in a set of data when the values are put in order of size.
Meiosis	A type of cell division where a cell divides twice to produce four genetically different gametes. It occurs in the reproductive organs.
Menstrual cycle	A monthly sequence of events during which the body prepares the lining of the uterus (womb) in case it receives a fertilised egg, and releases an egg from an ovary. The uterus lining then breaks down if the egg has not been fertilised.
Meristem tissue	Tissue found at the growing tips of plant shoots and roots that is able to differentiate.
Metabolism	All the chemical reactions that happen in a cell or the body.
Mitochondria	Structures in a cell which are the site of most of the reactions for aerobic respiration.
Mitosis	A type of cell division where a cell reproduces itself by splitting to form two identical offspring.
Mode (average)	The most common value in a set of data.
Monoclonal antibodies	Antibodies produced from lots of clones of a single white blood cell. This means that all the antibodies are identical and will only target one specific antigen.
Motor neurone	A nerve cell that carries electrical impulses from the CNS to effectors.
MRSA	$A strain \ of \ antibiotic-resistant \ bacteria. \ (Meticillin-resistant \ \textit{Staphylococcus aureus.})$
Mutation	A random change in an organism's DNA.
Mycoprotein	Protein produced by fungi, in particular Fusarium.

Glossary

	190
Myopia	Short-sightedness.
Natural selection	The process by which species evolve.
Negative feedback	A mechanism that restores a level back to optimum in a system.
Nervous system	The organ system in animals that allows them to respond to changes in their environment.
Neurone	A nerve cell. Neurones transmit information around the body, including to and from the CNS.
Non-communicable disease	A disease that cannot spread between individuals.
Nucleotide	A repeating unit in DNA and RNA that consists of a sugar, a phosphate group and a base.
Nucleus (of a cell)	A structure found in animal and plant cells which contains the genetic material.
Obesity	A condition where a person has an excessive amount of body fat, to the point where it poses a risk to their health.
Optimum dose	The dose of a drug that is most effective and has few side effects.
Optimum level (in the body)	A level of something (e.g. water, ions or glucose) that enables the body to work at its best.
Organ	A group of different tissues that work together to perform a certain function.
Organ system	A group of organs working together to perform a particular function.
Osmosis	The movement of water molecules across a partially permeable membrane from a region of higher water concentration to a region of lower water concentration.
Oxygen debt	The amount of extra oxygen your body needs after exercise to react with the build up of lactic acid and remove it from cells.
Partially permeable membrane	A membrane with tiny holes in it, which lets some molecules through it but not others.
Pathogen	A microorganism that causes disease, e.g. a bacterium, virus, protist or fungus.
Peer review	The process where scientists check each other's results and explanations to make sure they're scientific.
Permanent vacuole	A structure in plant cells that contains cell sap.
Phagocytosis	The process by which white blood cells engulf foreign cells and digest them.
Phenotype	The characteristics you have, e.g. brown eyes.
Phloem	A type of plant tissue which transports dissolved sugars around the plant.
Photosynthesis	The process by which plants use energy to convert carbon dioxide and water into glucose and oxygen.
Phototropism	The growth of a plant in response to light.
Placebo	A substance that is like a drug being tested, but which doesn't do anything.
Plasma	The liquid component of blood, which carries blood cells and other substances around the body.
Platelet	A small fragment of a cell found in the blood, which helps blood to clot at a wound.
Polydactyly	An inherited disorder, caused by a dominant allele, where a person has extra fingers or toes.
Precise result	A result that is close to the mean.
Predator	An animal that hunts and kills other animals for food.
Prey	An animal that is hunted and killed by another animal for food.
Primary consumer	An organism in a food chain that feeds on a producer.

Producer	An organism at the start of a food chain that makes its own food using energy from the Sun.
Prokaryotic cell	A small, simple cell, e.g. a bacterium.
Protease	A type of digestive enzyme that catalyses the breakdown of proteins into amino acids.
Protein	A large biological molecule made up of long chains of amino acids.
Protist	A type of pathogen. Protists are often transferred to other organisms by a vector.
Punnett square	A type of genetic diagram.
Pyramid of biomass	A diagram to represent the biomass at each stage of a food chain.
Quadrat	A square frame enclosing a known area. It is used to study the distribution of organisms.
Random error	A difference in the results of an experiment caused by things like human error in measuring.
Range	The difference between the smallest and largest values in a set of data.
Receptor	A group of cells that are sensitive to a stimulus (e.g. receptor cells in the eye detect light).
Recessive allele	An allele whose characteristic only appears in an organism if there are two copies present.
Reflex	A fast, automatic response to a stimulus.
Relay neurone	A nerve cell that carries electrical impulses from sensory neurones to motor neurones.
Repeatable result	A result that will come out the same if the experiment is repeated by the same person using the same method and equipment.
Reproducible result	A result that will come out the same if someone different does the experiment, or a slightly different method or piece of equipment is used.
Respiration	The process of breaking down glucose to transfer energy, which occurs in every cell.
Ribosome	A structure in a cell, where proteins are made.
Risk	The chance that a hazard will cause harm.
Risk factor	Something that is linked to an increased likelihood that a person will develop a certain disease.
Secondary consumer	An organism in a food chain that eats a primary consumer.
Selective breeding (artificial selection)	When humans artificially select the plants or animals that are going to breed, so that the genes for particular characteristics remain in the population.
Sensory neurone	A nerve cell that carries electrical impulses from a receptor in a sense organ to the CNS.
Sex chromosome (humans)	One of the 23rd pair of chromosomes, X or Y. Together they determine whether an individual is male or female.
Sexual reproduction	Where two gametes combine at fertilisation to produce a genetically different new individual.
Speciation	The development of a new species.
Species	A group of similar organisms that can reproduce to give fertile offspring.
Stable community	A community where all the species and environmental factors are in balance, so that the population sizes are roughly constant.
Statins	A group of medicinal drugs that are used to decrease the risk of heart and circulatory disease.
Stem cell	An undifferentiated cell which has the ability to become one of many different types of cell, or to produce more stem cells.
Stent	A wire mesh tube that's inserted inside an artery to help keep it open.
Stimulus	A change in the environment.

Stoma	A tiny hole in the surface of a leaf.
Structural adaptation	A feature of an organism's body structure that helps it to survive in its environment.
Synapse	The connection between two neurones.
Systematic error	An error that is consistently made every time throughout an experiment.
Tertiary consumer	An organism in a food chain that eats a secondary consumer.
Theory	A hypothesis which has been accepted by the scientific community because there is good evidence to back it up.
Thermoregulatory centre	An area of the brain which controls and monitors body temperature.
Tissue	A group of similar cells that work together to carry out a particular function.
Tissue culture (plants)	A method of cloning plants in which a few plant cells are put on a growth medium containing hormones and allowed to grow into new plants.
Toxicity	How harmful something is, e.g. a drug.
Toxin	A poison. Toxins are often produced by bacteria.
Transect	A line which can be used to study the distribution of organisms across an area.
Translocation	The movement of dissolved sugars around a plant.
Transpiration stream	The movement of water from a plant's roots, through the xylem and out of the leaves.
Trophic level	A stage in a food chain.
Tumour	A growth of abnormal cells.
Uncertainty	The amount of error your results might have.
Urea	A waste product produced from the breakdown of amino acids in the liver.
Vaccination	The injection of dead or inactive microorganisms, in order to produce an immune response that will help to protect you against a particular pathogen in the future.
Valid result	A result that is repeatable, reproducible and answers the original question.
Valve	A structure within the heart or a vein which prevents blood flowing in the wrong direction.
Variable	A factor in an investigation that can change or be changed (e.g. temperature or concentration).
Variation	The differences that exist between individuals.
Vasoconstriction	When blood vessels supplying the skin capillaries constrict (get narrower).
Vasodilation	When blood vessels supplying the skin capillaries dilate (get wider).
Vector (in genetic engineering)	Something used to transfer DNA into a cell, e.g. a virus or a bacterial plasmid.
Vector (in disease)	An organism that transfers a disease from one animal or plant to another, which doesn't get the disease itself.
Vein	A blood vessel that carries blood to the heart.
Virus	A tiny pathogen that can only replicate within host body cells.
White blood cell	A blood cell that is part of the immune system, defending the body against disease.
Xylem	A type of plant tissue which transports water and mineral ions around the plant.
Zero error	A type of systematic error caused by using a piece of equipment that isn't zeroed properly.

Index

condensation 112

condoms 48, 78

contact lenses 71

contraception 78

convex lenses 71

correlation 7, 9

corneas 70

control variables 4

coronary arteries 34

coronary heart disease 37

abiotic factors 107 abstinence 78 accommodation 70 accuracy (of results) 5 active sites 28 active transport 22 adaptations 108 adrenal glands 73, 80 adrenaline 80 adult cell cloning 100 aerobic respiration 62 alleles 91 alveoli 24, 33 ammonia 75 amylase 29-31 anaerobic respiration 62 ancestors 104 animal cells 11 anomalous results 5 antibiotic resistance 51, 103 antibiotics 51, 103 investigating the effects of 17 antibodies 36, 49

anti-diuretic hormone (ADH) antimicrobials 49 antiretroviral drugs 47 antitoxins 36, 49 aorta 34 apparatus 126 Archaea 104 arteries 35 artificial blood 38 artificial hearts 38 asexual reproduction 87, 89 aspirin 51 atria 34 auxins 81, 82

В bacteria 11, 16, 17, 46, 48, 104 balances 126 bar charts 6 barrier methods (of contraception) 78 basal metabolic rate 80 behavioural adaptations 108 Benedict's reagent 32 benign tumours 41 bias 2 bile 30 binary fission 16 binomial system 104 biodiversity 116-119 maintenance of 119

biogas 114 biological heart valves 38 biomass 122 biomass transfer 122 biotechnology 124 biotic factors 107 Biuret test 32 blind studies 52 body temperature 72 brain 69 breathing rate 33 bronchi 33 bronchioles 33 Bunsen burners 127

caffeine (effect on reaction time) 68 cancer 54 capillaries 35 carbohydrases 30 carbon cycle 113 catalysts 28 categoric data 6 cell cycle 15 division 15, 16 elongation 81 membranes 11, 20 walls 11 cells 11 drawing 13 specialised 14 central nervous system (CNS) 66, 67 cerebellum 69 cerebral cortex 69 chlorophyll 11, 57 chloroplasts 11, 57 cholesterol 37 chromosomes 15, 84 X and Y 90 cilia 49 ciliary muscles 70 class (classification) 104 classification 104 climate change 117

clinical trials 52

communicable diseases

complementary base pairing

computational models 1

concentration gradients 20

39. 46

communities 106

concave lenses 71

conclusions 9

cloning 100

cover slip (slides) 13 cuttings 82, 100 cystic fibrosis 93 cytoplasm 11 D Darwin, Charles 96, 97 decay 113-115 investigating 115 rate of 114 decomposers 120 defences (of plants) 55 deforestation 118 deletions (mutations) 86 dependent variables 4 detritus feeders 113 diabetes 74 dialysis 76 diaphragm (thorax) 33 diaphragms (contraceptive) 78 differentiation 14, 27 diffusion 20, 21, 23-25 digestive enzymes 30 digestive system 27, 31 digitalis 51 discrete data 6 distribution of organisms 110-112 DNA 15, 84, 85 nucleotides 85 dormancy (in seeds) 82 dosage 52 double-blind studies 52 double circulatory systems 33 drugs 51, 52 development of 52 Е ecosystems 106 maintenance of 119 effectors 65, 66 efficacy (of drugs) 52 electric heaters 127 embryo screening 93

electron microscopes 12 embryo transplants 100

endocrine system 73 endothermic reactions 57 environmental changes 112 environmental variation 95 enzymes 28-31 epidermal tissue 42 errors 5 ethene (in plants) 82 ethics (in investigations) 127 Eukaryota 104 eukaryotes 11, 46 evaluations 10 evaporation 112 evolutionary trees 104 evolution by natural selection 96, 97, 102

exchange surfaces 23-25 exercise 63 exothermic reactions 61 extinction 96 eyes 70, 71

fair tests 4 family (classification) 104 family trees 92 farming 123 fatty acids 30 fermentation 62 fertility 78, 79 fertility treatments 79 fight or flight response 80 filtration (kidneys) 75 Fleming, Alexander 51 flowering 82 focusing 70 follicle-stimulating hormone (FSH) 77, 79 food chains 109, 120, 121 food poisoning 48 food production 123 food security 123 fossils 101

gall bladder 31 gamete fusion 88 gametes 87, 88 gas exchange 24, 25 gas syringes 126 gene pools 98, 100 genes 15, 84 genetically modified crops 99

functional adaptations 108

fungi 46, 47

Index

genetic diagrams 90-93 disorders 93 engineering 99, 124 variants 86 variation 95 genomes 84 genotypes 91 genus 104 geotropism 81 germination 82 gibberellin 82 gills 25 glands 73 global warming 117 glucagon 74 glucose in the blood 74 test for 32 uses in plants 57 glycerol 30 glycogen 74 gonorrhoea 48 gradients 7 graphs 7 gravitropism 81 greenhouse gases 117 greenhouses 60 guard cells 44 gullet 31

н

habitats 106
haemoglobin 36
hazards 3
heart 34
heart valves 38
hereditary units 94
heterozygous organisms 91
HIV 47
homeostasis 65
homozygous organisms 91
hormones 73
hybridoma cells 53
hyperopia 71
hypotheses 1, 4

Ī

immune system 49 independent variables 4 indicators 126 'induced fit' model 28 infertility 79 inhibition zones 17, 18 insertions (mutations) 86 insulin 74 intercepts (on a graph) 7

interdependence 106 intrauterine devices (IUDs) 78 inverse correlations 7 proportion 60 square law 60 in vitro fertilisation (IVF) 79 iodine test (for starch) 29 irises 70 iris reflex 70 IVF 79

\mathbf{K}

kidney failure 76 kidneys 75, 76 kingdom (classification) 104

lactic acid 62, 63

т.

Lamarck, Jean-Baptiste 97 lamellae 25 large intestine 31 laser eye surgery 71 leaves gas exchange 25 tissues in 25 lenses 70, 71 light microscopes 12, 13 limiting factors (of photosynthesis) 57-59 line graphs 7 lines of best fit 7 Linnaean system 104 Linnaeus, Carl 104 lipases 30 lipids (test for) 32 liver 31, 74 'lock and key' model 28 long-sightedness 71 lungs 33 luteinising hormone (LH) 77, 79 lymphocytes 53

M

magnetic resonance imaging (MRI) 69
magnification 12
malaria 47
malignant tumours 41
mean (average) 6
mean division time 16
measles 47
measuring cylinders 126
mechanical heart valves 38
median 6

medulla (brain) 69 meiosis 88 Mendel, Gregor 94 menstrual cycle 77 meristem tissue 42 metabolism 61 microorganisms 46 culturing 17, 18 microscopes 12, 13 migration patterns 117 human 84 mitochondria 11 mitosis 15, 88 mode 6 models (scientific) 1 monoclonal antibodies 53, 54 motor neurones 66, 67 MRI scanning 69 mRNA 85 MRSA 51, 103 mucus 49 muscle cells 14 muscle fatigue 63 mutations 86, 95 mycoprotein 124 myopia 71

TAT

natural selection 96, 102 negative correlations 7 negative feedback systems 65 blood glucose 74 body temperature 72 thyroxine 80 water content 75 nerve cells 14 nervous system 66, 67 neurones 66, 67 neuroscientists 69 non-coding DNA 85, 86 non-communicable diseases 39 nucleotides 85 nucleus 11

O

oestrogen 77, 78
optic nerve 70
order (classification) 104
organ donors 76
organs 27
organ systems 27
osmosis 21
ovaries 73, 77
overfishing 123
ovulation 77
oxygen debt 63

pacemakers 34 painkillers 51 palisade mesophyll tissue 42 pancreas 31, 73, 74 parasites 46 partially permeable membranes 21 pathogens 46 pea plants 94 peat bogs 118 peer review 1, 52 pepsin 30, 31 percentage change 130 cover 111 percentiles 130 Petri dishes 17 phagocytosis 36, 49 phenolphthalein 115

phenotype variation 96 phloem cells 14 phloem tubes 42, 43 pH meters 126 photosynthesis 57-59 rate of 58, 59 phototropism 81 phylum (classification) 104 pipettes 126 pituitary gland 73, 77, 80 placebo effect 52 plant cells 11

phenotypes 91

diseases 55

growth 81, 82 hormones 81, 82 plasma 36 plasmids 11, 99 platelets 36 pleural membranes 33 pollution 116 polydactyly 93 population (human) 116 positive correlations 7 potantes 21 potometers 44, 128 preclinical testing 52 precipitation 112 precision (of results) 5

predator-prey cycles 109

primary consumers 109

progesterone 77, 78

prokarvotes 11, 16

predators 109

producers 109

prev 109

pregnancy tests 53

proteases 30

Index

proteins 85 function of 85 test for 32 protists 46, 47 puberty 77 pulmonary artery 34 pulmonary vein 34 Punnett squares 90, 91 pupils 70 pyramids of biomass 121

Q quadrats 110, 111

R random errors 5 sampling 129 range (of data) 6 rates of reaction 29 reaction times 68 receptors 65, 66 rectum 31 red blood cells 36 reflex arcs 67 relay neurones 67 repeatability (of data) 4, 5 replacement lens surgery 71 representational models 1 reproducibility (of data) 4, 5 resolution (microscopes) 12 respiration 61, 62 retinas 70 ribcage 33 ribosomes 11, 85 ripening (of fruit) 82 risk 3 risk factors 40 root hair cells 14, 22

S safety 3, 127 salivary glands 31 Salmonella 48 sample size 4 sampling 129 sclera 70 secondary consumers 109 secondary sexual

characteristics 77

rooting powder 82

rose black spot 47

rounding numbers 6

roots 81

selective breeding 98 selective reabsorption 75 sensory neurones 66, 67 sex hormones 77-79 sexually transmitted diseases sexual reproduction 87, 89 shivering 72 shoots 81 short-sightedness 71 SI units 8 significant figures 6 slides 13 small intestine 31 specialised cells 14 speciation 96, 102 species 102, 104 in classification 102 sperm cells 14, 77 spermicide 78 spinal cord 69 spongy mesophyll tissue 42 stains (microscope) 13 standard form 12 starch 29 test for 32 statins 37 stem cells 19 stents 37 sterilisation (contraception) stomach 31 stomata 25, 44 structural adaptations 108 subcellular structures 11, 14 substitutions (mutations) 86 Sudan III 32 sugars (test for) 32 superbugs 103 surface area to volume ratios survival of the fittest 96 suspensory ligaments 70

synapses 67 systematic errors 5 tables (of data) 6

sweat 72, 75

target organs 73 temperature receptors 72 tertiary consumers 109 testes 73, 77 testosterone 77 theories 1 thermometers 126 thermoregulatory centre 72

thorax 33 three-domain system of classification 104 thyroid gland 73, 80 thyroid stimulating hormone (TSH) 80 thyroxine 80 tissue culture 82, 100 tissues 27 tobacco mosaic virus 47 toxicity (of drugs) 52 trachea 33 transects 111 transpiration 43, 44, 112 rate 44 transplants (organ) 76 trophic levels 120-122 tumour markers 54 tumours 41 Type 1 diabetes 74 Type 2 diabetes 74

uncertainty 10 units 8 urea 75 urine 75

vaccination 48, 50 vacuoles 11 valid results 4 valves (heart) 34, 38 variables 4 variation 95 vasoconstriction 72 vasodilation 72 vectors 46, 48, 99 veins 35 vena cava 34 ventricles 34 villi 24 viruses 46, 47 vision defects 71 volumes (measuring) 126

w

Wallace, Alfred Russel 102 waste management 116 water baths 127 water cycle 112 weed killers 82 white blood cells 36, 49 wizardry 126 Woese, Carl 104

X and Y chromosomes 90

xvlem cells 14 xylem tubes 42, 43

yeast cells 62

zero errors 5



CGP books — they might just save your life...



OK, maybe not your Me. But when it comes to exam revision, CGP are the undisputed champions. You can order any of our books and cards (with next-day delivery!) from us, online or by phone:

www.cgpbooks.co.uk • 0800 1712 712

Or you'll find our range in any good bookshop, including:





Waterstones

Inspiring 9-1 GCSE Biology revision from CGP!

Gasp — this CGP Biology book is incredible! Take a deep breath and dive in...

- Brilliant notes for the Higher Level AQA course...
 Of course every topic and required practical!
- Stacks of exam-style questions...
 Yep, plus dozens of step-by-step video solutions

Trust CGP to warm up your cerebral cortex — it's the brainiest revision around \odot

P.S. CGP's matching Exam Practice Workbook is incredibly helpful too — don't miss it!



Filled to the brim with knowledgey goodness

